

Talon®-G

RODENTICIDE BAIT PACK (PELLETS)

For Effective Control of Commensal Rats

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI Americas Inc. 302/575-3000.

IF BAIT IS EATEN BY ANIMALS OR PETS, call University of Illinois Animal Poison Control Center 217/333-3611 or ICI Americas Inc. 302/575-3000 for 24-hour emergency assistance.

In case of significant spill, call CHEMTREC 800/424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated as in poisoning from overdose of bis(hydroxy coumarin). For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions given on the label. This warranty does not extend to the use of this product contrary to label instructions or under abnormal use conditions or under conditions not reasonably foreseeable to Seller and Buyer and no assurances are given by any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR MANUFACTURE OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE RELIEF SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Kills warfarin-resistant Norway Rats.
Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

Active Ingredient:

Brodilacoum:
3-(3'-4'-bromo-[1,1'-biphenyl]-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-
hydroxy-2H-1-benzopyran-2-one

Inert Ingredients	0.005%
Total	100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

May Be Harmful or Fatal if Swallowed.
Read Additional Precautionary Statements
on Side Panel




ICI Americas Inc.

Wilmington, Delaware 19897

EPA Reg No 10182-39
EPA Est No 2393 WI-1
Net Contents 4/100 x 50 gm (1.76 oz) bait pack boxes

Batch Number



STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container except for holding additional TALON rodenticide.

Storage: Store in original container only in a dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Cashier Disposal: Dispose of bait container in a sanitary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway Rats and Roof rats in and around homes, industrial, commercial, agricultural and public buildings. TALON may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Authorized by the USDA for use in federally inspected meat, poultry and egg processing plants.

SELECTION OF TREATMENT AREAS: Determine areas where rats will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floor and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS

Norway and Roof Rats: Apply 2 to 8 bait packs (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease. **Rats/Mice:** Open pack and apply $\frac{1}{4}$ to $\frac{1}{2}$ ounce of bait at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. **Rats and Mice:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS—CAUTION MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING. BAITS IF BAITS IS EATEN BY HUMANS, CALL A PHYSICIANS AT ONCE. For 24-hour emergency assistance call ICI America Inc. 302/575-3000 if BAITS IS EATEN BY ANIMALS OR PETS, CALL University of Illinois Animal Poison Control Center 217/333-3611 or ICI America Inc. 302/575-3000 for 24-hour emergency assistance. In case of significant spill, call CHEMTREC 800/424-9300.

NOTE TO PHYSICIANS OR PET OWNERS This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in preparing from overdoses of dietary coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks. (based on monitoring of prothrombin times) In severe cases, blood transfusions may be necessary.

EXTREME HAZARD HAZARDS This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

STORAGE AND DISPOSAL Store away from children and pets. Wrap empty or used container and put in trash.

Talon-G

RODENTICIDE BAIT PACK (PELLETS)
For Effective Control of Commercial Rats

Kills waterfowl-resistant Norway Rats.
Rodents may consume a lethal dose in one feeding
with live dead rodents appearing four or five days
after treatment begins.

ACTIVE INGREDIENT:

Brodifacoum,
1,3,4-(Epoxy-1,1-butenyl)-4-yl-
1,3,4-(Epoxy-1,1-butenyl)-1-naphthalenyl-4-
Hydroxy-2H-1-benzopyran-2-one

INERT INGREDIENTS

Total 100.000%

99.995%

KEEP OUT OF REACH OF CHILDREN

CAUTION

May Be Harmful or Fatal If Swallowed

Read Additional Precautionary Statements on Side Panel

ICI ICI Americas Inc.
Agricultural Chemicals Division
Wilmington, Delaware 19897

EPA Reg No 10102-39
EPA Est No 2393 WI-1
Net Contents 50 gons (1.76 oz.)
Batch Number

ACO 790-230R3 B/F

DIRECTIONS FOR USE It is a violation of federal law to use this product in a manner inconsistent with its labeling. See Instructions for control of Norway Rats and Roof rats in and around homes, industrial, commercial, agricultural and public buildings. TALON may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Authorized by the USDA for use in federally inspected meat, poultry and egg processing plants. Select areas of frequent activity. Determine areas where rats will most likely find and consume the bait. Generally, these areas are along walls, by grain piles, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS Norway and Roof Bait. Apply 2 to 8 bait packs (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease. Bait Bait. Open pack and apply 1/4 to 1/2 ounce of baits at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. Bait and Bait. Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to Disposal paragraph. To prevent reinfestation eliminate food, water, and harborage as much as possible. If reinfestation does occur repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

5.026157/C6