

17 FEB 1988

395
17
Mr. Miller
8'

Ms. Diane L. Ierley
ICI Americas, Inc.
Concord Pike & New Murphy Road
Wilmington, DE 19897

Dear Ms. Ierley:

Subject: Talon-G Rodenticide Bait Pack (Pellets)
EPA Registration No. 10182-39
Re: Labels Submitted in Response to the Agency's
Inert Policy
Your Letter Dated October 19, 1987

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable subject to the comment listed below.

- The "inerts of concern" statement must be in a type size larger than, or equal to, the heading used in the ingredients statement.

Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of this condition.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

William H. Miller
Product Manager (16)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

10744-I:Palmateer:CBI-19:KENCO:2/1/88:4/23/88:CB:LF:VO:EK:CB
CONCURRENCES

SYMBOL	ORIGINATOR						
.....						
SURNAME						
DATE						

Talon®-G

RODENTICIDE BAIT PACK (PELLETS)

For Effective Control of Commensal Rats

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI Americas Inc. 302/575-3000.

IF BAIT IS EATEN BY ANIMALS OR PETS, call University of Illinois Animal Poison Control Center 217/333-3611 or ICI Americas Inc. 302/575-3000 for 24-hour emergency assistance.

In case of significant spill, call CHEMTREC 800/424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer and user assumes the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Kills warfarin-resistant Norway Rats.

Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

Active ingredient:

Brodifacoum-

3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one 0.005%

Inert ingredients 99.995%

This product contains the toxic inert ingredient Rhodamine B. 100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

May Be Harmful or Fatal if Swallowed. Read Additional Precautionary Statements on Side Panel.



ICI Americas Inc.

Wilmington, Delaware 19897

EPA Reg. No. 13182-39

EPA Est. No. 2393 WI-1

Net Contents: 4/100 x 50 gm (1.76 oz) bait pack boxes

Batch Number.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage in container except for holding additional bait. Storage: Store in original container only and pets.

Pesticide Disposal: Wastes resulting from use should be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of bait container if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in violation of the labeling.

USE RESTRICTIONS: For control of Norway Rats in homes, industrial, commercial, agricultural, and around transport vehicles and terminal buildings. Do not use in sewers, storm drains, or in locations not accessible to rodents. Do not place bait in areas where it may be consumed by wildlife. Do not place bait in areas where it may be consumed by humans or surfaces that come in direct contact with humans.

Authorized by the USDA for use in feed processing plants.

SELECTION OF TREATMENT AREAS: Do not place bait in areas where it may be consumed by humans or surfaces that come in direct contact with humans. Generally, place bait in openings, in or beside burrows, in corners of walls, or in locations where rodents or other animals are likely to find much food as possible.

APPLICATION DIRECTIONS

Norway and Roof Rats: Apply 2 to 8 bait pack per placement. Maintain an uninterrupted bait station until signs of rat activity cease. Repeat baiting at intervals of 8 to 12 feet per placement. Baiting is needed at points of very high mouse activity. Refresh bait for 15 days or until signs of mouse activity are eliminated. Contaminated or spoiled bait immediately remove and unconsumed bait according to "Disposal" instructions. Eliminate food, water, and harborage as soon as possible. Where a continuous baiting program is required, repeat treatment. Where a continuous baiting program is required, repeat treatment. Where a continuous baiting program is required, repeat treatment. Where a continuous baiting program is required, repeat treatment.

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS—CAUTION MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT. IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI Americas Inc. 302/375-3000. IF BAIT IS EATEN BY ANIMALS OR PETS, CALL University of Illinois Animal Poison Control Center 717/233-3611 or ICI Americas Inc. 302/375-3000 for 24-hour emergency assistance. In case of significant spill, call CHEMTREC 800/424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN: This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of brodifacoum. For human cases, vitamin K₁ is indicated at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is indicated at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin time) or until normal blood coagulation is restored, if necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

STORAGE AND DISPOSAL: Store away from children and pets. Wrap empty or used container and put in trash.

Talon-G
RODENTICIDE BAIT (PELLETS)
For Effective Control of Commercial Rats

Kills warren-resistant Norway Rats,
Rodent-resistant house mice and
wild mice. Kills rats and mice
within 24 hours of eating bait. Kills
after treatment 30 days.

ACTIVE INGREDIENT:

Brodifacoum
3-(2,4-dinitrophenyl)-1,1-diphenyl-4-
1,2,3,4-tetrahydro-1-naphthalenyl-4-
hydroxy-2H-1-benzopyran-2-one

INERT INGREDIENTS:

Other ingredients constitute the balance.

0.005%
99.995%
100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Read Additional Precautionary Statements on Side Panel

ICI
ICI Americas Inc.
Agricultural Chemicals Division
Wilmington, Delaware 19857

EPA Reg. No. 10182-39
EPA Est. No. 2283 WA-1
Net Contents 50 gms (1.76 oz)
Batch Number

ACD 790-23683 B/F

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Use as directed. For control of Norway Rats and Roof rats in and around homes, industrial, commercial, agricultural and public buildings. TALON may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Authorized by the USDA for use in federally inspected meat, poultry and egg processing plants. Available in Tampered Areas: Determine areas where rats and mice likely find and consume the bait. Generally, these areas are along walls, by ground openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS: Review and Use Rate: Apply 2 to 8 bait packs (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease. Break into 1/2 ounce of baits at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. Bait and floor. Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and uneaten bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

ACCEPTED

10182-39

10182-39

5.07667% CO.