

PM16

REG# 10182-40

17 FEB 1988

315
17
206,343
1

Ms. Diane L. Ierley
ICI Americas, Inc.
Concord Pike & New Murphy Road
Wilmington, DE 19897

Dear Ms. Ierley:

Subject: Talon-G Rodenticide Bait Pack (Mini-Pellets)
EPA Registration No. 10182-40
Re: Labels Submitted in Response to the Agency's
Inert Policy
Your Letter Dated October 19, 1987

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable subject to the comment listed below.

- The "inerts of concern" statement must be in a type size larger than, or equal to, the heading used in the ingredients statement.

Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of this condition.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

William H. Miller
Product Manager (16)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

10744;I:Palmateer:CBT-19:KENCO:2/11/88:4/23/88:CB:LF:VO:EK:CB

CONCURRENCES

SYMBOL	ORIGINATOR						
SURNAME							
DATE							

OFFICIAL FILE COPY

ACCEPTED
with COMMENT
in EPA Letter

17 FEB

Under the Federal
Fungicide, and
as amended, for
registered under E

Talon-G

RODENTICIDE BAIT PACK (MINI-PELLETS)

For Effective Control of Commensal Mice and Rats

**PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI Americas Inc. Medical Emergency Information Center 1-800-327-8633 IF BAIT IS EATEN BY ANIMALS OR PETS, CALL ICI Americas Inc. Medical Information Center 1-800-327-8633 for 24-hour emergency assistance. In case of significant spill, call CHEMTREC 800/424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of bis-hydroxy coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Kills warfarin-resistant Norway Rats and house mice
Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.
Active ingredient:
Brodifacoum-
3-[3-(4'-bromo-[3,1'-biphenyl] 4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one
Inert ingredients
This product contains the toxic inert ingredient Rhodamine B.

	0.005%
	99.995%
	100.000%

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

May Be Harmful or Fatal if Swallowed
Read Additional Precautionary Statements
on Side Panel

This product is protected by U.S. patent numbers 3957824 and 4035505 which are the property of Ward Blenkinsop and Company Limited



ICI Americas Inc.
Wilmington, Delaware 19897

EPA Reg No 10182-40
EPA Est No 2393 WI-1
Net Contents 4/200 x 25gm (0.88oz) bait pack boxes

STORAGE AND LABELING

Do not contaminate water, food or feed by empty container except for holding additional Storage: Store in original container only in a and Pets.
Pesticide Disposal: Wastes resulting from disposed of on site or at an approved waste Container Disposal: Dispose of bait container incineration if allowed by state and local

DIRECTIONS FOR USE

It is a violation of Federal law to use this product labeling

USE RESTRICTIONS: For control of Norway Rats, mice, and house mice in homes, industrial, commercial, agricultural and be used in and around transport vehicles (Shipping terminal buildings. Do not use in sewers. This proof bait boxes or in locations not accessible to wildlife. Do not place bait in areas where there are or surfaces that come in direct contact with

Authorized by the USDA for use in federal processing plants

SELECTION OF TREATMENT AREAS: Determine and consume the bait. Generally, these openings, in or beside burrows, in corners and walls, or in locations where rodents or their much food as possible

APPLICATION DIRECTIONS

Norway and Roof Rats:
Apply 4 to 16 bait packs (usually at intervals of 8 to 12 feet) needed at points of very high mouse activity. Refresh bait for 15 days or until signs of mouse activity cease

House Mice:
Apply 1 bait pack at intervals of 8 to 12 feet needed at points of very high mouse activity. Refresh bait for 15 days or until signs of mouse activity cease

Replace contaminated or spoiled bait with dead animals and unconsumed bait according to reinfestation, eliminate food, water, and reinfestation does occur, repeat treatment. If reinfestation is present, establish permanent needed