

Talon

RODENTICIDE MINI-PELLETS

For Effective Indoor Control of Commensal Mice and Rats

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT.
IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE.
For 24-hour emergency assistance call 302-575-3000.
In case of significant spill, call CHEMTREC 800/424-9300.

This product may reduce the clotting ability of the blood and cause hemorrhaging. If this occurs, intravenous and oral administration of vitamin K₁ combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Vitamin K₁ is antidotal at doses of 10 to 20 mg.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

STORAGE: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

DISPOSAL: Bait that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternate procedures such as limited open burning.

Kills warfarin-resistant Norway Rats.
Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.
Active Ingredient:
Brodifacoum
3-[3-(4'-bromo[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4-hydroxy-2H-1-benzopyran-2-one
Inert Ingredients ...

Total

0.008%
99.992%
100.000%

May Be Harmful or Fatal if Swallowed
Read Additional Precautionary Statements on Side Panel



ICI Americas Inc.

Wilmington, Delaware 19897

EPA Reg. No. 10182-41
EPA Est. No. 523-IL-1
Net Contents: 4/5 kg (11.025 lb) boxes

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice in homes, industrial buildings and commercial buildings. Talon may also be used inside transport vehicles (ships, trains, aircraft) and inside sealed port or terminal buildings. For indoor use only. Treated baits must be placed in tamper proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Authorized by the USDA Food Safety and Quality Service for use in Federally inspected meat, poultry and egg processing plants.

SELECTION OF TREATMENT AREAS: After removing as much food, water and harborage as possible, determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed.

Apply 4 to 16 ounces of bait usually at intervals of 15 to 30 feet per placement. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of rat activity cease.

Apply 4 to 16 ounces of bait at intervals of 6 to 10 feet per placement. Larger placements up to 20 ounces may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. Repeat treatment when infestation recurs. Where continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.