

P-16 PLG# 10182-48

10182-48

10182-48

WeatherBlok® BAIT

Made with TALON® Rodenticide

Kills warfarin-resistant Norway Rats and House mice.
Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

Active ingredient:

Brodifacoum	
3-[3-(4-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one	0.005%
Inert ingredients	99.995%
Total	100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

May Be Harmful or Fatal if Swallowed. Read Additional Precautionary Statements on Side Panel.



This product is protected by U.S. patent numbers 3957824 and 4035505 which are the property of Ward Blenkinsop and Company Limited

EPA Reg No 10182-48 EPA Est No 30564-EN-1
Net Contents 500/20 gm bait blocks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container except for holding additional WEATHER BLOK bait.

STORAGE: Store in original container only in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Dispose of bait container in a sanitary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats, Roof rats and House mice in and around sewers, homes, industrial, commercial, agricultural and public buildings. WEATHER BLOK may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. WEATHER BLOK must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife such as in burrows which are then closed. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Authorized by the USDA for use in federally inspected meat, poultry, and egg processing plants.

APPLICATION DIRECTIONS

Norway and Roof Rats: Apply 8-23 WEATHER BLOKS (usually at intervals of 15-30 ft) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease. When infestations are limited to overhead areas in or around buildings, tie or fasten WEATHER BLOK securely in protected locations on fences, poles, beams or other overhead areas where signs indicate rodents are active.

For use in sewers, securely attach one end of a wire to WEATHER BLOK and the other end to a stationary structure such as the bottom step of a manhole ladder or to a sewer grate, allowing just enough wire for the block to rest on manhole benching. If benching is not present, suspend block a few inches above the high water line or place WEATHER BLOK on a board supported by opposing steps of the ladder. Securing WEATHER BLOK in this manner will prevent removal by rats or water. Place at least 15 WEATHER BLOKS per manhole. Maintain an uninterrupted supply of fresh bait for at least 10 days or until feeding stops.

House Mice: Apply 1 WEATHER BLOK at intervals of 8-12 feet per placement. Larger placements (2 Blocks) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. For use in sewers apply according to above directions for rats.

Made in U.K. ACD 790-279R1 F/F

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance, call ICI American Inc. 302/575-3000.

IF BAIT IS EATEN BY ANIMALS OR PETS, call the National Animal Poison Control Center 217/333-3611 or ICI American Inc. 302/575-3000 for 24-hour emergency assistance.

In case of significant spill, call CHEMTREC 800/424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of bishydroxy coumatrin. For human cases, vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer and User assumes the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

ACCEPTED

AUG 27 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10182-48

INK SELECTION
Pms 185

