



KILLS RATS AND MICE

RODENTICIDE BAIT PACK (PELLETS)

HAVOC

APPLICATION DIRECTIONS

Empty out bait bins. Apply 2 to 4 packs (amount of 15 to 30 bait per placement). Maintain an uninterrupted supply of fresh bait for 14 days or until signs of rat activity cease.

When Bait Shows Fresh and signs: 1) to 1% number of baits at intervals of 4 to 12 feet per placement. Larger placements (up to 2 corners) may be needed at points of very high rodent activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

Bait and Bait Station accompanied by baited bait receptacle. Contact and dispose of all bait receptacles and bait stations according to "Disposal" paragraph. To prevent contamination, always wash, rinse, and thoroughly dry bait stations. Do not reuse bait receptacle. When a continuous source of information is present, maintain permanent bait stations and approach bait as needed.

SOURCE OF BAIT AND USE

Bait stations may be used in accordance with directions on the label and to the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer or - Buyer assumes the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR DAMAGE OR OTHER SPECIAL OR PUNITIVE DAMAGES OR DAMAGES FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

This product is protected by US patent numbers 3957824 and 4025246 which are the property of ICI Americas Inc. and Company Limited. HAVOC is a registered trademark of ICI Americas Inc.

EPA Reg. No. 10162-01
EPA Est. No. 2182-00-1
Net Content: 1.65 lb (0.75 kg) net; 100 bait packs

ICI Americas Inc.
Agricultural Chemicals Division
Wilmington, Delaware 19807

ICI 706 00503 10

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty containers except for holding additional HAVOC rodenticide. Storage: Store in a special container only in a dry place inaccessible to children and pets.

Hazardous Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Disposal: Dispose of bait container as a secondary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the following:

Use Restrictions: For control of Norway Rats, Roof rats and House mice in and around homes, industrial, commercial, agricultural and public buildings. HAVOC may also be used in and around transient vehicles (trucks, buses, campers) and related parts of temporary buildings. Authorized by the USDA for use in Federally inspected meat, poultry and egg processing plants. Do not use in sewers. This product must be placed in traps, great bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.


Locations of baited areas: Determine areas where rats or mice will eat and store bait and consume the bait. Generally these areas are along walls, by ground openings or in hidden harbors in corners and unoccupied places between floors and walls or in crevices where rodents or their signs have been observed. Baiting on such bait as the label

10 bait packs

HAVOC

RODENTICIDE BAIT PACK (PELLETS)

KILLS RATS AND MICE



Kills warfarin-resistant Norway Rats and House Mice.

Rodents may consume a lethal dose in one feeding or first dead rodents appearing four or five days after treatment begins.

KEEP OUT OF REACH OF CHILDREN

CAUTION

May Be Harmful or Fatal if Swallowed. Read Additional Precautionary Statements on Side Panel.

Active Ingredients: Bromethalin
21.2% (w/w) + 0.000001% (w/w) = 21.2% (w/w)
Total: 21.2% (w/w)
Inert Ingredients: 78.8% (w/w)
Net Content: 1.65 lb (0.75 kg)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION
MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF BAIT IS EATEN BY HUMANS CALL A PHYSICIAN AT ONCE For 24-hour emergency information, call ICI Americas Medical Emergency Information-1-800-327-8633.

IF BAIT IS EATEN BY ANIMALS OR PETS, CALL VET IMMEDIATELY. Medical Information Center: 1-800-327-8633 for 24-hour emergency assistance.

In case of significant spill, call CHEM TALK 1-800-424-9300.

NOTE: TO PREVENT RATS OR MICE FROM BEING RE-BAITED.

This product may reduce the carrying ability of the food and cause malnutrition if rodents are not adequately fed or if rodents are not fed. Do not use in conjunction with other rodenticides or other pest control products. For human consumption, it is prohibited to consume of 10 to 20 mg (0.35 mg/kg) per pound (0.45 kg) of body weight. For other uses, it is prohibited to consume of 1 to 5 mg/kg. Personnel should wear a protective mask and gloves when handling bait. Personnel should wear a protective mask and gloves when handling bait. Personnel should wear a protective mask and gloves when handling bait.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

COPY 01
BEST AVAILABLE COPY

ACCEPTED
AUG 30 1983
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended (for US EPA Reg. No. 10162-01)
10162-01

PA 116 10182 61
-1-17



PRECAUTIONARY STATEMENTS. HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT.
IF BAIT IS EATEN BY HUMAN, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI American Inc. Medical Emergency Information Center 1-800-327-8633. **IF BAIT IS EATEN BY ANIMAL OR PET, CALL ICI American Inc. Medical Information Center 1-800-327-8633 for 24-hour emergency assistance or NOTE TO PHYSICIAN OR VETERINARIAN:** This product may reduce the clotting ability of the blood and cause hemorrhaging if poisoning occurs, and intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of heparin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin time). In severe cases, blood transfusion may be necessary.
ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

STORAGE AND DISPOSAL: Store away from children and pets. Wrap empty or used container and put in trash.

HOYOC

RODENTICIDE BAIT PACK (PELLETS)

KILLS RATS & MICE

Kills warfarin-resistant Norway Rats and House Mice. Rodenticidal contents lethal dose in one feeding with 10-15 pellets. No secondary poisoning five days after treatment begins.

Active Ingredients:
 Brodifacoum 33.3% (bore 1.1' (bophny) & n)
 1,2,3-triazolopyridin-4-yl-1-hydroxy-2-yl-benzoyl-2-yl
 Inert Ingredients 66.7%
 99.995%
 100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

May Be Harmful or Fatal if Swallowed. Read Additional Precautionary Statements on Back Panel.
 EPA Reg No 10182-61
 EPA Est No 2393 WI-1
 Net Contents 50 gm (1.76 oz) bait pack
 Batch Number



ICI Americas Inc.
 Agricultural Products
 Wilmington Delaware 19897

ACD 785-09284 J/G

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling and restrictions. For control of Norway Rats, Roof rats and House mice in and around homes, industrial, commercial, agricultural and public buildings. MAYOC may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Authorized by the USDA for use in Federally inspected meat, poultry, and egg processing plants. Do not use in sewers. This product must be placed in tamper proof bait baures or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. **Selective of Treatment Areas:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings in or beside burrows in corners and concealed places between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS Norway and Roof Rats: Apply 2 to 8 bait packs (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease. **Roof Rats:** Open pack and apply 1/4 to 1/2 ounce of bait at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. **Bats & Mice:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation eliminate food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present establish permanent bait stations and replenish bait as needed.

ACCEPTED
 AUG 30 1989
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended (FIFRA)
 Pesticide registered under
 EPA Reg. No. 10182-61



2