

Reg # 10182-61

PM-14

191

RECEIVED FROM
DATE MAY 9 1991

Precautionary statements. Hazards to humans and domestic animals—Caution. May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait. If bait is eaten by humans, call a physician at once. For 24-hour emergency assistance, call your local Poison Control Center. If bait is eaten by animals or pets, call your local Poison Control Center for 24-hour emergency assistance.
Usage to avoid hazards or restrictions. This product may reduce the clotting ability of the blood and cause hemorrhaging if poisoning occurs. Intracerebral and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of baitydracoumatin. For human cases, vitamin K₁ is indicated at doses of 10-20 mg (100 mg/kg). For animal cases, vitamin K₁ is indicated at 2.5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin time) in severe cases. Blood transfusions may be necessary.
Environmental hazards. This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals kept out of any body of water.

Storage and disposal: Store away from children and pets. Wrap empty or used container and put in trash.
Distributed by:
SALVAY Animal Health, Inc.
Mendota Heights, MN 55120-1135 USA
EPA Reg. No. 10182-61-116
EPA Est. No. 61382-WI-1



Salsbury™ Ropax™ Rodenticide Bait Pack (Pellets)

**Kills rats & mice,
including warfarin-
resistant Norway rats
and House mice.**

Directions for use: Use in a location of frequent use to use this product in a manner consistent with its labeling. Use restrictions: For control of Norway rats, Roof rats and House mice in and around homes, industrial, commercial, agricultural and public buildings. Salsbury Ropax may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Substances of treatment areas: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by ground openings, in or below basements, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.
Application directions: Norway and Roof rats: Apply 2 to 8 bait packs (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease. House mice: Open pack and apply 4 to 1/2 ounce of bait at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. Signs & notes: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replace bait as needed.

099001

NOV 1991