PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

AVOID CONTACT WITH SKIN AND EYES
AVOID BREATHING SPRAY MIST

STATEMENT OF PRACTICAL TREATMENT

ENVIRONMENTAL HAZARDS

PHYSICAL OR CHEMICAL HAZARDS

STORAGE AND DISPOSAL

NOTICE TO BUYER AND USER

EMULSIFIABLE CONCENTRATE

Keep Out of Reach of Children

CAUTION

See Side Panel For Additional Precautionary Statements and Statement of Practical Treatment

ACTIVE INGREDIENT

Permethrin-
(3-Phenoxyphenyl)methyl (±)cis, trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate 5.7%

INERT INGREDIENTS 94.3%

..... 100.0%

Contains 60 grams per liter (1.67 lb. active ingredient per quart)

ICI Americas Inc.

Agricultural Chemicals Division — Wilmington, Delaware 19897

EPA Reg. No. 10182-63
EPA Est. No. 4581-DE-1

Net Contents
ONE QUART

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.
Apply ECTIBAN according to the following table:

Target Species	Method of Application	Dilute	Application Rate
LACTATING AND NON-LACTATING DAIRY CATTLE AND GOATS, BEEF CATTLE, HORSES AND SHEEP			
Horn Flies, Face Flies, Stable Flies, House Flies, Horse Flies, Black Flies, Mosquitoes, Eye Gnats, Mange, Scabies, Ticks, Lice and Sheep Keds	Sprayer	1 quart to 25 gallons water (2 lbs ga)	1-2 quarts of spray per animal over whole body surface. For mange, lice and scabies be sure to thoroughly soak animal and apply a second treatment at 14 days
Horn Flies, Lice	Sprayer	1 quart to 50 ga water (1 lbs ga)	Spray until entire animal is thoroughly covered. For lice a second treatment at 14 days is recommended
Horn Flies, Face Flies, Stable Flies, Ear Ticks	Low Pressure Sprayer	1 quart to 2 ½ gal water (6 lbs gal)	1-2 oz spray per animal. Spot treat back, face, legs and ears
Horn Flies, Face Flies, Stable Flies, Ear Ticks	Backrubber Self Oiler	1 quart to 10 gal diesel oil (6 lbs ga)	Keep rubbing device charged. Results improved by daily forced use
POULTRY AND SWINE			
Lice, Northern Fowl Mites, Mange	Sprayer	1 quart to 25 gallons water (2 lbs ga)	POULTRY: 1 gal spray per 100 birds paying particular attention to vent. SWINE: Thoroughly soak including ears. Repeat at 14 days. Spray ears and change bedding. For maintenance prevention of non-clinical hog mange ECTIBAN spray concentration may be halved (1 quart to 50 gallons water)
Extended residual activity between treatment and slaughter:			
PREMISES – Dairies, Barns, Feedlots, Stables, Poultry Houses, Swine and Livestock Housing			
House Flies, Stable Flies, Other Manure Breeding Flies, Ants, Cockroaches, Crickets, Silverfish, Mosquitoes, Mites, Ticks	Sprayer	1 quart to 12½ gal water (5 lbs gal)	Spray to point of run-off. As a guide use 1 gal spray per 750 sq ft (400-1400 sq ft)
	Mist Sprayer	Ready to Use	Direct 4 oz onto 1000 sq ft of surface
	Ultra Low Volume Sprayer	1 quart to 12½ gal water (5 lbs gal)	4 oz spray per 1000 cu ft of airspace
EXTERIOR BUILDING STRUCTURES			
	Mist Sprayer	Ready to Use	1 can (5 oz) per 1000 sq ft of surface, concentrating on eaves, doorways and attics

SPECIMEN

BEST DOCUMENT AVAILABLE

10182-63