

RESTRICTED USE PESTICIDE.

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABEL FOR USE ON LETTUCECYMBUSH® 3E Insecticide
EPA Reg. No: 10182-65**PRECAUTIONARY STATEMENTS** — (Refer to container label for additional precautionary statements)

When mixing/loading and spraying, wear mid-forearm to elbow length chemical-resistant gloves, boots, or overshoes, one-piece overalls which have long sleeves or long pants, and face shield or goggles. Mixers/loaders should in addition wear a chemical-resistant apron.

Any article of clothing worn while applying this product must be cleaned before reusing. Clothing which has been drenched or heavily contaminated should be disposed of in accordance with state or local regulations.

Instead of clothing and equipment specified above, the applicator can use an enclosed tractor cab with filtered air supply.

DIRECTIONS FOR USE**It is a violation of federal law to use this product
in a manner inconsistent with its labeling**

Apply CYMBUSH® 3E Insecticide as shown in the following chart. Timing and frequency of applications should be based on insect populations reaching economic thresholds.

RE-ENTRY STATEMENTS: Do not treat areas while unprotected humans or domestic animals are present in the treatment area. Do not allow entry into treated areas without protective clothing until sprays have dried. State re-entry regulations will apply if they are more restrictive than those stated on this label. Consult with State regulatory officials.**DIRECTIONS FOR APPLICATION**

Crop	Target Pests	Rate (lbs a.i./A)	Remarks
Lettuce	Cabbage Looper	0.025 - 0.10	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment or 5 gallons per acre by air.
	Alfalfa Looper	(1.07 - 4.27 fl. oz./A)	
	Black Cutworm	fl. oz./A	
	Granulate Cutworm		
	Tobacco Budworm	0.04 - 0.10	Lower rates of CYMBUSH should be used under light to moderate insect pressure. Higher rates should be used to control heavy insect populations.
	Corn Earworm	(1.71 - 4.27 fl. oz./A)	
	Beet Armyworm	fl. oz./A	

Do not apply more than 1.5 lb a.i./A per season. Do not apply within 5 days of harvest. Do not plant rotational crops within 30 days of the last application.

All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

**This labeling must be in the possession of the user
at the time of pesticide application**

Not for sale or distribution after December 31, 1988. Not for use after December 31, 1989.