

1/23/85
PMS
10182-61

FUSILADE™ 4E Herbicide

Witchgrass added to pages 6 and 12.

JAN 23 1985
10182-61

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS
WARNING**

Causes eye irritation. Avoid contact with skin or clothing. Do not get in eyes, on skin or on clothing. Wear protective clothing and PVC gloves when spraying. Wear protective clothing, PVC gloves and apron and full face shield when handling or mixing concentrate. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT:

IF SWALLOWED: Do not induce vomiting. Call a physician immediately.
IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

IF ON SKIN: Immediately remove contaminated clothing and wash thoroughly with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIANS: Inducing vomiting as first aid for this substance may result in increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the herbicide solvent. Vomiting should be induced only under professional supervision. For 24 hour emergency information call (202) 575-3000.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply when weather conditions favor drift from target area.

This product is toxic to fish.
In case of a significant spill, call CHEMTREC (800) 424-9300.

PHYSICAL OR CHEMICAL HAZARDS: Do not use if there are holes or open flame.

STORAGE AND DISPOSAL

PRECAUTIONS: Do not contaminate water, food or feed by storage or disposal. Open burning is prohibited. Do not reuse empty container.

STORAGE: Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, wash up with sand, earth or synthetic absorbent. Remove to chemical waste area.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of pesticide products, spray machine or residue is a violation of Federal law. If these wastes need be disposed of by you according to label instructions, contact your State Pesticide or Environmental Control Agency or the hazardous waste representative of the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse for reusables. Then offer for recycling or reconditioning or purchase and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the described description on the label and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions or under conditions not reasonably foreseeable to Seller and Buyer and does not assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES OR SPECIAL DAMAGES IN ANY EVENT. THE USE OF THIS PRODUCT IS LIMITED TO THE USES AND CONDITIONS SPECIFICALLY LISTED ON THE LABEL AND THE USES AND CONDITIONS SPECIFICALLY LISTED ON THE LABEL. THE USE OF THIS PRODUCT FOR ANY OTHER PURPOSE IS AT THE USER'S SOLE RISK AND WITHOUT WARRANTY FROM THE SELLER.

ACD 761-3981 A/D



**POSTEMERGENCE HERBICIDE FOR CONTROL
OF PERENNIAL AND ANNUAL GRASS WEEDS**

ACTIVE INGREDIENT:	
butyl 2-(4-chlorophenyl)-2-propyl-4-pyridylphosphorylpropanoate	46.8%
INERT INGREDIENTS	53.2%
TOTAL	100.0%

*As of 10/1/81

FUSILADE 4E contains 4 pounds active ingredient per gallon.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the label. Refer to Supplemental Label for directions for use. This labeling must be in the possession of the user at the time of pesticide application.

**KEEP
OF RE
OF CHIL**

WARN

See Side Panel For
Precautionary State
Statement of Preca

EPA Est. No. 50
EPA Reg. No. 10

Net Content:
ONE QU

Batch No.:



FUSILADE 4E is a selective postemergence herbicide for control of annual and perennial grass weeds. FUSILADE 4E will provide effective control of grass weeds in conventional tillage, minimum tillage, and no-till plantings. FUSILADE 4E does not control broadleaf weeds or sedges (nutgrass).

CONTROL SYMPTOMS

FUSILADE 4E is a systemic herbicide which moves from the treated foliage into the shoots, roots, rhizomes, stolons, and growing points (meristematic regions) of treated grass weeds. Excellent control of a wide range of annual and perennial grass weeds will be obtained with FUSILADE 4E when applied as recommended on this label.

Growth of treated grass weeds stops soon after application. Symptoms include loss of vigor, yellowing and/or reddening, and eventual death to the treated grass plant. Symptoms are generally observed within one to three weeks, depending on grass weed species and environmental conditions.

APPLICATION DIRECTIONS

Timing

Apply to actively growing grass weeds before they exceed the recommended growth stages shown on this label.

Do not apply FUSILADE 4E if rainfall is expected within 1 hour.

Spray Additives (Cleared for use on growing crops)

ALWAYS ADD ONE OF THE FOLLOWING:

- Crop Oil Concentrate

Add a nonphytotoxic crop oil or a once refined vegetable oil concentrate containing 15-20% approved emulsifier; at 1% (1 quart per 25 gallons) of the finished spray volume.

- Nonionic Surfactant

Add nonionic surfactant containing at least 75% surface active agent, at 0.25% ($\frac{1}{4}$ pint per 25 gallons) of the finished spray volume.

Ground Application

Use a minimum of 5 gallons of spray volume per acre and 30-60 psi pressure. A sufficient spray volume and pressure should be used to provide complete coverage of the treated grass weed species.

Use higher volumes and pressure if the grass foliage is dense to ensure complete coverage.

DO NOT USE FLOOD TYPE OR OTHER SPRAY
NOZZLE TIPS WHICH DELIVER COARSE, LARGE DROPLET SPRAYS

DO NOT APPLY FUSILADE 4E WITH RECIRCULATING SPRAYERS, ROPEWICK APPLICATORS, CONTROLLED DROPLET APPLICATORS (CDA) OR ANY SIMILAR DEVICES.

Band Applications

Band application equipment should be adjusted to provide maximum coverage of grasses in the planted row. Thorough coverage can be obtained with a nozzle directed to each side of the row toward the grass weeds in the planted row. Application with a single nozzle over the top of the row is not recommended.

The use of band applications on perennial grasses may result in reinfestation of the treated band from the untreated middle and is not recommended.

Aerial Application

Use a minimum of 5 gallons per acre.

SPOT TREATMENTS

Mix FUSILADE 4E and crop oil concentrate or nonionic surfactant with water according to the amounts shown below. Spray to obtain thorough coverage, but do not spray to runoff. Retreat if necessary.

In cotton, applications must be made prior to boll set. Do not apply a total of more than 1 quart of FUSILADE 4E per acre per season.

In soybeans, applications must be made prior to bloom. Do not apply a total of more than 1 pint of FUSILADE 4E per acre per season.

SPOT SPRAY RECOMMENDATIONS

<u>To Make This Spray Volume</u>	<u>Add These Amounts</u>		
	<u>FUSILADE 4E</u>	<u>Crop/Vegetable Oil Concentrate</u>	<u>or</u> <u>Nonionic Surfactant</u>
1 gallon	1/2 fl. oz. (1 tbsp.)	1-1/4 fl. oz.	or 1/3 fl. oz.
25 gallons	3/4 pint	1 quart	1/2 pint
50 gallons	1-1/2 pints	2 quarts	1 pint
100 gallons	1-1/2 quarts	1 gallon	1 quart

5315

Cultivation

Cultivation no sooner than 3 days after application will often assist in grass weed control. Cultivation of treated grasses 7 days prior to or within 3 days after application of FUSILADE 4E may reduce weed control.

DIRECTIONS FOR USE ON COTTON AND SOYBEANS

ANNUAL AND PERENNIAL GRASS WEED CONTROL
COTTON AND SOYBEANS
REGION A

- AREAS WITH ADEQUATE RAINFALL
- FOR ALL STATES EXCEPT THE IRRIGATED AREAS OF ARIZONA, CALIFORNIA, NEW MEXICO, WEST OKLAHOMA, AND WEST TEXAS
- APPLY FUSILADE 4E TO ACTIVELY GROWING GRASSES
- APPLY FUSILADE 4E AT THE RECOMMENDED RATE TO GRASSES AT THE RECOMMENDED GROWTH STAGES AS OUTLINED
- DO NOT APPLY FUSILADE 4E TO GRASSES WHICH ARE STRESSED DUE TO MOISTURE, TEMPERATURE, OR LOW SOIL FERTILITY; GRASSES WHICH HAVE TILLERED, FORMED SEED HEADS, OR EXCEED RECOMMENDED GROWTH STAGES
- APPLY FUSILADE 4E WHEN THE FIRST GRASS WEED SPECIES IN A MIXED GRASS WEED POPULATION REACHES THE RECOMMENDED GROWTH STAGE FOR TREATMENT

ANNUAL AND PERENNIAL GRASS CONTROL RECOMMENDATIONS FOR REGION A

Annual Grass Species	Height (inches) Rate Pints/A	Remarks
		<ul style="list-style-type: none"> • When grass populations include grasses from all groups below, use the 1/2 pt/A rate of FUSILADE 4E for broad-spectrum control.

- Retreat at the recommended rate if needed to control late-germinating grasses.

Group I

Shattercane/Wildcane/Volunteer milo (<u>Sorghum bicolor</u>)	6-12	1/4
Volunteer corn (<u>Zea mays</u>)	12-18	1/4
Wild Proso Millet (<u>Panicum miliaceum</u>)	4-8	1/4

Group II

Seedling johnsongrass (<u>Sorghum halepense</u>)	2-4	3/8
Goosegrass (<u>Eleusine indica</u>)	2-4	3/8
Witchgrass (<u>Panicum capillare</u>)	2-4	3/8

Group III

Barnyardgrass (<u>Echinochloa crusgalli</u>)	2-4	1/2	
Broadleaf signalgrass (<u>Brachiaria platyphylla</u>)	2-4	1/2	
Fall panicum (<u>Panicum dichotomiflorum</u>)	2-4	1/2	
Itchgrass (<u>Rottboellia exaltata</u>)	2-4	1/2	
Junglerice (<u>Echinochloa colonum</u>)	2-4	1/2	
Large crabgrass (<u>Digitaria sanguinalis</u>)	2-4	1/2	
Smooth crabgrass (<u>Digitaria ischaemum</u>)	2-4 ^c	1/2	Length of runners
Southern sandbur (<u>Cenchrus echinatus</u>)	2-4	1/2	
Southwestern cupgrass (<u>Eriochloa gracilis</u>)	2-4	1/2	
Texas panicum (<u>Panicum texanum</u>)	2-4	1/2	
Wild oats (<u>Avena fatua</u>)	2-4	1/2	
Yellow foxtail (<u>Setaria lutescens</u>)	2-4	1/2	
Green foxtail (<u>Setaria viridis</u>)	2-4	1/2	
Giant foxtail (<u>Setaria faberi</u>)	4-6 ^d	1/2	
Red rice (<u>Oryza sativa</u>)	1-2 ^d	1/2	2-3 leaves
Volunteer Cereals			
Barley (<u>Hordeum vulgare</u>)	2-4	1/2	
Oats (<u>Avena sativa</u>)	2-4	1/2	
Rye (<u>Secale cereale</u>)	2-4	1/2	
Wheat (<u>Triticum aestivum</u>)	2-4	1/2	

Under conditions where rainfall is adequate following FUSILADE 4E application, soil residual activity may be observed. This may result in control or suppression of new flushes of annual grasses. The amount of residual control is dependent on the FUSILADE 4E rate used, grass density, and the soil type, with more activity being seen on coarse-textured soils low in organic matter.

Perennial Grass Species	Height (inches)	Rate (Pints/A)	Remarks
Rhizome johnsongrass (<u>Sorghum halepense</u>)	12-18	1/2	<ul style="list-style-type: none"> • Best results can be obtained if rhizomes or stolons are cut up by preplant tillage practices (discing, plowing etc.) • Two applications of the recommended rate of FUSILADE 4E are generally more effective for full season control than a single application of the same total amount. <p>Apply before the boot stage of growth. If new shoots emerge or regrowth occurs, make a second application of 3/8 pt/A when the johnsongrass is 6-12 inches tall.</p>
Bermudagrass (<u>Cynodon dactylon</u>)	3 or 6-12 inch runner (stolon) length	1/2	<p>If regrowth occurs, retreat at recommended height and rate. Bermudagrass is a low-growing perennial grass, and the crop canopy may shield it from herbicide sprays. Improve herbicide spray coverage by directing the spray under the crop canopy.</p>
Quackgrass (<u>Agropyron repens</u>)	less than 10 minimum of 4 leaves	1/2	<p>If regrowth occurs, retreat at the recommended rate when 3-5 quackgrass leaves are present.</p>
Wirestem muhly (<u>Muhlenbergia frondosa</u>)	4-12	1/2	<p>If regrowth occurs, retreat at the recommended rate two to six weeks after the first application.</p>

ANNUAL AND PERENNIAL GRASS WEED CONTROL
COTTON AND SOYBEANS
REGION B

- FOR THE IRRIGATED AREAS OF ARIZONA, CALIFORNIA, NEW MEXICO, WEST OKLAHOMA, AND WEST TEXAS
- APPLY FUSILADE 4E TO ACTIVELY GROWING GRASSES
- FUSILADE 4E MUST BE APPLIED WITHIN 7 DAYS AFTER IRRIGATION
- APPLY FUSILADE 4E AT THE RECOMMENDED RATE TO GRASSES AT THE RECOMMENDED GROWTH STAGES AS OUTLINED IN THE TABLE
- DO NOT APPLY FUSILADE 4E TO GRASSES WHICH ARE STRESSED DUE TO MOISTURE, TEMPERATURE, OR LOW SOIL FERTILITY: GRASSES WHICH HAVE TILLERED, FORMED SEED HEADS, OR EXCEED RECOMMENDED GROWTH STAGES
- APPLY FUSILADE 4E WHEN THE FIRST GRASS WEED SPECIES IN A MIXED GRASS WEED POPULATION REACHES THE RECOMMENDED GROWTH STAGE FOR TREATMENT

ANNUAL AND PERENNIAL GRASS CONTROL RECOMMENDATIONS
FOR REGION B

Annual Grass Species	Height (inches)	Rate (Pints/A)	Remarks
Group I			
Seedling johnsongrass (<u>Sorghum halepense</u>)	2-4	1/2	
Volunteer corn (<u>Zea mays</u>)*	12-24	1/2	• In cotton, retreat at the recommended rate if needed to control late germinating grasses or regrowth.
Group II			
Texas panicum (<u>Panicum texanum</u>)*	2-4	3/4	
Volunteer cereals	2-4	3/4	
Barley (<u>Hordeum vulgare</u>)			
Milo (<u>Sorghum bicolor</u>)			
Oats (<u>Avena sativa</u>)			
Wheat (<u>Triticum aestivum</u>)			

Group III

Barnyardgrass (<u>Echinochloa crus-galli</u>)	2-4	1
Large crabgrass (<u>Digitaria sanguinalis</u>)*	2-4	1
Yellow foxtail (<u>Setaria lutescens</u>)*	2-4	1
Field sandbur (<u>Cenchrus pauciflorus</u>)*	2-4	1
Junglerice (<u>Echinochloa colonum</u>)*	2-4	1

• In soybeans, do not apply more than one pint per acre per season.

Perennial Grass Species

Rhizome johnsongrass (<u>Sorghum halepense</u>)	12-18	3/4 to 1
Bermudagrass (<u>Cynodon dactylon</u>)	3 or 4-8 runner (stolon) length	3/4 to 1

• Best results can be obtained if rhizomes and stolons are cut up by preplant tillage practices (discing, plowing, etc.)

Apply before the boot stage. In cotton, a second application is necessary for season long control. Retreat at 12-18 inches.

For Arizona and California, 1 pint per acre is recommended. In cotton, retreat at recommended height and rate for season long control.

*Not recommended for control of this grass in Arizona and California.

SOYBEAN TANK-MIX FUSILADE 4E HERBICIDE/BLAZER 2L HERBICIDE
Refer to Supplemental Label for directions for use.

LABELED CROPS

Remarks and Precautions

COTTON

Do not apply a total of more than 1 quart of FUSILADE 4E per acre per season to cotton.

Do not harvest cotton within 100 days of application.

Do not apply after boll set.

Do not graze or harvest for forage or hay.

Do not plant rotational grass crops such as corn, sorghum and cereals within 60 days after last application.

SOYBEANS

Do not apply a total of more than 1 pint of FUSILADE 4E per acre per season to soybeans.

Make last application before bloom.

Do not graze or harvest for forage or hay.

Do not plant rotational grass crops such as corn, sorghum and cereals within 60 days after last application.

DIRECTIONS FOR USE NONCROP/NONFOOD USES

Not for use or storage in or around the home.

NONCROP AREAS

FUSILADE 4E herbicide can be used to control grass weeds in areas such as airports, electric transformer stations and substations, pipeline pumping stations, around buildings, storage yards, fence lines, parkways, roadsides, utility rights-of-way, and other noncropland areas.

NOTE: IF SPRAYING AREAS ADJACENT TO ORNAMENTAL TREES, SHRUBS OR GROUND COVERS:

DO NOT APPLY TO FOLIAGE OF ORNAMENTAL TREES, SHRUBS OR GROUND COVERS UNLESS THEY ARE LISTED IN TABLE I OF THIS ORNAMENTAL SECTION OF THIS LABEL AND A NONIONIC SURFACTANT IS BEING USED.

FALLOW LAND

FUSILADE 4E herbicide can be used to control grass weeds during fallow periods.

Do not plant rotational crops other than cotton and soybeans within 60 days after last application.

NONBEARING TREE CROPS/VINES/ASPARAGUS/JOJOBA

FUSILADE 4E herbicide can be used to control grass weeds in nonbearing groves, orchards or vineyards or for site preparation prior to transplanting.

APPLY AS A DIRECTED SPRAY TO PREVENT CONTACT OF SPRAY WITH FOLIAGE OF TREES OR VINES.

DO NOT APPLY TO TREES OR VINES THAT WILL BE HARVESTED WITHIN ONE YEAR AFTER APPLICATION.

DO NOT APPLY TO ASPARAGUS FROM WHICH SPEARS WILL BE HARVESTED WITHIN ONE YEAR OF TREATMENT.

FUSILADE 4E is recommended on the following nonbearing crops:

- | | |
|------------|--------------|
| Almonds | Macadamia |
| Apples | Nectarines |
| Apricots | Olives |
| Asparagus | Oranges |
| Avocados | Peaches |
| Berries | Pears |
| Cherries | Pecans |
| Dates | Pistachios |
| Figs | Plums |
| Filberts | Pomegranates |
| Grapes | Prunes |
| Grapefruit | Tangelos |
| Jjoba | Tangerines |
| Lemons | Walnuts |
| Limes | |

ORNAMENTALS

FUSILADE 4E herbicide can be used to control grass weeds in many newly transplanted and established nongrassy ornamentals, trees, shrubs and ground covers. See the following Tables I and II for specific recommended uses. (USE ONLY NONIONIC SURFACTANT ON ORNAMENTALS.)

Application of FUSILADE 4E may be made over-the-top to ornamentals listed in Table I. Apply as a directed spray to prevent contact of spray with foliage of ornamentals listed in Table II.

APPLICATION DIRECTIONS

Timing

Apply to actively growing grass weeds before they exceed the recommended growth stages shown on this label.

Do not apply FUSILADE 4E if rainfall is expected within 1 hour.

Spray Additives (Cleared for use on growing crops)

ALWAYS ADD ONE OF THE FOLLOWING:

- Crop Oil Concentrate

Add a nonphytotoxic crop oil or a once refined vegetable oil concentrate containing 15-20% emulsifier; at 1% (1 quart per 25 gallons) of the finished spray volume.

Do not use when making applications to ornamentals.

• Nonionic Surfactant

Add nonionic surfactant containing at least 75% surface active agent, at 0.25% ($\frac{1}{4}$ pint per 25 gallons) of the finished spray volume.

A nonionic surfactant is recommended for use when treating grass infested ornamentals.

ANNUAL AND PERENNIAL GRASS CONTROL RECOMMENDATIONS FOR NONCROP/NONFOOD USES

Use 1 pint of FUSILADE 4E per acre and 1% crop oil concentrate (1 quart per 25 gallons) or .25% ($\frac{1}{4}$ pint per 25 gallons) of a nonionic surfactant. Use only nonionic surfactant on ornamentals. Spray to obtain thorough coverage of grass foliage, but not to runoff. Repeat treatments may be necessary to provide full-season control. Use standard pesticide hollow cone or flat fan nozzles.

Apply FUSILADE 4E to actively growing grasses before they exceed the recommended growth stages shown below:

Apply 1.0 pt/A of FUSILADE 4E herbicide.

Grass

Perennial Grasses

Growth Stage

Johnsongrass (<u>Sorghum halepense</u>)	12-18" tall and before boot stage
Bermudagrass (<u>Cynodon dactylon</u>)	3" tall or with 4-8" runners
Quackgrass (<u>Agropyron repens</u>)	3-5 leaves, not more than 10" tall
Guineagrass (<u>Panicum maximum</u>)	6-12" tall and before seedhead initiation
Wirestem muhly (<u>Muhlenbergia frondosa</u>)	4-12" tall and before seedhead initiation

Annual Grasses

Growth Stage 2-8" tall, before tillering and/or heading

- Barnyardgrass (Echinochloa crusgalli)
- Broadleaf signalgrass (Brachiaria platyphylla)
- Fall panicum (Panicum dichotomiflorum)
- Field sandbur (Cenchrus pauciflorus)
- Foxtail, green (Setaria viridis)
- Foxtail, giant (Setaria faberi)
- Foxtail, yellow (Setaria lutescens)

128. L

Goosegrass (Eleusine indica)
Guineagrass, seedling (Panicum maximum)
Itchgrass (Rottboellia exaltata)
Johnsongrass, seedling (Sorghum halepense)
Junglerice (Echinochloa colonum)
Large crabgrass (Digitaria sanguinalis)
Prairie cupgrass (Eriochloa contracta)
Red rice (Oryza sativa)
Shattercane/Wildcane/Volunteer milo (Sorghum bicolor)
Smooth crabgrass (Digitaria ischaemum)
Southern sandbur (Cenchrus echinatus)
Southwestern cupgrass (Eriochloa gracilis)
Texas panicum (Panicum texanum)
Volunteer barley (Hordeum vulgare)
Volunteer oats (Avena fatua)
Volunteer rye (Secale cereale)
Volunteer corn (Zea mays)
Volunteer wheat (Triticum aestivum)
Wild oats (Avena fatua)
Wild proso millet (Panicum miliaceum)
Witchgrass (Panicum capillare)

TABLE I

OVER-THE-TOP APPLICATIONS may be applied to the following ornamentals:

Use Only Nonionic Surfactant on Ornamentals

Alyssum (Alyssum spp.)
Arborvitae (Thuja spp.)
Cinquefoil (Potentilla spp.)
Cotoneaster (Cotoneaster spp.)
Dogwood (Cornus spp.)
Douglas fir¹/Pseudotsuga spp.)
English ivy (Hedera helix)
Euonymus (Euonymus spp.)
Fir¹/Abies spp.)
Forsythia (Forsythia spp.)
Hemlock¹/Tsuga spp.)
Holly (Ilex spp.)
Japanese snowball (Viburnum spp.)
Lilyturf (Liriope spp.)
O'Connor's legume (Trifolium fragiferum)
Periwinkle (Vinca minor)
Pine (Pinus spp.)
Privet (Ligustrum spp.)
Rhododendron (excluding Azalea)
Spruce¹/Picea spp.)
Yew¹/Taxus spp.)

¹/DIRECTED APPLICATION recommended from budbreak, through initial growth hardening.

NOTICE TO BUYER: It is impossible to test every species and variety of ornamental or nursery plants. Plant tolerance to FUSILADE has been found to be acceptable in the specified genera and species. Neither the manufacturer nor the seller has determined whether or not FUSILADE can safely be used on plants not specified on this label. The professional user should determine if FUSILADE can be used safely prior to use.

TABLE II

DIRECTED APPLICATIONS

to avoid ornamental foliage are recommended for the following:

- Ajuga (Ajuga spp.)
- Almond (flowering) (Prunus spp.)
- Ash (Fraxinus spp.)
- Azalea (Rhododendron spp.)
- Barberry (Berberis spp.)
- Beech (Fagus spp.)
- Birch (Betula spp.)
- Boxwood (Buxus spp.)
- Cherry (flowering) (Prunus spp.)
- Crabapple (Malus spp.)
- Crepe Myrtle (Lagerstroemia indica)
- Elm (Ulmus spp.)
- Eucalyptus (Eucalyptus spp.)
- Firethorn (Pyracantha spp.)
- Magnolia (Magnolia spp.)
- Maple (Acer spp.)
- Marigold (Calandula spp.)
- Mountainash (Sorbus spp.)
- Oak (Quercus spp.)
- Peach (flowering) (Prunus spp.)
- Pear (flowering) (Pyrus spp.)
- Petunia (Petunia spp.)
- Photinia (Photinia spp.)
- Pittosporum (Pittosporum spp.)
- Plum (flowering) (Prunus spp.)
- Quince (Chaenomeles spp.)
- Redbud (Cercis spp.)
- Spirea (Spirea spp.)
- Gardenia (Gardenia spp.)
- Geranium (Pelargonium spp.)
- Horsechestnut (Aesculus spp.)
- Impatiens (Impatiens spp.)
- Juniper (Juniperus spp.)
- Sweet-gum (Liquidambar spp.)
- Tulip-tree (Liriodendron spp.)
- Willow (Salix spp.)
- Zinnia (Zinnia spp.)

GENERAL REMARKS AND PRECAUTIONS

Clean sprayer thoroughly before and after the application of FUSILADE 4E. Failure to clean sprayer thoroughly after

application may result in injury to any grass crop subsequently sprayed, including corn, sorghum, small grains, sugarcane, rice and turf.

Avoid drift to all other crops and nontarget areas. Grass crops are highly susceptible to FUSILADE 4E.

Do not plant rotational grass crops such as corn, sorghum and cereals within 60 days after last application.

Do not tank mix FUSILADE 4E with herbicides or other pesticides, with liquid fertilizers or with any other additives except as specified on this label or other supplemental labeling. Sequential applications of other herbicides except as specified on supplemental labeling within five days before or after FUSILADE 4E application has sometimes resulted in crop injury and reduced grass control.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer and User assumes the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.