

**ACCEPTED**

AUG 30 1968

Under the recent Insecticide, Fungicide, and  
Rodenticide Act, as amended, for the  
pesticide registered under  
EPA Reg. No. 10182-76

# HAVOC™

**RODENTICIDE  
PELLETS**

**KILLS RATS  
& MICE**



**Kills warfarin-resistant  
Norway Rats and House Mice.**

**Rodents may consume a lethal dose in one feeding  
with first dead rodents appearing four or five days after  
treatment begins.**

**Active Ingredient:**

<b>Brodifacoum-</b>	
3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-	
1,2,3,4-tetrahydro-1-naphthalenyl]-4-	
hydroxy-2H-1-benzopyran-2-one	0.005%
<b>Inert Ingredients</b>	<b>99.995%</b>
<b>Total</b>	<b>100.000%</b>

**KEEP OUT OF REACH  
OF CHILDREN**

**CAUTION**

May Be Harmful or Fatal if Swallowed.  
Read Additional Precautionary Statements  
on Side Panel.



EPA Reg. No. 10182-78  
EPA Est. No. 523-IL 1  
Net Contents: 45 lb (20.4 kg)

**STORAGE AND DISPOSAL**

**STORAGE:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**DISPOSAL:** Bait that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures. Dispose of bait container in a sanitary landfill or by incineration if allowed by State and local authorities.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS:** For control of Norway Rats, Roof rats and House mice in and around homes, industrial, commercial, agricultural and public buildings. HAVOC may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in temper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces the come in direct contact with food.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

**APPLICATION DIRECTIONS**

**Norway and Roof Rats:**  
Apply 4 to 16 ounces of bait (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

**House Mice:**  
Apply ¼ to ½ ounce of bait at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

**Rats & Mice:**  
Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

**MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY  
FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH  
HANDS AFTER HANDLING BAIT.**

**IF BAIT IS EATEN BY HUMANS. CALL A PHYSICIAN AT ONCE.**  
For 24-hour emergency assistance call ICI Americas  
Medical Emergency Information-1-800-327-8633  
**IF BAIT IS EATEN BY ANIMALS OR PETS.**  
For 24-hour emergency assistance call ICI Americas  
Medical Emergency Information-1-800-327-8633

In case of significant spill, call CHEMTREC 800/424-9300.

**NOTE TO PHYSICIAN OR VETERINARIAN**

This product may reduce the clotting ability of the blood and cause hemorrhaging. If this occurs, intramuscular and oral administration of vitamin K, combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Vitamin K<sub>1</sub> is antidotal at doses of 10 to 20 mg (not mg/kg) and should be given and repeated as necessary (based on monitoring of prothrombin times).

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

**NOTICE TO BUYER AND USER:** Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer and User assumes the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.