

REFLEX PA Label
RS-121,088B

P4123
10182-83
Pg 10, 3

KEEP OUT
OF REACH
OF CHILDREN

WARNING

REFLEX® 2LC Herbicide

See Side Panel for Additional
Precautionary Statements and
Statement of Practical Treatment

EPA Reg. No. 10182-83
EPA Est. No.
U.S. Patent No. 4,285,723

Net Contents: ONE U.S. GALLON
(3.79 Liters)

For Postemergence Control of Broadleaf Weeds in Soybeans

ACTIVE INGREDIENT:

Sodium salt of fomesafen	
5-[2-chloro-4-(trifluoromethyl)phenoxy]-N-	
(methylsulfonyl)-2-nitrobenzamide.	22.8%*
INERT INGREDIENTS.	77.2%
TOTAL.	100.0%

*Equivalent to 21.7% fomesafen or 2 pounds fomesafen active ingredient per gallon.

ICI (Roundel) Agricultural Products
ICI Americas Inc.
Wilmington, Delaware 19897

Jan. 9, 1989
10182-83

REFLEX 2LC IN SOYBEANS USE RATES AND WEEDS CONTROLLED

REGIONS 1 & 2

REFER TO MAP FOR DEFINITION OF SPECIFIED
GEOGRAPHIC REGIONS

Region 1- Includes the following states or portion of states where REFLEX 2LC may be applied: Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma (east of U.S. Highway 75 and east of Indian Nation Parkway), South Carolina, Tennessee, and Texas (counties east of U.S. Highway 75 and Interstate 45, additional counties of ~~Austin~~, Brazoria, Colorado, Fort Bend, Harris, Matagorda, Waller, and Wharton).

Region 2- Includes the following states or portion of states where REFLEX 2LC may be applied: Connecticut, Delaware, Indiana, Illinois (counties south of Interstate 70), Kentucky, Maine, Maryland, Massachusetts, Michigan (excluding the upper peninsula), Missouri (counties of Bollinger, Butler, Cape Girardeau, Dunklin, Madison, Mississippi, New Madrid, Pemiscot, Perry, Ripley, Scott, Stoddard, and Wayne), New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, and West Virginia.

Application Rates for Weed Growth Stages

Weeds Controlled ^c	Maximum Growth Stage Controlled At	
	1 Pt/Acre # of Leaves	1½ Pt/Acre # of Leaves
Amaranth, Palmer	4	6
Amaranth, Spiny	2	4
Anoda, Spurred	--	2
Carpetweed	6" Diameter Size	Unlimited Size
Citron (Wild Watermelon)	2	4
Cocklebur, Common ^d	2	4
Copperleaf, Hophornbeam	2	4
Copperleaf, Virginia	2	4
Crotalaria, Showy	4	6

USE OF REFLEX 2LC WITH FUSILADE 2000 + BASAGRAN - SOYBEANS ONLY

REFLEX 2LC, FUSILADE 2000 and BASAGRAN may be used together in a tank-mix as a postemergence program for broad spectrum weed control in soybeans. The tank mix should be applied at the recommended rates and growth stages to susceptible annual grass and broadleaf weed species in a manner consistent with respective labels.

The REFLEX 2LC/FUSILADE 2000/BASAGRAN tank-mix should be applied to actively growing weeds. Do not apply this tank-mix if weeds are stressed due to unfavorable temperatures, drought and/or low soil fertility. Do not apply the REFLEX 2LC/FUSILADE 2000/BASAGRAN tank-mix if soybeans show injury from prior herbicide applications.

Use a nonionic surfactant or crop oil concentrate. Apply in a minimum of 15 gallons per acre of spray mixture with spray pressures of 40-60 PSI at the nozzle tip.

Use of REFLEX 2LC With Butyrac 200 - Soybeans Only

REFLEX 2LC and Butyrac 200 may be applied in a tank mix for improved postemergence control of annual morningglory species (Entireleaf, Ivyleaf and Tall), giant ragweed and cocklebur in soybeans.

The tank mix should be applied to actively growing weeds before they exceed the growth stages described on the REFLEX 2LC label. The most consistent control is obtained when applications are made to actively growing seedlings under conditions of high soil moisture and high humidity. The most restrictive labeling of either product used applies in tank mixtures.

SEE SOYBEAN TANK MIX USE PRECAUTIONS SECTIONS FOR ADDITIONAL INFORMATION.