

# FUSILADE® 2000

**Ground Application** - Use 5-40 gallons of total spray mixture per acre and 40-60 psi pressure. Sufficient spray volume and pressure should be provided to provide complete coverage of the foliage.

Use higher volumes and increase the pressure to 60 psi if foliage is dense to ensure complete coverage.

**Band Applications** - Band application equipment should be adjusted to provide maximum coverage of foliage in the planted row. Thorough coverage can be obtained with a nozzle directed to each side of the row towards the foliage in the planted row. Applications with a single nozzle over the top of the row are not recommended.

Do not use band applications when the grass is a perennial. Reinfestation of the treated band can result from the untreated middle.

**Aerial Application** - Use a minimum of 5 gallons of finished spray volume per acre. If the foliage is dense, increase the spray volume to 10 gallons per acre to improve coverage.

**Mixing** - Fill the spray tank half-full with water and add the recommended amounts of FUSILADE 2000 1E + AMBUSH or CYMBUSH 3E while the agitator is running. Then add crop oil concentrate or nonionic surfactant and the remaining quantity of water and mix thoroughly.

### USE RESTRICTIONS

- AMBUSH AND CYMBUSH ARE RESTRICTED USE PESTICIDES FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.
- Always read and follow the restrictions and limitations for all products whether used alone or in tank mix. The most restrictive labeling of any product used applies in tank mixtures.
- Do not apply a total of more than 96 fl. oz. of FUSILADE 2000 1E herbicide per acre per season to cotton (1 lb ai/acre).
- Do not apply a total of more than 12 pints of AMBUSH per acre per season (3 lbs ai/acre).
- Do not apply a total of more than 1.0 lb ai of CYMBUSH 3E per acre per season.
- Do not harvest cotton within 90 days of FUSILADE 2000 1E application.
- Do not apply FUSILADE 2000 1E after boll set.
- Do not graze or harvest for forage or hay.
- Do not plant rotational crops such as corn, sorghum and cereals within 60 days after the last tank mix application containing FUSILADE 2000 1E.
- Avoid drift to all other crops and nontarget areas.
- Thoroughly flush spray tanks with water before and after each use.
- Apply to foliage when grasses are actively growing. When weeds are under stress use AMBUSH and CYMBUSH alone to control insects.

### DIRECTIONS FOR USE

#### NONFOOD USES

Not for use or storage in or around the home.

#### FALLOW LAND

FUSILADE 2000 1E herbicide can be used to control grass weeds during fallow periods.

Do not plant rotational crops other than cotton and soybeans within 60 days after last application.

### NONBEARING CROPS

FUSILADE 2000 1E herbicide can be used to control grass weeds in nonbearing groves, orchards or vineyards or for site preparation prior to transplanting.

APPLY AS A DIRECTED SPRAY TO PREVENT CONTACT OF SPRAY WITH FOLIAGE OF TREES OR VINES.

DO NOT APPLY TO TREES, VINES OR OTHER LISTED CROPS THAT WILL BE HARVESTED WITHIN ONE YEAR AFTER APPLICATION.

DO NOT APPLY TO ASPARAGUS FROM WHICH SPEARS WILL BE HARVESTED WITHIN ONE YEAR OF TREATMENT.

FUSILADE 2000 1E herbicide is recommended on the following nonbearing crops:

Acerola (West Indian Cherry)*	Lemons
Almonds	Limes
Apples	Macadamia
Apricots	Mango*
Asparagus	Nectarines
Avocados	Olives
Berries	Oranges
Cherries	
Coffee	Peaches
Dates	Pears
Figs	Pecans
Filberts	Pistachios
Ginseng*	Plums
Grapes	Pomegranates
Grapelruit*	Prunes
Guava*	Tangelos
Jojoba	Tangerines
Kiwi*	Walnuts

\*Not applicable in California

### ORNAMENTALS

FUSILADE 2000 1E herbicide can be used to control grass weeds in many newly transplanted and established non-grassy ornamentals, trees, shrubs and ground covers. See the following Tables I through IV for specific recommended uses.

Application of FUSILADE 2000 1E may be made over-the-top to ornamentals listed in Table I. Plants listed in Tables II through IV should have a directed spray applied to prevent contact of spray with foliage. If the growth habit of the plant or other factors do not allow a directed spray, an over-the-top application can be made to plants in Table II-IV. However, there is the possibility of damage in the range listed for each table when applied at label rates.

DO NOT APPLY TO ORNAMENTALS THAT MAY BE HARVESTED FOR FOOD WITHIN ONE YEAR AFTER APPLICATION.

**NOTICE TO BUYER AND USER:** It is impossible to test every species and variety or cultivar of ornamental or nursery plants under all conditions. Plant tolerance of pesticides vary as conditions vary. Plant tolerance of FUSILADE 2000 1E at label rates has been found to be acceptable within the ranges specified for the indicated genera and species. Neither the manufacturer nor the seller has determined whether or not FUSILADE 2000 1E can safely be used on plants not specified on this label. The professional user should determine if FUSILADE 2000 1E can be used safely prior to use.

### APPLICATION DIRECTIONS - NON - FOOD USES

**Timing** - For non-food use apply to actively growing grass weeds before they exceed the recommended growth stages shown on this page.

Do not apply FUSILADE 2000 1E herbicide if rainfall is expected within 1 hour.

HERBICIDES

ACCEPTED  
APR 9 1986  
U.S. Dept. of Internal Security  
EPA Reg. No. 10182-88