

outer label
1416 10182-93

ACCEPTED
AUG 30 1993
Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 10182-93

HOVOC®

RODENTICIDE BAIT PACK (PELLETS)

KILLS RATS & MICE

Kills warfarin-resistant
Norway Rats and House Mice.

Rodents may consume a lethal dose in one feeding with first
dead rodents appearing four or five days after treatment begins.

Active Ingredient:

Brodifacoum-	
3-[3-(4-bromo-[1,1'-biphenyl]-4-yl)-	
1,2,3,4-tetrahydro-1-naphthalenyl]-4-	
hydroxy-2H-1-benzopyran-2-one	0.005%
Inert Ingredients	99.995%
	100.000%

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

May Be Harmful or Fatal if Swallowed.
Read Additional Precautionary
Statements on Side Panel



ICI Americas Inc.
Agricultural Chemicals Division
Wilmington, Delaware 19897

EPA Reg. No. 10182-93
EPA Est. No. 2393 W-1
Net Contents: 5/25 gm (0.99 oz) bait packs

10/1

2

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container except for holding additional HAVOC rodenticide.

Storage: Store in original container only in dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of bait container in a sanitary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway Rats, Roof rats and House mice in and around homes, industrial, commercial, agricultural and public buildings. HAVOC may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS

Norway and Roof Rats:

Apply 4 to 16 bait packs (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

House Mice:

Apply 1 bait pack at intervals of 8 to 12 feet per placement. 2 bait packs may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

Rats & Mice:

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

26/11

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance, call ICI Americas Medical Emergency Information-1-800-327-8633

IF BAIT IS EATEN BY ANIMALS OR PETS. For 24-hour Emergency assistance call ICI Americas Medical Emergency Information Center - 1-800-327-8633

In case of significant spill, call CHEMTREC 800/424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN
This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS
This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer and User assumes the risk of any such use. **SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.**

30/1

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic life. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

STORAGE AND DISPOSAL

Refer to label on container for full storage and disposal directions.

DIRECTIONS FOR USE

Refer to label on container for full use directions.



EPA Reg No 10182-93
EPA Est No 523 K-1
Net Contents 25 gm (0.88 oz)

Havoc™

RODENTICIDE BAIT PACK (PELLETS)

For Effective Control of Commercial, Home and Wild

Kills warfarin-resistant Norway Rats and House Mice.
Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

Active Ingredient:	
Brodifacoum	
3334 (bromo) 1,1-bis(4-chlorophenyl) 4-yl-1,2,3,4-tetrahydro-1-naphthylidene-4-hydroxy-2H-1-benzopyran-2-one	0.000%
Inert Ingredients	99.999%
Total	100.000%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

May be Harmful or Fatal if Swallowed. Read Additional Precautionary Statements on Back Panel.

PRECAUTIONARY STATEMENTS

HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF SWALLOWED OR INHALED, CALL A PHYSICIAN AT ONCE for 24 hour emergency assistance, call ICI Americas Medical Emergency Information-1-800-327-8633. IF SWALLOWED BY HUMANS OR PETS, FOR 24-hour Emergency assistance call ICI Americas Medical Emergency Information Center - 1-800-327-8633. **NOTE TO PHYSICIAN OR VETERINARIAN**

This product may reduce the clotting ability of the blood and cause hemorrhaging if this occurs, an irregular and oral administration of vitamin K₁ combined with blood transfusions are indicated as in the case of hemorrhage caused by over-dosage of bromadiolone. Vitamin K₁ is present at doses of 10 to 20 mg/kg and should be given and repeated as necessary based on monitoring of prothrombin time.

Under the Federal Insecticide, Fungicide and Rodenticide Act, it is unlawful to sell, distribute, or use any pesticide registered under EPA Reg. No. 10182-93

1/2/94

