

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060. Approval expires 2-28-95

|  |   |   |  |
|--|---|---|--|
|  | United States<br><b>Environmental Protection Agency</b><br>Washington, DC 20460 | <input type="checkbox"/> Registration     | OPP Identifier Number<br><br><b>264382</b> |
|  |   | <input type="checkbox"/> Amendment        |  |
|  |   | <input checked="" type="checkbox"/> Other |  |

**Application for Pesticide - Section I**

|   |   |   |
|---|---|---|
| 1. Company/Product Number<br>10182-96   | 2. EPA Product Manager<br>George T. LaRocca   | 3. Proposed Classification<br><input type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product (Name) (Alt. Brand Name for<br>Warrior Insecticide Karate Insecticide)   | PM#<br>03   |   |
| 5. Name and Address of Applicant (Include ZIP Code)<br>Zeneca Agricultural Products<br>P.O. Box 15458<br>Wilmington, DE 19850-5458<br><input type="checkbox"/> Check if this is a new address | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar on the basis of its composition and labeling to:<br>EPA Reg. No. _____ <b>NOV 30 1998</b><br>Product Name _____ |   |

**Section - II**

|  |   |
|--|---|
| <input type="checkbox"/> Amendment - Explain below.                            | <input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated 1/13/98 |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application.  |
| <input checked="" type="checkbox"/> Notification - Explain below.              | <input type="checkbox"/> Other - Explain below.   |

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

The addition of rice, which is already approved on the Karate Insecticide label.

**Section - III**

|  |   |  |   |   |                                  |
|--|---|--|---|---|----------------------------------|
| 1. Material This Product Will Be Packaged In:  |   |  |   | 2. Type of Container  |                                  |
| Child-Resistant Packaging<br><input type="checkbox"/> Yes<br><input type="checkbox"/> No                     | Unit Packaging<br><input type="checkbox"/> Yes<br><input type="checkbox"/> No | Water Soluble Packaging<br><input type="checkbox"/> Yes<br><input type="checkbox"/> No |   | <input type="checkbox"/> Metal                              | <input type="checkbox"/> Plastic |
| * Certification must be submitted  |   | If "Yes" Unit Packaging wgt.   | No. per container   | <input type="checkbox"/> Glass                              | <input type="checkbox"/> Paper   |
|  |   | If "Yes" Package wgt.  | No. per container   | <input type="checkbox"/> Other (Specify) _____              |                                  |
| 3. Location of Net Contents Information<br><input type="checkbox"/> Label <input type="checkbox"/> Container |   | 4. Size(s) Retail Container  |   | 5. Location of Label Directions<br><input type="checkbox"/> |                                  |
| 6. Manner in Which Label is Affixed to Product   |   |  | <input type="checkbox"/> Lithograph<br><input type="checkbox"/> Paper glued<br><input type="checkbox"/> Stenciled | <input type="checkbox"/> Other _____                        |                                  |

**Section - IV**

|  |                                  |   |
|--|----------------------------------|---|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)  |                                  |   |
| Name<br>Nancy Bockrath   | Title<br>Regulatory Assistant    | Telephone No. (Include Area Code)<br>302-886-1066 |
| <b>Certification</b><br>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. |                                  | 6. Date Application Received<br>(Stamped)         |
| 2. Signature<br>   | 3. Title<br>Regulatory Assistant |   |
| 4. Typed Name<br>Nancy Bockrath  | 5. Date<br>11/16/98              |   |

2725

**RESTRICTED USE PESTICIDE**  
**Due to Toxicity to Fish and Aquatic Organisms**  
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

# WARRIOR®

*For Use to Control Certain Insects on Listed Crops*

**KEEP OUT OF REACH  
OF CHILDREN  
DANGER  
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

|  |               |
|--|---------------|
| <b>ACTIVE INGREDIENT:</b>  |               |
| Lambda-cyhalothrin   |               |
| [1 $\alpha$ (S*),3 $\alpha$ (Z)]-(±)-cyano-(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate | 13.1%         |
| <b>INERT INGREDIENTS:</b>  | 86.9%         |
| <b>TOTAL</b>   | <b>100.0%</b> |

WARRIOR® Insecticide contains 1 pound of active ingredient per gallon and is an emulsifiable concentrate.  
EPA Reg. No. 10182-96

### STATEMENT OF PRACTICAL TREATMENT

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.  
**IF SWALLOWED:** Do not induce vomiting. Call a physician immediately.  
**IF INHALED:** Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.  
**IF IN EYES:** Hold eyelids open and flush with steady, gentle stream of water. Remove any contact lenses and repeat flushing procedure for several minutes. Get medical attention if irritation persists.  
**NOTE TO PHYSICIAN:** Induced vomiting as first aid for this substance may result in increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent. Vomiting should be induced only under professional supervision.

**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).**  
**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.**


See inside for Precautionary Statements.

COMPLETE DIRECTIONS FOR USE

CR1098

## ZENECA

**NOTIFICATION**  
**NOV 30 1998**



**CONDITIONS OF SALE  
AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

*The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.*

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

TABLE OF CONTENTS

STATEMENT OF PRACTICAL TREATMENT ..... 4

PRECAUTIONARY STATEMENTS ..... 4

DIRECTIONS FOR USE ..... 5

AGRICULTURAL USE REQUIREMENTS ..... 5

STORAGE AND DISPOSAL ..... 6

GENERAL DIRECTIONS FOR USE ..... 6

Spray Drift Precautions ..... 7

Chemigation ..... 8

Use Precautions - Sprinkler Irrigation Application ..... 8

Crop Uses: Alfalfa, Alfalfa Grown for Seed ..... 9

    Cole Crops ..... 11

    Corn: Field Corn, Popcorn, Seed Corn ..... 12

    Sweet Corn ..... 13

    Cotton ..... 14

    Lettuce (Head and Leaf) ..... 15

    Onion (Bulb) and Garlic ..... 16

    Peanut ..... 16

    Rice ..... 17

    Sorghum (Grain) ..... 18

    Soybean ..... 19

    Sunflower ..... 20

    Tomato and Tomatillo ..... 21

    Wheat, Wheat Hay and Triticale ..... 21

Rate Conversion Chart ..... 22

5 8 25

### STATEMENT OF PRACTICAL TREATMENT

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF SWALLOWED:** Do not induce vomiting. Call a physician immediately.

**IF INHALED:** Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**IF IN EYES:** Hold eyelids open and flush with steady, gentle stream of water. Remove any contact lenses and repeat flushing procedure for several minutes. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** Induced vomiting as first aid for this substance may result in increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent. Vomiting should be induced only under professional supervision.

**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).**  
**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.**

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### **DANGER**

**CORROSIVE. CAUSES SKIN DAMAGE. MAY BE FATAL IF SWALLOWED OR INHALED. CAUSES EYE INJURY. HARMFUL IF ABSORBED THROUGH SKIN. MAY CAUSE ALLERGIC SKIN REACTIONS. DO NOT** get on skin, in eyes, or on clothing. **DO NOT** breathe spray mist.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 to 30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

#### **Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistant category selection chart.

Applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading
- For exposures in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)
- For exposures outdoors, dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**PHYSICAL AND CHEMICAL HAZARDS**

Combustible liquid. Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

*Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.*

This labeling must be in the possession of the user at the time of application.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton  $\geq$  14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage and disposal.

**STORAGE:** Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL**

Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**FOR BULK/MINI-BULK, E-Z HANDLER® AND BOOMERANG CONTAINERS**

Return container to ZENECA, Inc. for reuse with seal intact and in salable condition.

**CONTAINER PRECAUTIONS**

Before refilling RETURNABLE CONTAINERS, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

**REFILL ONLY WITH WARRIOR® Insecticide.** The contents of RETURNABLE CONTAINERS cannot be completely removed by cleaning. Refilling with materials other than WARRIOR Insecticide will result in contamination and may weaken container.

After filling and before transporting, check for leaks.

*Do not refill or transport damaged or leaking container.*

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

**GENERAL DIRECTIONS FOR USE**

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air or 10 gallons per acre by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, WARRIOR Insecticide may be applied before, during or after planting. For soil incorporated applications, use higher rates for improved control.

*Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.*

*If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.*

**WARRIOR® Insecticide may be applied in the following states or portions of states:** Arizona, California, Colorado, Connecticut, Delaware, Florida (all counties except Baker, Bay, Calhoun, Columbia, Duval, Escambia, Franklin, Gadsden, Gulf, Hamilton, Holmes, Jackson, Jefferson, Liberty, Leon, Madison, Nassau, Okaloosa, Santa Rosa, Suwannee, Wakulla, Walton, and Washington), Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Jersey, New Hampshire, New York, North Dakota, Ohio, Oregon, Pennsylvania, Puerto Rico, Rhode Island, South Dakota, Tennessee (counties of Stewart, Robertson, Montgomery, Cheatam, and Sumner), Utah, Vermont, Virginia (counties of Accomac and Northampton), Washington, West Virginia, Wisconsin, and Wyoming.

**SPRAY DRIFT PRECAUTIONS**

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.**

- Do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams, marshes, pot holes, or natural ponds; estuaries and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultralow volume (ULV) application is made.
- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.
- Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.
- Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
- Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
- Do not cultivate within 10 feet of the aquatic area so as to allow growth of a vegetative filter strip.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.
- Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

In the State of New York, a 25 foot vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial or ground application. For aerial applications, the 25 foot vegetated non-cropped buffer strip for runoff protection would be part of the larger 150 foot buffer strip (or 450 foot buffer strip for ULV application) required for spray drift.



**CHEMIGATION**

**SPRINKLER IRRIGATION APPLICATION:** Apply WARRIOR Insecticide at rates and timing described elsewhere in this label. Apply with center pivot or continuous-move equipment distributing 1/2 acre-inch or less during treatment. In general, use the least amount of water required for proper distribution and coverage. If stationary systems (solid sets, handlines or wheel lines other than continuous-move) are used, WARRIOR Insecticide should be injected into no more than the last 20-30 minutes of the set. Do not apply when winds are greater than 10-15 mph to avoid drift or wind skips. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

**USE PRECAUTIONS - SPRINKLER IRRIGATION APPLICATION**

- A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B. *Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.*
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Do not apply through chemigation systems connected to public water systems.

Apply WARRIOR® Insecticide as shown in the following charts:



| SPRAY RECOMMENDATIONS                    |   |             |           |  |
|--|---|-------------|-----------|--|
| Crop                                     | Target Pests  | Rate        |           | Remarks  |
|  |   | lb ai/A     | fl oz/A   |  |
| Alfalfa,<br>Alfalfa<br>grown for<br>seed | Alfalfa Caterpillar<br>Cutworm Spp.<br>Green Cloverworm<br>Looper Spp.<br>Velvetbean Caterpillar<br>Webworm Spp.<br>Leafhopper Spp.<br>Threecornered Alfalfa Hopper   | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air or 10 gallons per acre by ground. When foliage is dense and/or pest populations are high 5-10 gallons per acre by air or 20 gallons per acre by ground and higher use rates are recommended. Use higher rates for increased residual control.</li> <li>Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. Avoid direct application to bee shelters.</li> <li>Do not apply more than 0.03 lb ai (0.24 pints) per acre per cutting. Do not apply more than 0.12 lb ai (0.96 pints) per acre per season. Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay.</li> </ul> |
|  | Armyworm<br>Corn Earworm<br>Fall Armyworm <sup>1</sup><br>Western Yellow-striped Armyworm<br>Yellow-striped Armyworm<br>Alfalfa Weevil<br>Bean Leaf Beetle (Adult)<br>Blister Beetle Spp.<br>Clover Leaf Weevil Spp.<br>Clover Root Borer (Adult)<br>Clover Root Curculio Spp. (Adult)<br>Clover Stem Borer (Adult)<br>Cowpea Curculio (Adult)<br>Cowpea Weevil (Adult)<br>Cucumber Beetle Spp. (Adult)<br>Egyptian Alfalfa Weevil<br>Grape Colaspis (Adult)<br>Green June Beetle (Adult)<br>Japanese Beetle (Adult)<br>Mexican Bean Beetle<br>Pea Weevil (Adult) | 0.02-0.03   | 2.56-3.84 |  |

11725

| SPRAY RECOMMENDATIONS                    |   |         |         |  |
|--|---|---------|---------|--|
| Crop                                     | Target Pests  | Rate    |         | Remarks  |
|  |   | lb ai/A | fl oz/A |  |
| Alfalfa,<br>Alfalfa<br>grown for<br>seed | Sweet Clover Weevil (Adult)<br>Whitefringed Beetle Spp.<br>(Adult)<br>Meadow Spittlebug<br>Plant Bug Spp. including<br>Lygus Spp. <sup>1</sup><br>Stink Bug Spp.<br>Alfalfa Seed Chalcid (Adult)<br>Blue Alfalfa Aphid<br>Cowpea Aphid<br>Green Peach Aphid <sup>2</sup><br>Pea Aphid<br>Spotted Alfalfa Aphid<br>Thrips Spp.<br>Grasshopper Spp.<br><br>Beet Armyworm <sup>1</sup><br>Blotch Leafminer <sup>2</sup><br>Spider Mites <sup>2</sup> | 0.03    | 3.84    | <ul style="list-style-type: none"> <li>• Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>• Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air or 10 gallons per acre by ground. When foliage is dense and/or pest populations are high 5-10 gallons per acre by air or 20 gallons per acre by ground and higher use rates are recommended. Use higher rates for increased residual control.</li> <li>• Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. Avoid direct application to bee shelters.</li> <li>• Do not apply more than 0.03 lb ai (0.24 pints) per acre per cutting. Do not apply more than 0.12 lb ai (0.96 pints) per acre per season. Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay.</li> </ul> <p><sup>1</sup> For control of first and second instar only.<br/> <sup>2</sup> Suppression only.<br/> <sup>3</sup> See resistance statement under General Directions for Use.</p> |

| SPRAY RECOMMENDATIONS  |  |             |           |   |
|--|--|-------------|-----------|---|
| Crop   | Target Pests   | Rate        |           | Remarks   |
|  |  | lb ai/A     | fl oz/A   |   |
| Broccoli,<br>Brussels<br>Sprouts,<br>Cabbage,<br>Cavalo,<br>Broccoli,<br>Cauliflower,<br>Chinese<br>Broccoli<br>(gai ion),<br>Chinese<br>Cabbage<br>(napaa),<br>Chinese<br>Mustard<br>Cabbage<br>(gai choy),<br>Kohirabi | Alfalfa Looper<br>Cabbage Looper<br>Imported Cabbageworm<br>Southern Cabbageworm<br>Cutworm Spp.<br>Cabbage Webworm  | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>• Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>• Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>• Do not apply within 1 day of harvest.</li> <li>• Do not apply more than 0.24 lb ai (1.92 pints) per acre per season.</li> </ul> <p><sup>1</sup>For control of first and second instar only.<br/> <sup>2</sup>Suppression only.<br/> <sup>3</sup>See resistance statement under General Directions for Use.</p> |
|  | Diamondback Moth <sup>1</sup><br>Armyworm<br>Beet Armyworm <sup>2</sup><br>Fall Armyworm <sup>1</sup><br>Yellow-striped Armyworm<br>Corn Earworm<br>Flea Beetle Spp.<br>Japanese Beetle (Adult)<br>Vegetable Weevil (Adult)<br>Grasshopper Spp.<br>Leafhopper Spp.<br>Plant Bug Spp. including<br>Lygus Spp. <sup>3</sup><br>Stink Bug Spp.<br>Meadow Spittlebug<br>Aphid Spp. <sup>3</sup><br>Whitefly Spp. <sup>3</sup><br>Thrips Spp. <sup>3</sup><br>Spider Mite Spp. <sup>3</sup> | 0.02-0.03   | 2.56-3.84 |   |

| SPRAY RECOMMENDATIONS                      |   |             |           |   |
|--|---|-------------|-----------|---|
| Crop                                       | Target Pests  | Rate        |           | Remarks   |
|  |   | lb ai/A     | fl oz/A   |   |
| Corn<br>Field Corn<br>Popcorn<br>Seed Corn | Cutworm Spp.<br>Western Bean Cutworm <sup>1</sup><br>Corn Earworm <sup>1</sup><br>Green Cloverworm<br>Meadow Spittlebug   | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods.</li> <li>Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3- to 5-day intervals if needed. WARRIOR Insecticide may only suppress heavy infestations and/or subsequent migrations.</li> <li>For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program use a minimum of 3.84 fl oz/A (0.03 lb ai/A).</li> <li>Do not apply within 21 days of harvest. Do not apply more than 0.12 lb ai (0.96 pints) per acre per season. Do not apply more than 0.06 lb ai (0.48 pints) after silk initiation. Do not apply more than 0.03 lb ai (0.24 pints) after corn has reached the milk stage (yellow kernels with milky fluid).</li> <li>For control before the larva bores into the plant stalk or ear.</li> <li>For control of the first and second instar only.</li> <li>Suppression only.</li> <li>See resistance statement under General Directions for Use.</li> </ul> |
|  | Tobacco Budworm <sup>1,2</sup><br>European Corn Borer <sup>1</sup><br>Southwestern Corn Borer <sup>1</sup><br>Stalk Borer <sup>1</sup><br>Hop Vine Borer <sup>1</sup><br>Armyworm <sup>2</sup><br>Fall Armyworm <sup>2</sup><br>Yellow-striped Armyworm <sup>2</sup><br>Webworm Spp.<br>Flea Beetle Spp.<br>Western Corn Rootworm Beetle (Adult)<br>Northern Corn Rootworm Beetle (Adult)<br>Southern Corn Rootworm Beetle (Adult)<br>Mexican Corn Rootworm Beetle (Adult)<br>Bean Leaf Beetle<br>Cereal Leaf Beetle<br>Japanese Beetle (Adult)<br>Sap Beetle (Adult)<br>Stink Bug Spp.<br>Grasshopper Spp.<br>Corn Leaf Aphid <sup>1</sup><br>Bird Cherry-Oat Aphid <sup>2</sup><br>English Grain Aphid <sup>2</sup> | 0.02-0.03   | 2.56-3.84 |   |
|  | Beet Armyworm <sup>2,3</sup><br>Chinch Bug<br>Green Bug <sup>3,4</sup>  | 0.03        | 3.84      |   |

| SPRAY RECOMMENDATIONS |  |           |           |   |
|-----------------------|--|-----------|-----------|---|
| Crop                  | Target Pests   | Rate      |           | Remarks   |
|                       |  | lb ai/A   | fl oz/A   |   |
| Sweet Corn            | Corn Earworm<br>Fall Armyworm <sup>1</sup><br>Southern Armyworm <sup>1</sup><br>Beet Armyworm <sup>3</sup><br>Yellow-Striped Armyworm <sup>1</sup><br>Cutworm Spp.<br>Western Bean Cutworm<br>Webworm Spp.<br>European Corn Borer<br>Southwestern Corn Borer<br>Common Cornstalk Borer<br>Western Corn Rootworm Beetle (Adult)<br>Northern Corn Rootworm Beetle (Adult)<br>Southern Corn Rootworm Beetle (Adult)<br>Mexican Corn Rootworm Beetle (Adult)<br>Japanese Beetle (Adult)<br>Sap Beetle (Adult)<br>Flea Beetle Spp.<br>Tarnished Plant Bug<br>Stink Bug Spp.<br>Chinch Bug<br>Aster Leafhopper<br>Grasshopper Spp.<br>Aphid Spp. <sup>2</sup><br>Spider Mite Spp. <sup>2</sup> | 0.02-0.03 | 2.56-3.84 | *Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 4 or more days. <i>Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods and should be targeted for control before insects enter the stalk or ear.</i><br><br>*Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage and ears (if present). When applying by air, apply in a minimum of 2 gallons of water per acre.<br><br>*For control of adult corn rootworm beetles ( <i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program use a minimum of 3.2 fl oz/A (0.025 lb ai/A).<br><br>*Do not apply within 1 day of harvest.<br><br>*Do not apply more than 0.48 lb ai (3.84 pints) per acre per season.<br><br>For control of first and second instar only.<br><sup>1</sup> Suppression only.<br><sup>2</sup> See resistance statement under General Directions for Use. |
|                       | Corn Silkyfly (Adult) <sup>1</sup>   | 0.03      | 3.84      |   |

| SPRAY RECOMMENDATIONS |   |            |           |   |
|-----------------------|---|------------|-----------|---|
| Crop                  | Target Pests  | Rate       |           | Remarks   |
|                       |   | lb ai/A    | fl oz/A   |   |
| Cotton                | Cutworm Spp.<br>Tobacco Thrips<br>Soybean Thrips  | 0.015-0.02 | 1.92-2.56 | <ul style="list-style-type: none"> <li>• Apply as required by scouting, usually at intervals of 5 to 7 days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>• Apply with ground or air equipment using sufficient water to obtain full coverage of foliage.</li> <li>• Applications may also be made with equipment adapted and calibrated for ULV sprays. WARRIOR Insecticide may be mixed with once-refined vegetable oil and applied in a minimum of at least one quart of finished spray per acre.</li> <li>• Under light bollworm/budworm infestation levels, 0.02 lb ai/A may be applied in conjunction with intense field monitoring.</li> <li>• For boll weevil control spray on a 3 to 5 day schedule.</li> <li>• When applied according to label directions for control of cotton bollworm and tobacco budworm, WARRIOR Insecticide also provides ovicidal control of unhatched <i>Heliothis</i> spp. eggs.</li> <li>• Do not apply within 21 days of harvest.</li> <li>• Do not graze livestock in treated areas.</li> <li>• Do not apply more than 1.6 pints (0.2 lb ai) per acre per season.</li> <li>• Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include AMBUSH<sup>1</sup> insecticide, Ammo<sup>2</sup> insecticide, Asana<sup>3</sup> XL insecticide, Baythroid<sup>2</sup> emulsifiable pyrethroid insecticide, Capture<sup>2</sup> insecticide/miticide, Danitol<sup>2</sup> Spray insecticide/miticide, Decis<sup>1</sup> insecticide, Fury<sup>2</sup> insecticide, KARATE<sup>2</sup> insecticide, Mustang<sup>1</sup> insecticide, Pounce<sup>1</sup>, Scout X-TRA<sup>2</sup> insecticide, SynerGin<sup>2</sup> insecticide, and WARRIOR<sup>4</sup> IEC insecticide.</li> </ul> <p>For control of the first and second instar only.<br/> <sup>1</sup>Suppression only.<br/> <sup>2</sup>See resistance statement under General Directions for Use.</p> |
|                       | Lygus Bug Spp. <sup>2</sup><br>Pink Bollworm<br>Cabbage Looper<br>Cotton Leafperforator<br>Saltmarsh Caterpillar<br>Cotton Leafworm<br>Cotton Fleahopper  | 0.02-0.03  | 2.56-3.84 |   |
|                       | Cotton Bollworm<br>Tobacco Budworm <sup>1</sup><br>Boll Weevil<br>Fall Armyworm<br>Beet Armyworm <sup>1</sup><br>European Corn Borer<br>Brown Stink Bug<br>Green Stink Bug<br>Southern Green Stink Bug<br>Twospotted Spider Mite <sup>2</sup><br>Cotton Aphid <sup>2</sup><br>Bandedwing Whitefly <sup>2,3</sup><br>Sweetpotato Whitefly <sup>2,3</sup> | 0.025-0.04 | 3.20-5.12 |   |

| SPRAY RECOMMENDATIONS      |   |             |           |   |
|----------------------------|---|-------------|-----------|---|
| Crop                       | Target Pests  | Rate        |           | Remarks   |
|                            |   | lb ai/A     | fl oz/A   |   |
| Lettuce<br>(Head and Leaf) | Alfalfa Looper<br>Cabbage Looper<br>Imported Cabbageworm<br>Cutworm Spp.<br>Saltmarsh Caterpillar<br>Green Cloverworm   | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>• Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>• Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>• Do not apply within 1 day of harvest.</li> <li>• Do not apply more than 0.3 lb ai (2.4 pints) per acre per season.</li> </ul> <p><sup>1</sup>For control of first and second instar only.<br/> <sup>2</sup>Suppression only.<br/> <sup>3</sup>See resistance statement under General Directions for Use.</p> |
|                            | Diamondback Moth <sup>1</sup><br>Armyworm<br>Beet Armyworm <sup>1,2</sup><br>Fall Armyworm <sup>1</sup><br>Southern Armyworm<br>Corn Earworm<br>Tobacco Budworm <sup>1</sup><br>European Corn Borer<br>Flea Beetle Spp.<br>Japanese Beetle (Adult)<br>Vegetable Weevil (Adult)<br>Grasshopper Spp.<br>Leafhopper Spp.<br>Plant Bug Spp. Including<br>Lygus Spp. <sup>3</sup><br>Stink Bug Spp.<br>Meadow Spittlebug<br>Aphid Spp. <sup>2</sup><br>Whitefly Spp. <sup>2,3</sup><br>Spider Mite Spp. <sup>1</sup> | 0.02-0.03   | 2.56-3.84 |   |



| SPRAY RECOMMENDATIONS   |  |             |           |  |
|-------------------------|--|-------------|-----------|--|
| Crop                    | Target Pests   | Rate        |           | Remarks  |
|                         |  | lb ai/A     | fl oz/A   |  |
| Onion (Bulb) and Garlic | Cutworm Spp.<br>Seedcorn Maggot (Adult)<br>Onion Maggot (Adult)<br>Leafminer Spp. (Adult)  | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>• Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>• Use the higher label rates as thrips population increases and avoid rescue situations.</li> <li>• Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>• For thrips control by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturers use directions) may enhance the deposition of the spray and increase plant coverage.</li> <li>• Do not apply within 14 days of harvest. Do not apply more than 0.24 lb ai (1.92 pints) per acre per season.</li> <li>For control of the first and second instar only.</li> <li><sup>1</sup>Suppression only.</li> <li><sup>2</sup>See resistance statement under General Directions for Use.</li> </ul> |
|                         | Amyworm Spp. <sup>1</sup><br>Onion Thrips<br>Tobacco Thrips<br>Western Flower Thrips <sup>2,3</sup><br>Flower Thrips <sup>4</sup><br>Aphid Spp. <sup>5</sup><br>Plant Bug Spp.<br>Stink Bug Spp. | 0.02-0.03   | 2.56-3.84 |  |
| Peanut                  | Cutworm Spp.<br>Green Cloverworm<br>Velvetbean Caterpillar<br>Red-necked Peanut Worm<br>Potato Leafhopper  | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>• Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>• Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>• Do not apply within 14 days of harvest. Do not apply more than 0.12 lb ai (0.96 pints) per acre per season.</li> <li>• Do not graze livestock in treated areas. Do not use treated vines or hay for animal feed.</li> <li>For control of the first and second instar only.</li> <li><sup>1</sup>Suppression only.</li> <li><sup>2</sup>See resistance statement under General Directions for Use.</li> </ul>  |
|                         | Corn Earworm<br>Fall Armyworm<br>Bean Leaf Beetle<br>Southern Corn Rootworm (Adult)<br>Vegetable Weevil<br>Whitefringed Beetle (Adult)<br>Stink Bug Spp.<br>Tobacco Thrips<br>Grasshopper Spp.   | 0.02-0.03   | 2.56-3.84 |  |
|                         | Beet Armyworm <sup>1,3</sup><br>Soybean Looper <sup>2,3</sup><br>Lesser Cornstalk Borer <sup>2</sup><br>Spider Mite Spp. <sup>2</sup><br>Aphid Spp. <sup>2</sup>                                 | 0.03        | 3.84      |  |

| SPRAY RECOMMENDATIONS |  |            |           |  |
|-----------------------|--|------------|-----------|--|
| Crop                  | Target Pests   | Rate       |           | Remarks  |
|                       |  | lb ai/A    | fl oz/A   |  |
| Rice                  | True Armyworm<br>Fall Armyworm<br>Yellow-striped Armyworm<br>Rice Water Weevil (Adult)<br>Rice Stink Bug<br>Chinch Bug<br>Grasshopper Spp.<br>Leafhopper Spp.<br>Oat Birdcherry Aphid<br>Green Bug | 0.025-0.04 | 3.20-5.10 | <ul style="list-style-type: none"> <li>• Apply as required by scouting. Timing and frequency of application should be based upon local economic thresholds. Determine the need for repeat applications, usually at intervals of 5-7 days, by scouting.</li> <li>• WARRIOR Insecticide can be safely used when propanil products are being used for weed control.</li> <li>• Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 5 gallons of water per acre.</li> <li>• For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations.</li> <li>• For control of rice water weevil in water seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations.</li> <li>• Green bug is known to have many biotypes. WARRIOR Insecticide may only provide suppression. If satisfactory control is not achieved with the first application of WARRIOR Insecticide, a resistant biotype may be present. Use alternate chemistry for control.</li> <li>• Do not release flood water within 7 days of an application.</li> <li>• Do not apply more than 0.12 lb ai (0.96 pints) per acre per season. Do not apply more than 0.08 lb ai (0.64 pints) per acre within 28 days of harvest or more than 0.04 lb ai (0.32 pints) per acre within 21 days of harvest.</li> <li>• Do not apply within 21 days of harvest.</li> <li>• Do not use treated rice fields for the aquaculture of edible fish and crustacea.</li> <li>• Do not apply as an ultra-low volume (ULV) spray.</li> </ul> |

| SPRAY RECOMMENDATIONS |   |            |           |  |
|-----------------------|---|------------|-----------|--|
| Crop                  | Target Pests  | Rate       |           | Remarks  |
|                       |   | lb ai/A    | fl oz/A   |  |
| Sorghum<br>(Grain)    | Cutworm Spp.<br>Sorghum Midge   | 0.015-0.02 | 1.92-2.56 | <ul style="list-style-type: none"> <li>• Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>• Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>• For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5-day intervals if needed.</li> <li>• For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3- to 5-day intervals if needed. WARRIOR insecticide may only suppress heavy infestations and/or subsequent migrations.</li> </ul> |
|                       | Armyworm<br>Beet Armyworm <sup>1</sup><br>Fall Armyworm <sup>1</sup><br>Yellow-striped Armyworm <sup>1</sup><br>Corn Earworm<br>Webworm Spp.<br>European Corn Borer <sup>2</sup><br>Southwestern Corn Borer <sup>2</sup><br>Lesser Cornstalk Borer <sup>2</sup><br>Flea Beetle Spp.<br>Stink Bug Spp.<br>Grasshopper Spp. | 0.02-0.03  | 2.56-3.84 |  |
|                       | Chinch Bug  | 0.03       | 3.84      | <ul style="list-style-type: none"> <li>• Do not apply within 30 days of harvest. Do not apply more than 0.08 lb ai (0.64 pints) per acre per season. Do not apply more than 0.06 lb ai (0.48 pints) per acre per season after crop emergence. Do not apply more than 0.02 lb ai (0.16 pints) per acre per season once crop is in soft dough stage.</li> <li>• Do not graze livestock in treated areas or harvest for fodder, silage or hay.</li> </ul>   |

<sup>1</sup>For control of the first and second instar only.  
<sup>2</sup>For control before the larva bores into the plant stalk.  
<sup>3</sup>See resistance statement under General Directions for Use.

| SPRAY RECOMMENDATIONS |   |             |           |  |
|-----------------------|---|-------------|-----------|--|
| Crop                  | Target Pests  | Rate        |           | Remarks  |
|                       |   | lb ai/A     | fl oz/A   |  |
| Soybean               | Corn Earworm<br>Velvetbean Caterpillar<br>Green Cloverworm<br>Cabbage Looper<br>Painted Lady (Thistle) Caterpillar<br>Saltmarsh Caterpillar<br>Woollybear Caterpillar<br>Cutworm Spp.<br>Bean Leaf Beetle<br>Mexican Bean Beetle<br>Western Corn Rootworm Beetle (Adult)<br>Northern Corn Rootworm Beetle (Adult)<br>Southern Corn Rootworm Beetle (Adult)<br>Mexican Corn Rootworm Beetle (Adult)<br>Three-Cornered Alfalfa Hopper<br>Potato Leafhopper<br>Thrips Spp. | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>Do not graze or harvest treated soybean forage, straw or hay for livestock feed.</li> <li>Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program use a minimum of 2.56 fl oz/A (0.02 lb ai/A).</li> <li>Do not apply within 45 days of harvest. Do not apply more than 0.06 lb ai (0.48 pints) per acre per season.</li> </ul> For control of the first and second instar only.<br>*Suppression only.<br>*See resistance statement under General Directions for Use. |
|                       | Armyworm <sup>1</sup><br>Fall Armyworm <sup>1</sup><br>Yellow-striped Armyworm <sup>1</sup><br>Tobacco Budworm <sup>1</sup><br>Webworm Spp.<br>European Corn Borer<br>Silverspotted Skipper<br>Japanese Beetle (Adult)<br>Blister Beetle Spp.<br>Stink Bug Spp.<br>Plant Bug Spp.<br>Grasshopper Spp.   | 0.025-0.03  | 3.20-3.84 |  |
|                       | Beet Armyworm <sup>1,2</sup><br>Soybean Looper <sup>2,3</sup><br>Lesser Cornstalk Borer <sup>2</sup><br>Spider Mite Spp. <sup>4</sup>   | 0.03        | 3.84      |  |

| SPRAY RECOMMENDATIONS         |                                    |             |   |  |
|-------------------------------|------------------------------------|-------------|---|--|
| Crop                          | Target Pests                       | Rate        |   | Remarks  |
|                               |                                    | lb ai/A     | fl oz/A   |  |
| Sunflower                     | Sunflower Beetle                   | 0.015-0.025 | 1.92-3.20   | <ul style="list-style-type: none"> <li>• Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>• Apply with ground or air equipment using sufficient water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>• Do not apply within 45 days of harvest. Do not apply more than 0.12 lb ai (0.96 pints) per acre per season. Do not apply more than 0.09 lb ai (0.72 pints) per acre per season after bloom initiation. Do not apply as an ultra low volume (ULV) spray.</li> </ul> |
|                               | Cutworm Spp.                       |             |   |  |
|                               | Sunflower Moth                     | 0.02-0.03   | 2.56-3.84   |  |
|                               | Banded Sunflower Moth              |             |   |  |
|                               | Fall Armyworm <sup>1</sup>         |             |   |  |
|                               | Woollybear Caterpillar             |             |   |  |
|                               | Spotted Cabbage Looper             |             |   |  |
|                               | Painted Lady (Thistle) Caterpillar |             |   |  |
|                               | Seed Weevil (Adult)                |             |   |  |
|                               | Stem Weevil (Adult)                |             |   |  |
| Head-Clipper Weevil (Adult)   |                                    |             |   |  |
| Japanese Beetle (Adult)       |                                    |             |   |  |
| Sunflower Maggot (Adult)      | 0.03                               | 3.84        | <ul style="list-style-type: none"> <li><sup>1</sup>For control of first and second instar only.</li> <li><sup>2</sup>Suppression only.</li> <li><sup>3</sup>See resistance statement under General Directions for Use.</li> </ul> |  |
| Leafhopper Spp.               |                                    |             |   |  |
| Meadow Spittlebug             |                                    |             |   |  |
| Stinkbug Spp.                 |                                    |             |   |  |
| Grasshopper Spp.              |                                    |             |   |  |
| Beet Armyworm <sup>1,2</sup>  |                                    |             |   |  |
| Spider Mite Spp. <sup>3</sup> |                                    |             |   |  |

22 7 25

| SPRAY RECOMMENDATIONS           |  |             |           |   |
|---------------------------------|--|-------------|-----------|---|
| Crop                            | Target Pests   | Rate        |           | Remarks   |
|                                 |  | lb ai/A     | fl oz/A   |   |
| Tomato and Tomatillo            | Cabbage Lopper<br>Cutworm Spp.<br>Homworm Spp.   | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>Do not apply within 5 days of harvest. Do not apply more than 0.36 lb ai (2.88 pints) per acre per season.</li> <li>Do not use on varieties in which the mature tomatoes will be less than one inch in diameter (such as cherry tomatoes).</li> <li><sup>1</sup>For control of first and second instar only.</li> <li><sup>2</sup>Suppression only.</li> <li><sup>3</sup>See resistance statement under General Directions for Use.</li> </ul>   |
|                                 | Tomato Fruitworm<br>Tobacco Budworm <sup>1</sup><br>Tomato Pinworm<br>Beet Armyworm <sup>1,2</sup><br>Southern Armyworm <sup>1</sup><br>Yellow-striped Armyworm <sup>1</sup><br>Fall Armyworm <sup>1</sup><br>European Corn Borer<br>Leafminer Spp. <sup>2</sup><br>Colorado Potato Beetle <sup>3</sup><br>Flea Beetle Spp.<br>Grasshopper Spp.<br>Leafhopper Spp.<br>Aphid Spp. <sup>1,2</sup><br>Whitefly Spp. <sup>1,2</sup><br>Meadow Spittlebug<br>Stink Bug Spp.<br>Plant Bug Spp. | 0.02-0.03   | 2.56-3.84 |   |
| Wheat, Wheat Hay, and Triticale | Cutworm Spp.<br>Army Cutworm   | 0.015-0.025 | 1.92-3.20 | <ul style="list-style-type: none"> <li>Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.</li> <li>Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.</li> <li>For chinch bug control, repeat applications at 3- to 5-day intervals if needed. WARRIOR Insecticide may only suppress heavy infestations and/or migrations.</li> <li>Greenbug is known to have many biotypes. WARRIOR Insecticide may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.</li> <li>Do not apply within 30 days of harvest.</li> <li>Do not apply more than 0.06 lb ai (0.48 pints) per acre per season.</li> <li><sup>1</sup>Best control is obtained before insects begin to roll leaves. Once wheat has started to boot, WARRIOR Insecticide may provide suppression only. Higher rates and increased coverage will be necessary.</li> <li><sup>2</sup>See resistance statement under General Directions for Use.</li> </ul> |
|                                 | Armyworm<br>Fall Armyworm<br>Yellow-striped Armyworm<br>Flea Beetle Spp.<br>Cereal Leaf Beetle<br>Stink Bug Spp.<br>English Grain Aphid <sup>1</sup><br>Russian Wheat Aphid <sup>1</sup><br>Bird Cherry-Oat Aphid <sup>1</sup><br>Grasshopper Spp.   | 0.02-0.03   | 2.56-3.84 |   |
|                                 | Grass Sawfly   | 0.025-0.03  | 3.20-3.84 |   |
|                                 | Chinch Bug<br>Greenbug <sup>2</sup>  | 0.03        | 3.84      |   |

**RATE CONVERSION CHART**

| <b>LB AI<br/>PER ACRE</b> | <b>FL OZ<br/>PER ACRE</b> | <b>PINTS<br/>PER ACRE</b> | <b>TREATED ACRES<br/>PER GALLON</b> |
|---------------------------|---------------------------|---------------------------|-------------------------------------|
| 0.015                     | 1.92                      | 0.12                      | 66                                  |
| 0.02                      | 2.56                      | 0.16                      | 50                                  |
| 0.025                     | 3.20                      | 0.20                      | 40                                  |
| 0.03                      | 3.84                      | 0.24                      | 33                                  |
| 0.04                      | 5.12                      | 0.32                      | 25                                  |

AMBUSH®, WARRIOR®, E-Z HANDLER® and KARATE® are trademarks of a ZENECA Group Company.  
 Ammo®, Capture®, Fury®, Mustang® and Pounce® are trademarks of FMC Corporation.  
 Asana® is a trademark of E. I. du Pont de Nemours & Company.  
 Baythroid® is a trademark of Mobay Corporation.  
 Danitol® is a trademark of Sumitomo Chemical Co., LTD.  
 DECIS® and SCOUT-XTRA® are registered trademarks of Hoechst Schering AgrEvo S.A. de C.V.  
 SynerGin™ is a trademark of AgrEvo USA Company.

©1998 ZENECA Inc.

24 7 25

Made in U.S.A.

**ZENECA** Ag Products

**ZENECA Inc.**  
Wilmington, Delaware 19850-5458

CR1098



25 9 25

**RESTRICTED USE PESTICIDE**  
**Due to Toxicity to Fish and Aquatic Organisms**  
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



*For Use to Control Certain Insects on Listed Crops*

**KEEP OUT OF REACH  
OF CHILDREN  
DANGER  
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**ACTIVE INGREDIENT:**

Lambda-cyhalothrin  
[1 $\alpha$ (S\*),3 $\alpha$ (Z)]-(s)-cyano-(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate . . . . . 13.1%

**INERT INGREDIENTS:** . . . . . 86.9%

**TOTAL** . . . . . 100.0%

WARRIOR<sup>®</sup> Insecticide contains 1 pound of active ingredient per gallon and is an emulsifiable concentrate.

EPA Reg. No. 10182-96

**STATEMENT OF PRACTICAL TREATMENT**

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.  
**IF SWALLOWED:** Do not induce vomiting. Call a physician immediately.  
**IF INHALED:** Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.  
**IF IN EYES:** Hold eyelids open and flush with steady, gentle stream of water. Remove any contact lenses and repeat flushing procedure for several minutes. Get medical attention if irritation persists.  
**NOTE TO PHYSICIAN:** Induced vomiting as first aid for this substance may result in increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent. Vomiting should be induced only under professional supervision.

**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).**  
**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.**

See attached booklet for complete Directions for Use.

CR1098.

**ZENECA**