

Reg # 10182-98 p. 1/5
PM 10

[Box Label - Front Panel]

"MOUSER"

READY FOR USE BAIT BOX

ACCEPTED
NOV 4 1986
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10182-98

- Contains enough product to kill up to 10 mice.
- Kills warfarin-resistant mice
- No bait handling
- Effective in damp conditions
- Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT:

Brodifacoum-	
3-[5-(4'-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one.	0.005%
Inert Ingredients.	99.995%
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Read additional precautionary statements on side panel.

ICI Americas Inc.	EPA Est. No.
Agricultural Chemicals Div.	EPA Reg. No. 10182-
Wilmington, DE 19897	Net Contents: 1/20 gm Bait Blok

This product is protected by US patent numbers 3957824 and 4035505 which are the property of Ward Blenkinsop and Company Limited

Made in U.K.

[Box Label - Back Panel]

PRECAUTIONARY STATEMENTS, HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

CAUTION

Harmful if swallowed. Keep away from humans, domestic animals and pets.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI Americas Inc. 302/575-3000.

IF BAIT IS EATEN BY ANIMALS OR PETS, call the National Animal Poison Control Center 217/333-3611 or ICI Americas Inc. 302/575-3000 for 24-hour emergency assistance.

NOTE TO PHYSICIAN OR VETERINARIAN:

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

STORAGE AND DISPOSAL:

Storage: Store only in original container, in a dry place inaccessible to children and pets.

Disposal: Do not reuse empty container. Securely wrap in newspaper and discard in trash. Do not allow children to play with empty baiters.

NOTICE TO BUYER AND USER:

Seller warrants that this product conforms to the chemical description of the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

c 3/5

[Box Label - Side Panel]

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS, AND NONTARGET WILDLIFE. For control of house mice in and around homes, industrial, commercial, agricultural and public buildings. "MOUSER" may also be used in and around transport vehicles (ships, trains, aircrafts) and related port or terminal buildings. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

APPLICATION DIRECTIONS:

Place "MOUSER" at points of mouse activity. Place along walls, in corners, and concealed places. Mice will enter more readily when the box is placed against a wall or other vertical surface. Check the "MOUSER" after 3 days for signs of feeding (mouse droppings or consumption of block visible through transparent bottom). Stations not touched after 3 days should be moved to a new location where mouse activity is high. One "MOUSER" could be sufficient for a small domestic infestation. Where mice are active in more than one place and for large areas, use one "MOUSER" for every 24 sq. yards of floor space. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water and harborage as much as possible. Where a continuous source of infestation is present, establish a permanent bait program and replenish bait as needed.

[Mouse Station Label - Front Panel]

"MOUSER" Ready for Use Bait Box

Kills Mice

Active Ingredient:

Brodifacoum-	
3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-	
naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one.	0.005%
Inert Ingredients.	99.995%
TOTAL	<u>100.000%</u>

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Read additional precautionary statements on back panel.

[Mouse Station Label - Back Panel]

PRECAUTIONARY STATEMENTS, HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

CAUTION

DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS, AND NONTARGET WILDLIFE.

Harmful if swallowed. Keep away from humans, domestic animals and pets.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI Americas Inc. 302/575-3000.

IF BAIT IS EATEN BY ANIMALS OR PETS, call the National Animal Poison Control Center 217/333-3611 or ICI Americas Inc. 302/575-3000 for 24-hour emergency assistance.

NOTE TO PHYSICIAN OR VETERINARIAN:

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

STORAGE AND DISPOSAL:

Storage: Store only in original container, in a dry place inaccessible to children and pets.

Disposal: Do not reuse empty container. Securely wrap in newspaper and discard in trash. Do not allow children to play with empty baiters.

DIRECTIONS FOR USE:

Refer to label on box for full use directions.

ICI Americas Inc.
Agricultural Chemicals Div.
Wilmington, DE 19897

EPA Est. No.
EPA Reg. No. 10182-
Net Contents: 1/20 gm Bait Blok
Made in U.K.

G1/120585elp15