

PM 25
10182-103

page 1 of 3

114501387

10182 103

RESTRICTED USE PESTICIDE
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

Conversion Table
GRAMOXONE SUPER Rate/ha to be Applied

Rate/ha	Rate/lb/a	Rate/kg/a	Rate/gal
475	71	083	18
275	197	125	12
1 1/3	213	259	6
1 1/2	240	281	5.3
2	320	375	4
2 2/3 (1 1/3 qt)	427	500	3
4	640	750	2
5 1/3 (2 2/3 qt)	853	1000	1.5



SUPPLEMENTAL PAMPHLET
DIRECTIONS FOR USE
READ WORKER SAFETY RULES

 **ICI Americas Inc.**
Agricultural Chemicals Division
Wilmington, Delaware 19897

EPA Reg No 10182-103



POISON

**DO NOT POUR FROM ORIGINAL CONTAINER
EXCEPT FOR IMMEDIATE USE.**

CLOVER

DORMANT SEASON TREATMENT (Established Pastures) - For use in Washington, Oregon, Colorado, Idaho, Montana, Utah, Nevada, Wyoming, and California (counties of Del Norte, Siskiyou, Madoc, Shasta, Lassen, Plumas, Sierra and Nevada). Apply for Deciduous or Ryegrass, Bluegrass, Cheatgrass, Dogfennel, Chickweed, and Trifolium in Dormant Clover, except Annual Clover.

Apply 2 1/2 to 4 pts GRAMOXONE SUPER Herbicide per acre (use high rate for cheatgrass) in 20 to 100 gals spray mix by ground as a broadcast application. Apply by air in 5 to 10 gals spray mix per acre. Apply to established stands after the last fall cutting when the crop is dormant but before spring growth starts. Do not apply if regrowth following the last fall cutting is more than 2" tall. Clover foliage present at the time of application will be burned. Apply when the broadleaf weeds and grasses are succulent and growth is from 1" to 6" tall. Add Spreader (non-ionic) at 8 to 32 oz / 100 gals spray mix. Do not pasture animals in treated fields before first cutting. Do not graze, cut or harvest within 60 days of application for clover. Do not apply more than once per season.

Insert
Copy A
Attached



CORN

PRE-PLANT OR PRE-EMERGENCE TREATMENT.

Apply 1 1/2 to 2 1/2 qts GRAMOXONE SUPER Herbicide per sprayed acre as a band treatment over the crop row, or as a broadcast treatment, prior to, during, or after planting - but before emergence of the crop. Aerial Application - 5 to 10 gals water per acre. Broad Application - Use 20 to 60 gals, diluted spray. Add Spreader (non-ionic) at 8 to 32 oz per 100 gals of diluted spray. Seedbeds should be formed as far ahead of planting and treatment as possible to permit maximum weed and grass emergence. Seeding should be done with a minimum amount of soil disturbance. Weeds and grasses emerging after application will not be controlled. Crop plants emerged at time of application will be killed.

TANK MIX RECOMMENDATIONS - For Kill of Existing Vegetation and Residual Weed Control Where Cans Will Be Planted Directly into a Cover Crop, Established Sod, or in Previous Crop Residues. Tank-Mix with Atrazine.

Apply 1 1/2 to 2 1/2 pts GRAMOXONE SUPER Herbicide plus 2 1/2 to 3 1/2 lbs Atrazine 80 Wettable per acre, as a broadcast spray. Refer to Atrazine 80 Wettable label for directions, limitations and cautions. Apply 20 to 60 gals diluted spray per acre. Add Spreader (non-ionic)

at 8 oz per 100 gals diluted spray. Add Atrazine to spray tank first, mixing thoroughly in water. Then add GRAMOXONE SUPER Herbicide and surfactant. Constant agitation recommended at all times.

TANK-MIX WITH AATREX® 80 W AND PRINCEP® 80 W.

Apply 1 1/2 to 2 1/2 pts GRAMOXONE SUPER Herbicide plus 1 1/2 to 2 1/2 lbs PRINCEP 80 W plus 1 1/2 to 2 1/2 lbs AATREX 80 W per sprayed acre prior to, during, or after planting - but before crop emergence. Use the low rates of PRINCEP 80 W and AATREX 80 W on light soils, and the higher rates on heavy clay soils. Use 20 to 60 gals of diluted spray per sprayed acre. Add Spreader (non-ionic) at 8 oz per 100 gals diluted spray. Add the AATREX 80 W to the spray tank while agitating, then add the PRINCEP 80 W, GRAMOXONE SUPER Herbicide, and Spreader (non-ionic). Continue agitation during application. Refer to the AATREX 80 W and PRINCEP 80 W labels for specific precautionary statements.

TANK-MIX WITH BLADEX® 80 W.

Apply 1 1/2 to 2 1/2 pts GRAMOXONE SUPER Herbicide plus 1 1/2 to 5 lbs BLADEX 80 W, depending on soil type, as a broadcast spray. Apply 20 to 60 gals diluted spray per acre. Add Spreader (non-ionic) at 1 qt per 100 gals diluted spray. Add BLADEX 80 W to spray tank first, mixing thoroughly in water. Then add GRAMOXONE SUPER Herbicide and surfactant. Constant agitation recommended at all times. Do not mix with liquid fertilizer. Refer to BLADEX 80 W label for directions, limitations and cautions.

TANK-MIX WITH LASSO® AND ATRAZINE.

Apply 1 1/2 to 2 1/2 pts GRAMOXONE SUPER Herbicide (0.25 to 0.5 lb active) plus 4 to 6 pts LASSO (2.0 to 3.0 lbs active) plus 1.25 to 2.5 lbs Atrazine 80 Wettable (1.0 to 2.0 lbs active) per treated acre prior to, during, or after planting, but before crop emergence. Use 20 to 60 gals of diluted spray per treated acre. Add Spreader (non-ionic) at 8 oz per 100 gals of diluted spray. Add the Atrazine to the spray tank while agitating, then add the LASSO, GRAMOXONE SUPER Herbicide and Spreader (non-ionic). Continue agitation during application. Refer to Atrazine and LASSO labels for cautions and for specific rates for a given soil type.

TANK-MIX WITH DUAL® AND AATREX.

Apply 1 1/2 to 2 1/2 pts GRAMOXONE SUPER Herbicide per acre plus DUAL plus AATREX as a broadcast ground spray before, during, or

BEST AVAILABLE COPY

3 of 3

Apply 1 1/2 to 2 1/2 pints of GRAMOXONE SUPER per acre. Always add a nonionic surfactant at one pint per 100 gallons of diluted spray. Apply in a minimum of 10 gallons per acre by ground or 3 gallons per acre by air to obtain thorough coverage. For best results by ground, use flat-fan nozzles spaced at 20 inch intervals with boom height adjusted to obtain a 30% spray pattern overlap. For improved emerged weed control or extended weed control GRAMOXONE SUPER may be tank-mixed with other registered herbicides. Refer to the label(s) of each herbicide for directions, limitations, cautions and for a list of weeds controlled.