

RESTRICTED USE PESTICIDE
Due to Acute Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

ACCEPTED
JAN 22 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10182-111

Inside of Front Cover

CYCLONE® Herbicide
A Weed, Grass, and Harvest Aid Herbicide
COMPLETE DIRECTIONS FOR USE

96 NOV 29 P 3:05

RECD EPA/119/DPD1

CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL**

BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

TABLE OF CONTENTS

	Page
STATEMENT OF PRACTICAL TREATMENT	
PRECAUTIONARY STATEMENTS	
Hazards to Humans and Domestic Animals	
Personal Protective Equipment	
Environmental Hazards	
DIRECTIONS FOR USE	
Agricultural Use Requirements	
Storage & Disposal	
General Information	
Rainfastness	
Application	
Use of a Nonionic Surfactant or Crop Oil Concentrate	
Nozzle Selection	
Spray Carrier	
Spray Volume	
Drift Minimization	
Application Timing	
Burndown of Grass Cover Crops or Volunteer Cereals	
Environmental Conditions	
Spot Spraying	
Tank Mixing For Improved Burndown	
Limitations, Precautions and Warnings	
SPECIFIC USE RECOMMENDATIONS	
Alfalfa	
Almonds	
Asparagus	
Beans, Dry (Harvest-Aid)	
Cassavas, Taniars & Yams	
Chemical Fallow	
Clover and Other Legumes	
Corn	
Cotton	
Easter Lilies	
Fallow Land	
Grasses (for seed)	
Guar	
Guava	
Hops	
Mint	
Onions (seeded) and Garlic	
Passion Fruit	
Peanuts	
Pigeon Peas	
Pineapple	
Potato	
Resin Soaking	
Safflower	
Small Fruits	
Small Grains	

Sorghum (grain)
Soybeans
Sugar Beets
Sugarcane
Sunflower
Taro, Dryland (Hawaii Only)
Trees & Vines
Tyfon
Vegetables
Non-Agricultural Use Requirements

OTHER USES

Conservation Reserve, Federal Set-Aside, Conservation Compliance
Noncrop Uses
Pasture Reseeding

Conversion Table

Storage & Disposal

KEEP OUT OF REACH OF CHILDREN

O
DANGER POISON
X
PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

~~ONE SWALLOW CAN KILL!~~

~~SYMPTOMS ARE PROLONGED AND PAINFUL. ONSET OF SYMPTOMS MAY BE
DELAYED FOR UP TO 3 DAYS AFTER SWALLOWING.~~

- NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.
- ~~IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN STATEMENT OF
PRACTICAL TREATMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.~~
- DO NOT USE OR STORE IN OR AROUND THE HOME.
- DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.

ACTIVE INGREDIENT:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride) 30.3%

INERT INGREDIENTS: 69.7%

Total 100.0%

Contains 2 lbs. paraquat cation per gallon.
Contains stench (odor) and emetic.

EPA Reg. No. 10182-111

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: IMMEDIATELY INDUCE VOMITING by touching back of throat with finger. Administer fluids and **Drink 1 or 2 glasses of water** and induce further vomiting. **If person is unconscious do not give anything by mouth and do not induce vomiting. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician or Poison Control Center and GET TO A HOSPITAL OR PHYSICIAN MEDICAL FACILITY FAST. TAKE CONTAINER AND LABELING WITH YOU. PROMPT TREATMENT IS ESSENTIAL AND MUST BE INITIATED BEFORE SIGNS AND SYMPTOMS APPEAR. SEE NOTE TO PHYSICIANS.**

IF INHALED: IMMEDIATELY Remove victim to fresh air. or get away from spray mist. Exposure may cause irritation, nose bleeds, and may lead to poisoning. Stop and check spray procedure. **If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.** The odor of this product is from the stenching agent which has been added, not from paraquat.

IF IN EYES: IMMEDIATELY wash eyes with **Hold eyelids open and flush with a steady, gentle stream of water** for at least 15 minutes, and get medical attention. Contact may cause eye injury. **CALL A PHYSICIAN. Get medical attention.**

IF ON SKIN: IMMEDIATELY Wash with **plenty of soap and water.** Prolonged contact will cause severe irritation. Contact with irritated skin or a cut or repeated contact with intact skin may result in poisoning. **GET MEDICAL ATTENTION. Get medical attention.**

Read and follow **PRECAUTIONARY STATEMENTS:**

NOTE TO PHYSICIANS: CALL ZENECA'S MEDICAL EMERGENCY INFORMATION NETWORK AT 1-800-327-8633 (Dial 1-800-F-A-S-T-M-E-D) at any hour to obtain toxicology and medical management consultation, a supply of bentonite clay, and paraquat analysis analyses. **Prompt treatment is essential and must be initiated immediately before signs and symptoms appear. Symptoms are prolonged and painful and may be delayed for days after swallowing. Symptoms following ingestion may be delayed up to 3 days. To be effective, treatment for paraquat poisoning must begin IMMEDIATELY. Treatment consists of may include binding paraquat in the gut with suspensions of bentonite clay or charcoal and/or removal of paraquat from the blood by prolonged charcoal hemoperfusion or continuous hemodialysis. Probable mucosal damage may contraindicate the use of gastric lavage.**

FOR 24 HOUR EMERGENCY MEDICAL ASSISTANCE, CALL ZENECA'S MEDICAL EMERGENCY INFORMATION NETWORK AT 1-800-(327-8633) (1-800-F-A-S-T-M-E-D) FOR 24 HOUR EMERGENCY MEDICAL ASSISTANCE.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident Call CHEMTREC AT 1-800-424-9300 FOR A CHEMICAL EMERGENCY such as spill, leak, fire or accident.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

O
DANGER POISON
X

~~FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. CAUSES IRREVERSIBLE EYE DAMAGE. CAUSES SKIN IRRITATION. MAY CAUSE ALLERGIC SKIN REACTION.~~

~~DO NOT get in eyes, on skin, or on clothing. DO NOT breathe spray mist. WASH SPLASHES from skin and eyes immediately.~~

~~May be fatal if swallowed. Fatal if inhaled. Harmful if absorbed through skin. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Do not breathe spray mist. Avoid contact with skin.~~

~~IMPORTANT: Inhalation is an unlikely route of exposure due to low vapor pressure and large spray droplet size, but mucosal irritation or nose bleeds may occur. Prolonged contact with this concentrated product can irritate your skin.~~

Personal Protective Equipment

Applicators and other handlers (other than Mixers and Loaders) must wear:

- Long-sleeve shirt and long pants
- Waterproof gloves
- Shoes plus socks
- ~~Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)~~

Mixers and Loaders must wear:

- Long-sleeve shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Face shield
- Chemical-resistant apron

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

WILDLIFE: This product is toxic to wildlife. ~~Birds and other wildlife in treated areas may be killed.~~ Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DRIFT: CYCLONE® herbicide is a contact herbicide that kills ~~desiccates~~ all green plant tissue. Do not apply under conditions involving possible drift to food, forage, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use, or consumption. Do not apply when weather conditions favor drift from treated areas. To avoid drift, do not make aerial applications during periods of thermal inversion. Refer to the local state laws, regulations, and guidelines for proper application to avoid off-target movement.

CHEMICAL HAZARD: This product reacts with aluminum to produce hydrogen gas which may form a highly combustible gas mixture. ~~Such gas could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette, or other ignition source.~~ Do not mix or store in containers, spray tanks, nurse tanks, or such systems made of aluminum or having aluminum fittings. ~~After dilution with water to spray concentrations, spray systems made of, or containing aluminum parts, may be used but should not be used for extended (overnight) storage.~~

EQUIPMENT/CONTAINER: Flush all spray equipment with water after use each day. CYCLONE is corrosive to aluminum. Aluminum spray equipment and aluminum aircraft structures that are exposed to spray solution or spray drift should be flushed thoroughly with water immediately after use.

DIRECTIONS FOR USE

It is a violation of Federal law and may endanger the applicator or third parties to use this product without complying with all label directions and warnings. ~~In a manner inconsistent with its labeling.~~ Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT USE AROUND HOME GARDENS, SCHOOLS, RECREATIONAL PARKS, GOLF COURSES OR PLAYGROUNDS.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), ~~notification to workers~~, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

For Preplant or Preemergence (Broadcast or Banded), Chemical Fallow, Postemergence Directed Spray, Early Postemergence Broadcast in Peanuts and Dormant Season Applications, and "Between Cutting" Applications in Alfalfa: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

For Harvest Aid and Desiccation Applications: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 ~~24~~ hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- ~~Coveralls over long-sleeved shirt and long pants~~
- Waterproof gloves
- ~~Chemical-resistant footwear plus socks~~
- ~~Chemical-resistant headgear for overhead exposure~~

~~Shoes plus socks~~
~~Protective eyewear~~

~~Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.~~

~~READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. READ STATEMENT OF PRACTICAL TREATMENT AND PRECAUTIONARY STATEMENTS ON PAGES _____~~

~~NOTE: DO NOT USE AROUND HOME GARDENS, SCHOOLS, RECREATIONAL PARKS, GOLF COURSES, OR PLAYGROUNDS.~~

~~DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.~~

~~IMPORTANT: The effect of swallowing CYCLONE is far more severe than that which usually results from skin contact or inhalation of spray mist. This product should not cause injury if used according to label instructions. Follow the label as if your life depends on it.~~

GENERAL INFORMATION

CYCLONE herbicide is a contact herbicide used to control or suppress a broad spectrum of emerged weeds. CYCLONE herbicide controls most small annual weeds – both broadleaves and grasses, and suppresses perennial weeds by destroying green foliage. CYCLONE herbicide can also be used as a crop desiccant at harvest.

CYCLONE herbicide is formulated as a liquid which contains 2 lbs. of active ingredient per gallon. The formulation contains a nontoxic odor and an emetic (an agent which will induce vomiting if the product is swallowed). The odor is included in the formulation to help prevent accidental ingestion of CYCLONE herbicide.

CYCLONE herbicide is rapidly absorbed by green plant tissue and interacts with the photosynthetic process to produce superoxides which destroy the plant cells. CYCLONE herbicide requires actively growing green plant tissue to function. Thorough coverage of all green foliage is essential for effective weed control. CYCLONE is not as effective on drought-stressed weeds, weeds with little green foliage (i.e., mowed or cut weeds), or mature woody bark of trees and vines.

Clay and organic matter rapidly tie up CYCLONE herbicide. As a result, CYCLONE herbicide has no residual soil activity to affect later-planted crops or later germinating weeds.

RAINFASTNESS

Because CYCLONE herbicide is rapidly absorbed by the weed foliage, rain occurring 30 minutes or more after application will have no effect on the activity of CYCLONE herbicide.

APPLICATION

Since CYCLONE herbicide is a contact-type herbicide, it is essential to obtain complete coverage of target weeds to get good control. Improper application technique and/or application to large, stressed, or mown weeds will usually result in unacceptable weed control. Good coverage is also essential for proper performance in harvest-aid applications. See details below for specific application instructions. ~~Do not apply this product through any type of irrigation system.~~

USE OF A NONIONIC SURFACTANT OR CROP OIL CONCENTRATE

ALWAYS ADD ONE OF THE FOLLOWING:

Failure to add one of the following will result in reduced performance of CYCLONE herbicide.

Nonionic Surfactant: Add nonionic surfactant containing 75% or more surface-active agent at 0.125% v/v (1 pint per 100 gallons), OR add a nonionic surfactant containing 50-74% surface-active agent at 0.25% v/v (2 pints per 100 gallons), of the finished spray volume for ground applications. For aerial applications, add a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons) of the finished spray volume.

Crop Oil Concentrate: Add a non-phytotoxic crop oil concentrate containing 15-20% approved emulsifier, at 1.0% v/v (1 gallon per 100 gallons) of the finished spray volume for ground applications. For aerial applications, add 1 pint of crop oil concentrate per acre. Do not use crop oil concentrate when using CYCLONE for cotton harvest aid.

NOZZLE SELECTION

The use of flat-fan nozzles will result in the most effective application of CYCLONE herbicide. Flood nozzles are generally not as good as flat fans since they produce large uneven droplets. The use of flood nozzles may result in reduced weed control due to inadequate coverage.

WHEN SPRAYING LESS THAN 20 GALLONS OF SPRAY CARRIER PER ACRE, USE ONLY FLAT FAN NOZZLES AS RECOMMENDED IN THE CHART BELOW.

RECOMMENDED NOZZLES, PRESSURES AND SETUP		
	Nozzle Type	
	Flat Fan	Flood
Maximum Size	8	15
Spray Pressure (at nozzle)	30-50 P.S.I.	30-50 P.S.I.
Maximum Nozzle Spacing	30"	40"
Direction of Spray Pattern	Down	Down
Maximum Speed	10 mph	10 mph
Spray Overlap (at each edge)	30%	50%

(Typesetter: Insert nozzle illustrations here)

Using nozzles, pressures, or setups different from the above chart will result in reduced control.

SPRAY CARRIER

Always use clean water (free of mud or clay), clear liquid nitrogen, or complete clear liquid fertilizers as the carrier when spraying CYCLONE herbicide. Muddy water, or suspension-type fertilizers containing clay, can inactivate CYCLONE herbicide. Never use suspension-type fertilizers containing clay as the spray carrier. If using a complete clear liquid fertilizer containing high phosphate levels as the spray carrier, always use the higher rate of CYCLONE and surfactant.

NOTE: When using liquid fertilizers such as 28% N as a carrier, it is important that nonionic surfactant still be used with CYCLONE herbicide. Liquid fertilizer carriers cannot substitute for surfactant.

RATES OF CYCLONE

Follow recommended rates listed with each use of CYCLONE herbicide. Use the higher label rates when weeds are dense or large. Also, use higher label rates for harvest aid when crop vegetation is dense.

SPRAY VOLUME

Follow recommended minimum spray volumes listed with each use of CYCLONE herbicide. These are minimum volumes only, and spray volumes should be increased as necessary to obtain complete coverage of the target weed or plant without runoff from the foliage.

WHEN SPRAYING LESS THAN 20 GALLONS OF SPRAY CARRIER PER ACRE, TARGET WEEDS SHOULD NOT EXCEED 6 INCHES IN HEIGHT.

DRIFT MINIMIZATION

DO NOT spray if conditions of thermal inversion exist, or if wind direction and speed may cause spray drift onto adjacent areas. With both aerial and ground applications, the use of a drift control or spray deposition additive (in accordance with label directions) may reduce the potential for drift to occur. Drift minimization is the responsibility of the applicator. Refer to the local state laws, regulations, and guidelines for proper application to avoid off-target movement.

When CYCLONE is applied at less than 10 gallons per acre finished spray volume, a drift control or spray deposition additive **SHOULD** be used. Refer to the additive label for use directions.

APPLICATION TIMING

CYCLONE herbicide should be applied to emerged weeds when they are small. Weeds 1 to 6" in height are the easiest to control. Larger weeds may be more difficult to control. When weeds have been grazed or mowed, thus removing much of the green foliage, allow the weeds to regrow to a height of 2 to 4 inches before spraying if possible. Similarly, when forage or grain crops have been harvested prior to spraying, weeds present in the field will also have been cut. To allow for adequate green foliage to remain on weeds in this situation, raise cutter bars as high as possible from the ground to cut stubble and weeds at a greater height.

BURNDOWN OF GRASS COVER CROPS OR VOLUNTEER CEREALS

When using CYCLONE herbicide for control of grass cover crops or volunteer cereals, best results are obtained when CYCLONE herbicide is applied prior to tillering or after boot stage. This is especially important with a wheat cover crop or volunteer wheat. Treatments made between tillering and boot stage will generally not provide complete control. Do not expect complete control of perennial cover crops.

ENVIRONMENTAL CONDITIONS

CYCLONE herbicide is active over a wide range of environmental conditions. Cool weather (below 55°) will slow the activity of CYCLONE herbicide, as will cloudy, overcast weather, but will not affect performance.

In dry areas, dust stirred up by high winds or equipment tires can coat weed leaves and reduce CYCLONE herbicide activity. Avoid applying CYCLONE herbicide in extremely dusty conditions.

SPOT SPRAYING

When only small areas are to be sprayed with labeled applications, it is advantageous to mix small quantities of CYCLONE herbicide. To aid in mixing small quantities, the following table should be consulted.

If The Broadcast Rate Per Acre for CYCLONE is:	Add The Following Amount of CYCLONE To 1 Gallon of Water
1 pt.	3/8 fl. oz.
2 pts.	1/2 fl. oz.
2 1/4 pts.	3/4 fl. oz.
4 pts.	1-1/4 fl. oz.

Always add 1/3 to 1/2 fl. oz. of a nonionic surfactant for each gallon of spray. When spot spraying in this manner, spray to thoroughly wet the foliage, but not to the point of runoff.

TANK MIXING FOR IMPROVED BURNDOWN OF DIFFICULT WEEDS AND RESIDUAL WEED CONTROL

Photosynthetic Inhibitor Herbicides

Difficult weeds can often be controlled by tank mixing CYCLONE herbicide with other herbicides. The addition of herbicides which are also photosynthetic inhibitors (PSI) will slow the activity of CYCLONE, allowing CYCLONE herbicide to thoroughly distribute itself within the treated leaf. The resulting level of control is usually greater than if CYCLONE herbicide was applied alone.

CYCLONE herbicide may be applied in tank mixture with the following PSI herbicides:

Aatrex® herbicide
Atrazine herbicide
Bicep® herbicide
Bicep Lite® herbicide
Bladex® herbicide
Canopy® herbicide
Extrazine® herbicide
Gemini® herbicide
Griffex® herbicide
Lariat® herbicide
Lexone® herbicide
Linex® herbicide
Lorox® herbicide
Lorox Plus™ herbicide
Preview® herbicide
Princep® herbicide
Sencor® herbicide

Refer to respective product label(s) for rates of application, directions for use, limitations, cautions and for a list of weeds controlled.

Improved Weed Control With PSIs

Control of difficult weeds listed below and annual grass control will be enhanced by the addition of a PSI herbicide. For best results a second application is needed.

Marestail	Prickly lettuce
Pennsylvania Smartweed	Lambsquarters
Velvetleaf	Morningglory
Malva (Cheeseweed)	Cheatgrass
Fall Panicum	Knotweed
Giant Ragweed	Volunteer wheat
Kochia	Perennial Weeds (suppression only)
Cocklebur	Barnyardgrass
Tansymustard	Broadleaf signalgrass
	Sedges

Improved Control of Perennial and Annual Broadleaf Weeds

When perennial broadleaf weeds such as Canada thistle, bindweed, dandelion, etc. or difficult to control annual broadleaf weeds such as giant ragweed or morningglory are present, tankmixes with 2,4-D ester (Low Volatile), 2,4-DB or Banvel® herbicide, where labeled, will help improve control. Tank mixing the amine formulation of 2,4-D with CYCLONE herbicide may result in reduced grass control.

Order of Tank Mixing

In general, CYCLONE herbicide tankmixes with other products should be mixed as follows:

1. Fill spray tank 1/2 full with clean water or other approved carriers such as clear liquid fertilizer.
2. Begin tank agitation and continue throughout mixing and spraying.
3. Add dry formulations (WP, DF, etc.) to tank.
4. Add liquid formulations (SC, EC, L, etc.) to tank.
5. Add CYCLONE herbicide to tank.
6. Add nonionic surfactant to tank.
7. Fill remainder of spray tank.

Always refer to labels of other pesticide products for mixing directions and precautions which may differ from those outlined here.

Since many of the herbicides listed on this label are available in several types of formulations, it is advisable to perform a jar test to check physical compatibility.

LIMITATIONS, PRECAUTIONS AND WARNINGS

- **DO NOT** use around home gardens, schools, recreational parks, or playgrounds.
- In preplant and preemergence (to the crop) uses, do not apply to soils lacking clay minerals, i.e., peat, muck, pure sand, artificial planting media.
- Seedbeds and plantbeds should be formed as far ahead of planting and treatment as possible to permit maximum weed and grass emergence prior to treatment.
- Seeding or transplanting should be done with a minimum amount of soil disturbance.
- Weeds and grasses emerging after application of CYCLONE herbicide will not be controlled or suppressed.
- Unless otherwise indicated, crop plants emerged at time of application may be severely injured or killed if contacted by sprays of CYCLONE.
- **PRODUCT PERFORMANCE CONCERNS:** Concerns with the performance of CYCLONE herbicide must be reported to the retail dealer or to a ZENECA representative within 14 days of application. Weeds emerging after application or regrowth of perennial weeds do not constitute a legitimate product performance complaint.

SPECIFIC USE RECOMMENDATIONS

The following tables indicate use patterns, rates, minimum spray volumes, preharvest intervals and other precautions, restrictions and comments specific to each crop. Read and follow directions carefully.

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>ALFALFA</u> New seedlings (California only)	Broadcast	See Table __ page __	Ground: 10 gal Air: 5 gal	70	<ul style="list-style-type: none"> • Apply during late winter or early spring. • Do not cut or harvest within 70 days after application. • Do not apply more than once during the first growing season. • Caution: Seedling alfalfa stands will be reduced and replanting may be necessary. • Not recommended for seedling alfalfa grown for seed. • Alfalfa foliage present at time of application will be burned.
<u>ALFALFA</u> (No-till or conventional planting)	Preplant or Preemergence Broadcast or Banded Over-Row	2 - 4 pints (use a minimum of 3 pts. in California)	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Apply prior to emergence of the crop. • Crop plants emerged at time of application will be killed. • Seeding should be done with a minimum amount of soil disturbance.

16/1/1

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p>ALFALFA</p> <p>Dormant season on established plantings</p> <p>Region A - See map at end of Alfalfa section.</p>	Broadcast	2 - 3-1/8 pints	Ground: 10 gal Air: 5 gal	42	<ul style="list-style-type: none"> • For control of weeds, including bluegrass, chickweed, henbit, downy brome, ryegrass, cheatgrass, dogfennel, tansymustard, london rocket, sowthistle, rescue brome, wild oats, and other winter annuals; and suppression of perennial weeds. • Do not apply if <i>fall regrowth</i> following last fall cutting is greater than 6", or if <i>spring regrowth</i> is more than 2 inches. • Apply to well-established stands (at least one year old) after the crop is dormant. • Alfalfa foliage present at the time of application will be burned which may reduce the yield of the first cutting. • Do not cut or harvest within 60 days of application. • Do not apply more than once per season. • Tank mix with metribuzin (Lexone or Sencor) for improved burndown of weed vegetation and residual weed control. Consult the metribuzin product label for a list of weeds controlled, rates of application, and precautions.

17/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p><u>ALFALFA</u></p> <p>Dormant season Tankmix with Velpar® L herbicide</p> <p>Region A - See map at end of Alfalfa section.</p>	Broadcast	1½ - 2½ pints	Ground: 10 gal Air: 10 gal	42	<ul style="list-style-type: none"> • For control of weeds such as chickweed, downy brome and tansymustard. • Use the 1½ pt. rate of CYCLONE when weeds and grasses are less than 4" tall. • Mix with 1 to 2 qts. of Velpar L per acre. • Use the lower rate of Velpar L on loamy sands or sandy loams. Refer to Velpar L label for directions, limitations, cautions and for a list of weeds controlled. • Apply once to established alfalfa stands during the dormant season. • Do not apply if <i>fall regrowth</i> following last fall cutting is greater than 6 inches, or if <i>spring regrowth</i> is more than 2 inches. • Do not apply to alfalfa during the first season after seeding. • Temporary chlorosis may occur on alfalfa regrowth. • Stress which may be caused in part by low fertility, disease, insects, winterkill, over cutting, drought or frost may increase the chances of crop injury. • DO NOT USE on gravelly or rocky soils, exposed subsoils, hardpan, sand or poorly drained alkaline soils as crop injury, including mortality, may result. • Do not cut or harvest within 42 days of application.

18/91

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p>ALFALFA</p> <p>Dormant Season</p> <p>On established plantings: Region B- See map at end of Alfalfa section.</p> <p>On fall-seeded, newly established stands less than 1-year-old: Region A- See map at end of Alfalfa section.</p> <p>On fall-seeded, newly established stands less than 1-year-old: Region B- See map at end of Alfalfa section.</p>	<p>Broadcast</p> <p>Broadcast</p> <p>Broadcast</p>	<p>1 - 2 pints</p> <p>1 - 2 pints</p> <p>0.5 - 1-1/4 pints</p>	<p>Ground: 10 gal Air: 5 gal</p> <p>Ground: 10 gal Air: 5 gal</p> <p>Ground: 10 gal Air: 5 gal</p>	<p>60</p>	<ul style="list-style-type: none"> • For desiccation of weeds, including London rocket, sowthistle, rescue brome, wild oats, chickweed, ryegrass, bluegrass, cheatgrass, dogfennel, tansymustard, henbit, downy brome, and other winter annuals; and suppression of perennial weeds. • Apply during late fall or winter months after the last fall cutting and before first spring cutting. • Do not apply if <i>regrowth</i> after grazing or cutting is more than 2 inches. • Do not harvest within 60 days of application. • CAUTION: Applications to alfalfa that is not dormant, or has broken dormancy, may result in stand and/or yield reductions. Replanting may be necessary. Green alfalfa foliage present at the time of application will be burned. • Total hay yield of first cutting may be reduced in alfalfa fields with severe weed infestation. This reduction will usually be directly proportionate to the loss of weed weight. • Do not apply more than once per season. • Tank mix with metribuzin (Lexone or Sencor) for improved burndown of weed vegetation and residual weed control in <i>dormant established</i> (at least 1-year old) alfalfa. Consult the metribuzin product label for a list of weeds controlled, rates of application, and precautions. • Do not apply tankmix with metribuzin on newly established (less than 1-year old) alfalfa. <p>California</p> <ul style="list-style-type: none"> • For desiccation of weeds including bluegrass, ryegrass, shepherd's-purse, chickweed, tansymustard, foxtail, sowthistle and groundsel. • Use high rate if ryegrass, shepherd's-purse, sowthistle or groundsel is present.

19/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p>ALFALFA</p> <p>Between-cuttings treatment in established plantings. (Includes first year alfalfa) (All states East of the Rocky Mountains)</p> <p>(INSERT MAP BETWEEN THE ALFALFA AND ASPARAGUS TABLES)</p>	Broadcast	1½ pints	Ground: 10 gal	30	<ul style="list-style-type: none"> • Weeds much beyond the seedling stage and the stubble of weeds cut off during harvest will be less affected by this treatment. • Apply immediately after alfalfa has been removed for hay or silage. • Do not treat more than 5 days after cutting. • CAUTION: First year alfalfa stands and yields may be reduced if alfalfa is allowed to regrow more than two (2) inches. • Alfalfa foliage present at time of application will be burned. • In arid areas where moisture is limited, weed control may be reduced. • Do not cut or harvest within 30 days of application. • Make 1 to 3 applications, as needed, during the growing season. These sprays may be applied in addition to a dormant application. • For first year alfalfa, do not apply more than twice during the first growing season.

19/01

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>ALMONDS</u>	Directed Spray	1¼ - 4 pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • Do not allow spray to contact green stems (except suckers) or foliage. • Use a shield or wrap plant when spraying around young trees or vines. • Do not graze treated areas. • Do not feed cover crops grown in treated areas to livestock. • Do not apply when nuts to be harvested are on the ground. • For mature woody weeds, perennial weeds, late germinating weeds and green suckers, retreatment or spot treatments may be necessary.
<u>ASPARAGUS</u>	Preplant or Preemergence Broadcast or Banded Over-Row	2 - 4 pints	Ground: 10 gal Air:* 5 gal	-	<ul style="list-style-type: none"> • Apply prior to emergence of the crop. • Crop plants emerged at time of application will be killed.
<u>ASPARAGUS</u> Preemergence to established plantings at least 2 years old	Broadcast or Band Over-Row	2 - 4 pints	Ground: 10 gal Air: 5 gal	6	<ul style="list-style-type: none"> • Apply prior to emergence of crop or after last harvest. • Crop plants emerged at time of planting will be killed. • May be tank-mixed with Princep herbicide for residual control. Refer to Princep label for specific directions, limitations, cautions and for a list of weeds controlled.

21/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
BEANS, DRY Sweet lupin White sweet lupin White lupin Grain lupin Adzuki beans Asparagus beans Black beans Broad beans Field beans Garbonzo beans Kidney beans Lablab beans Lima beans Moth beans Mung beans Navy beans Pinto beans Rice beans Snap beans Tepary beans Urd beans Wax beans Blackeyed peas Chickpeas Cowpeas	Harvest-Aid	1½ - 2¼ pints	Ground: 20 gal Air: 5 gal	7	<ul style="list-style-type: none"> • Add spreader (nonionic) at 1 quart per 100 gallons of spray mix. • For vining type beans or bush type with lush growth, use a single application of the higher rate. • May also be applied as a split application. DO NOT make more than 2 applications or exceed a total of 1.5 pints per acre. The split application may improve vine coverage. • Apply when the crop is mature and at least 80% of the pods are yellowing and mostly ripe with no more than 40% (<u>bush type beans</u>) or 30% (<u>vine type beans</u>) of the leaves still green in color. • DO NOT apply when weather conditions favor spray drift. A drift control agent may be included to reduce spray drift. • NOT REGISTERED FOR USE ON DRY BEANS IN CALIFORNIA
BEANS, DRY Crowder peas Southern peas Catjang Guar					

22/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
CASSAVAS, TANIERS & YAMS (Puerto Rico only)	Shielded Post Directed Spray	2½ pints	Ground: 50 gal Air: 5 gal	90	<ul style="list-style-type: none"> • Apply when weeds are succulent and growth is 1-6 inches. • On cassavas and taniers, do not make more than 3 applications per crop season. • On yams do not make more than 2 applications per crop season. • Do not allow spray to contact cassavas, tanier or yam plants as injury may result. • Do not spray under windy conditions. • Do not graze treated areas or feed treated forage to livestock.

23/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
CHEMICAL FALLOW General Information			Ground: 5 gal Air: 5 gal See Precautions, Restrictions and Comments		<ul style="list-style-type: none"> • Use higher spray volumes for better coverage as density of stubble, crop residue or weeds increase. • To control volunteer wheat or downy brome, fall-applied treatments generally work best with CYCLONE. If possible, tank mix with Atrazine or Bladex for maximum burndown and residual control. • Apply from immediately after harvest up to emergence of the newly seeded crop as a broadcast or band treatment. • Cut wheat as high as possible to avoid cutting weeds too short, and allow the weeds to grow at least 2-3 inches after harvest before applying CYCLONE. • The addition of Banvel or 2,4-D ester (Low Volatile) may aid in the suppression of emerged perennial broadleaf weeds and large annual broadleaf weeds. • Refer to 2,4-D ester (Low Volatile), Banvel or residual herbicide label(s) for directions, limitations, cautions and for a listing of weeds controlled. • For extended weed control during the fallow period, tankmixes with registered residual herbicide combinations other than those listed on this label are permissible. • Weeds taller than 6 inches may not be controlled. • Weeds and grasses emerging after application will not be controlled. • Crop plants emerged at the time of application will be killed. • By ground application, apply 5-60 gallons of spray mix per acre. If applying at <10 GPA by ground, utilize the following additional precautions: <ul style="list-style-type: none"> - Do not apply with floaters or exceed a speed of 10 MPH - Apply with flat fan nozzles only at 30-40 PSI - Apply only in a tankmix with atrazine at a minimum of 0.5 lbs ai/acre; - By air, apply in 5-10 gallons of spray mix per acre.

24
 / 61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
CHEMICAL FALLOW Continuous Wheat 2-3 Month Recropping Interval	Broadcast	Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts	Ground: 5 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Make application at least 45 days prior to seeding. • Use at least 2 pints of CYCLONE per acre with a PSI (see page __) for volunteer wheat or downy brome control in the spring. • Tank mix with Bladex for enhanced burndown, however, low rates (less than 2 lbs active) of Bladex may not provide complete control of volunteer wheat and downy brome. • Refer to the Bladex label for specific use rates for your soil type, use directions, cautions and a list of weeds controlled. • Refer to the Chemical Fallow General Information section.
CHEMICAL FALLOW Wheat-Fallow- Wheat Rotations Fall applied after harvest (seeded 12-14 months later)	Broadcast	Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts	Ground: 5 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Spray before weeds produce seed. • Volunteer wheat and downy brome control are better with late August or early September applications. • Tank mix with Atrazine and/or Bladex, Marksman® herbicide, or Command® herbicide for enhanced burndown and residual weed control. • Tank mix with Sencor 75DF for burndown and residual control of grass and broadleaf weeds. • Refer to the product labels for specific use rates for your soil type, use directions, cautions and a list of weeds controlled. • Refer to the Chemical Fallow General Information section.

25
 19/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p>CHEMICAL FALLOW</p> <p>Wheat-Fallow- Wheat Rotations Spring applied (seeded 3-5 months later)</p>		<p>Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts</p>			<ul style="list-style-type: none"> • Application should be made March 1 to April 15, prior to spring rains to conserve moisture. • Volunteer wheat is easier to control after the boot stage, but soil moisture loss will be greater. • Use at least 2 pints of CYCLONE per acre with a PSI (see page ___) for volunteer wheat or downy brome control in the spring. • Tank mix with Bladex for enhanced burndown, however, low rates (less than 2 lbs active) of Bladex may not provide complete control of volunteer wheat and downy brome. • Refer to the Bladex label for specific use rates for your soil type, use directions, cautions and a list of weeds controlled. • Refer to the Chemical Fallow General Information section. • Tank mix with Sencor 75DF for burndown and residual control of grass and broadleaf weeds. • Refer to the Sencor 75DF label for use rates for your soil type, use directions, cautions, and weeds controlled.
<p>CHEMICAL FALLOW</p> <p>Wheat-Annual Crop*-Wheat Rotations (Fall applied in wheat stubble)</p>	Broadcast	<p>Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts</p>	<p>Ground: 5 gal Air: 5 gal</p>	-	<ul style="list-style-type: none"> • Tank mix with Atrazine and/or Bladex or Marksman for enhanced burndown and residual weed control. Refer to the labels for specific use rates for your soil type, use directions, cautions, and a list of weeds controlled. • Spray after wheat harvest and before weeds produce seed. If grasses such as foxtails or barnyardgrass recover, respray before they develop seed. • Volunteer wheat and downy brome are easier to control with late August to November applications. • Refer to the Chemical Fallow General Information section.

24/11

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p>CHEMICAL FALLOW</p> <p>Wheat-Annual Crop*-Wheat Rotations (Spring applied prior to planting an annual crop*)</p> <p>* Approved Annual Crops are grain sorghum, corn, wheat, or proso millet.</p>		<p>Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts</p>	<p>Ground: 5 gal Air: 5 gal</p>	<p>-</p>	<ul style="list-style-type: none"> • Tank mix with Atrazine and/or Bladex for enhanced burndown and residual weed control. Refer to the labels for specific use rates for your soil type, use directions, cautions, and a list of weeds controlled. • Use at least 2 pints of CYCLONE per acre with a PSI (see page __) for volunteer wheat or downy brome control in the spring. • Tank mix with Bladex for enhanced burndown, however, low rates (less than 2 lbs active) of Bladex may not provide complete control of volunteer wheat and downy brome. • Follow the Atrazine recommendations pertaining to soil pH and recropping intervals. • Refer to the Chemical Fallow General Information section.

27/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p><u>CLOVER AND OTHER LEGUMES¹</u></p> <p>Dormant Season</p> <p>On established plantings: Region A- See map at end of Alfalfa section.</p> <p>On established plantings: Region B- See map at end of Alfalfa section.</p> <p>On fall-seeded, newly established stands less than 1-year-old: Region A- See map at end of Alfalfa section.</p> <p>On fall-seeded, newly established stands less than 1-year-old: Region B- See map at end of Alfalfa section.</p>	<p>Broadcast</p> <p>Broadcast</p> <p>Broadcast</p> <p>Broadcast</p>	<p>2 - 3-1/8 pints</p> <p>1 - 2 pints</p> <p>1 - 2 pints</p> <p>1/2 - 1-1/4 pints</p>	<p>Ground: 10 gal Air: 5 gal</p>	<p>60</p> <p>60</p> <p>60</p> <p>60</p>	<ul style="list-style-type: none"> For desiccation of weeds, including London rocket, sowthistle, rescue brome, wild oats, chickweed, ryegrass, bluegrass, cheatgrass, dogfennel, tansymustard, henbit, downy brome, and other winter annuals, and suppression of perennial weeds. Apply during late fall or winter months after the last fall cutting and before first spring cutting. Do not apply if <u>regrowth</u> after grazing or cutting is more than 2 inches. Do not harvest within 60 days of application. CAUTION: Applications to clover or other legumes that is not dormant, or has broken dormancy, may result in stand and/or yield reductions. Replanting may be necessary. Green clover or other legumes foliage present at the time of application will be burned. Clover or other legumes foliage present at the time of application will be discolored and temporarily stunted. Total hay yield of first cutting may be reduced in clover or other legumes fields with severe weed infestation. This reduction will usually be directly proportionate to the loss of weed weight. Do not apply more than once per season. <p>California</p> <ul style="list-style-type: none"> For desiccation of weeds including bluegrass, ryegrass, shepherd's-purse, chickweed, tansymustard, foxtail, sowthistle and groundsel. Use high rate if ryegrass, shepherd's-purse, sowthistle or groundsel is present. <p>¹Other legumes include velvetbean, lespedeza, lupine, sainfoin, trefoil, vetch, crown vetch, and milk vetch.</p>
<p><u>CORN</u> <u>FIELD CORN</u> <u>POPCORN</u> <u>SWEETCORN</u> <u>SEED CORN</u> (Used alone)</p>	<p>Preplant or Preemergence Broadcast or Banded Over Row</p>	<p>Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts</p>	<p>Ground: 10 gal Air: 5 gal</p>		<ul style="list-style-type: none"> Includes field, fresh, sweet, forage, fodder and popcorn. Seedbeds should be formed as far ahead of planting and treatment as possible to permit maximum weed and grass emergence. Seeding should be done with a minimum amount of soil disturbance. Weeds and grasses emerging after application will not be controlled. Crop plants emerged at time of application will be killed.

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments																										
<p><u>CORN</u> Tank Mixes for No-till/Reduced Till</p>	<p>Preplant or Preemergence Broadcast or Banded Over Row</p>	<p>Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts</p>	<p>Ground: 10 gal Air: 5 gal**</p>	<p>-</p>	<ul style="list-style-type: none"> • Apply as a broadcast spray before, during or after planting, but before crop emergence. • For improved burndown or residual control, CYCLONE may be tank mixed with the following herbicides: <table style="margin-left: 20px; border: none;"> <tr> <td>2,4-D Ester (Low Volatile)</td> <td>Harness</td> </tr> <tr> <td>AAtrex/Atrazine</td> <td>Harness Xtra</td> </tr> <tr> <td>Banvel</td> <td>Harmony® Extra herbicide</td> </tr> <tr> <td>Bicep</td> <td>Lasso® herbicide</td> </tr> <tr> <td>Bicep Lite® herbicide</td> <td>Linex</td> </tr> <tr> <td>Bladex</td> <td>Lorox</td> </tr> <tr> <td>Dual® herbicide</td> <td>Modown® herbicide</td> </tr> <tr> <td>Extrazine</td> <td>Princep</td> </tr> <tr> <td>Frontier</td> <td>Prowl® herbicide</td> </tr> <tr> <td>Griffex</td> <td>Simazine</td> </tr> <tr> <td>Guardsman</td> <td>Surpass EC</td> </tr> <tr> <td></td> <td>Surpass 100</td> </tr> <tr> <td></td> <td>TopNotch</td> </tr> </table> • CYCLONE may also be tank mixed with AMBUSH® Insecticide. • Refer to respective product label(s) for rates of application, directions for use, limitations, cautions, and for a list of weeds or insects controlled. <p>**Refer to respective product labels to determine if these products can be applied by air.</p>	2,4-D Ester (Low Volatile)	Harness	AAtrex/Atrazine	Harness Xtra	Banvel	Harmony® Extra herbicide	Bicep	Lasso® herbicide	Bicep Lite® herbicide	Linex	Bladex	Lorox	Dual® herbicide	Modown® herbicide	Extrazine	Princep	Frontier	Prowl® herbicide	Griffex	Simazine	Guardsman	Surpass EC		Surpass 100		TopNotch
2,4-D Ester (Low Volatile)	Harness																														
AAtrex/Atrazine	Harness Xtra																														
Banvel	Harmony® Extra herbicide																														
Bicep	Lasso® herbicide																														
Bicep Lite® herbicide	Linex																														
Bladex	Lorox																														
Dual® herbicide	Modown® herbicide																														
Extrazine	Princep																														
Frontier	Prowl® herbicide																														
Griffex	Simazine																														
Guardsman	Surpass EC																														
	Surpass 100																														
	TopNotch																														

10/29/96

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p><u>FIELD CORN</u> <u>POPCORN</u> <u>SWEETCORN*</u> <u>SEED CORN</u> *Currently not registered for use in California on sweet corn.</p>	<p>Postemergence Directed Spray (Including Hooded or Shielded)</p>	<p>1 - 2 pints</p>	<p>Ground: 10 gal</p>	<p>-</p>	<ul style="list-style-type: none"> • Apply when weeds are actively growing. • Use higher rate on larger or hard to control weeds. Weeds 6 inches or taller may not be controlled. • Severe damage and/or complete kill can occur if spray contacts corn plants. <p><u>HOODED OR SHIELDED SPRAYERS</u></p> <ul style="list-style-type: none"> • To avoid excessive crop phytotoxicity, use a hooded or shielded sprayer with skids or wheels on the spray boom to maintain spray height. • Apply by directing spray between the rows and using hooded or shielded sprayers to prevent spray contact with crop plants. <p><u>DIRECTED SPRAY WITHOUT HOODED OR SHIELDED SPRAYERS</u></p> <ul style="list-style-type: none"> • Apply when corn is at least 10 inches tall with nozzles arranged to spray no higher than the lower 3 inches of corn stalks. • Corn plants shorter than 10 inches may be injured and not recover (corn height measured from soil surface to top of whorl). • For corn greater than 20 inches tall, arrange the nozzles to spray no higher than the lower 1/3 of the corn stalks. • Corn foliage sprayed will be injured, but the crop will recover and develop normally.
<p><u>FIELD CORN ONLY</u> (grain, fodder, forage)</p>	<p>Postemergence directed spray USDA witchweed eradication program</p>	<p>2 pints</p>	<p>Ground: 10 gal</p>	<p>-</p>	<ul style="list-style-type: none"> • Initiate sprays in late June to early July and repeat in early August if regrowth occurs. • Follow application instructions in postemergence directed spray section above.
<p><u>FIELD CORN ONLY</u> (grain, fodder, forage) 2,4-D Amine Tankmix</p>	<p>Postemergence directed spray USDA witchweed eradication program</p>	<p>8 fl oz + 0.5 lbs 2,4-D Amine AE</p>	<p>Ground: 10 gal</p>	<p>-</p>	<ul style="list-style-type: none"> • Apply as a directed spray onto grassy weeds and witchweed before witchweed blooms. Reapply if regrowth occurs. • Follow application instructions in postemergence directed spray section above.

30/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>COTTON</u> (Used alone)	Preplant or Preemergence Broadcast	2 - 4 pints	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Apply prior to, during or after planting, but before crop emergence. • For fallow bed treatment, beds should be preformed to permit maximum weed and grass emergence prior to treatment. • Seeding should be done with a minimum of soil disturbance.
<u>COTTON</u> Goal® herbicide Tankmix	Preplant or Fallow Bed	1 - 4 pints	Ground or Air: 10 gal	-	<ul style="list-style-type: none"> • Refer to Goal label for specific use directions and restrictions, and weeds controlled.
<u>COTTON</u> Bladex Tankmix	Preplant Broadcast	2 - 4 pints	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Refer to Bladex label for specific use directions and restrictions and weeds controlled.
<u>COTTON</u> Other Tankmixes	Preplant or Preemergence			-	<ul style="list-style-type: none"> • Apply as a broadcast spray before, during or after planting, but before crop emergence. • For improved burndown or residual control, CYCLONE may be tank mixed with the following herbicides: <ul style="list-style-type: none"> - Cotoran® herbicide - Meturon® herbicide - Cotton-Pro® herbicide - Caparol® herbicide - Diuron - Harmony Extra - MSMA - Zorlak® herbicide - Dual - Prowl • When tank mixing with Cotoran DF or Meturon DF, follow mixing instructions in the "Order of Tank Mixing" section carefully and maintain constant agitation. • When tank mixing with any of the herbicides listed above, refer to that product's label for specific directions and restrictions and for a list of weeds controlled.

31/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>COTTON</u>	Harvest Aid			3	<p><u>Harvest Aid Use Precautions</u> (Applies to all sections)</p> <ul style="list-style-type: none"> • Do not pasture livestock in treated fields or feed treated foliage. • Do not apply to cotton within 3 days before harvest. • Repeat application if necessary. Do not exceed a total of 2 pts. per acre as a harvest aid. • May be tankmixed with other cotton harvest aid materials known to be effective by the local expert. Unless otherwise instructed in this label, refer to tankmix product label for rates, directions, limitations and cautions. • CYCLONE can be applied in a tankmix with methyl-parathion and/or KARATE® insecticide. • Nodes above cracked bolls (NACB) timing is for guidance and is not intended to restrict the local expert in their use of the product.
<u>SOUTHERN COTTON</u> Harvest aid for boll opening and defoliation (Tank mix with phosphate and chlorate defoliant)	Broadcast	8 fl. oz. plus 1 pt. phosphate or 1 gal chlorate	Ground: 10 gal Air: 5 gal	7	<ul style="list-style-type: none"> • Development of immature bolls will be inhibited. • Apply when 80% or more of the bolls are open and the remaining bolls to be harvested are mature.
<u>SOUTHERN COTTON</u> Additional tankmixes for boll opening and defoliation	Broadcast	4 - 6 oz.	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • To aid in defoliation and opening of mature bolls, CYCLONE may be tank mixed with the following products: ACCELERATE® defoliant FOLEX® defoliant DEF® defoliant HARVADE® harvest growth regulant DROPP® defoliant PREP™ PGR ETHEPHON plant growth regulant • Apply when 60% or more of the bolls are open and the remaining bolls to be harvested are mature. • Development of immature bolls will be inhibited. • Refer to tank mix product label for rate, directions, limitations and cautions.

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<p><u>SOUTHERN COTTON</u></p> <p>Post Defoliation-To aid in opening of mature bolls and to desiccate green weeds</p>	Broadcast	1 - 2 pints	Ground: 10 gal Air: 5 gal	3	<ul style="list-style-type: none"> • Use higher rate if weed infestation is heavy or dense. • Apply when 75% or more of the bolls are open and remaining bolls to be harvested are mature. • Development of immature bolls will be inhibited. • After a defoliation or conditioning application has been made, delay desiccation application of CYCLONE approximately 3-7 days to minimize leaf sticking.
<p><u>WESTERN COTTON</u></p> <p>Harvest aid for boll opening and early defoliation</p>	Broadcast	1/3 - 1/2 pint plus phosphate or sodium chlorate; and/or other compatible harvest aid products.	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Use higher rate of CYCLONE on rank cotton. • Do not use more than 1/2 pt. of CYCLONE for early defoliation as excessive desiccation may occur. • Early defoliation timing is when 60% or more of the bolls are open and the remaining bolls to be harvested are mature (approximately 4 NACB). • Development of immature bolls will be inhibited. • Do not use more than 4.0 lbs. of actual sodium chlorate defoliant per acre at this early defoliation timing.
<p><u>WESTERN COTTON</u></p> <p>Harvest aid for boll opening and mid-to-late defoliation</p>	Broadcast	1/2 - 1 pt. alone or tank mix with sodium chlorate or phosphate defoliant and/or other compatible harvest aid products.		3 (Alone)	<ul style="list-style-type: none"> • In desert cotton areas or on rank vigorous cotton, use the 21 fl oz rate of CYCLONE. • Mid-to-late defoliation timing is when 75% or more of the bolls are open and remaining bolls to be harvested are mature (approximately 3 or fewer NACB). • Development of immature bolls will be inhibited.

33/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments								
<p><u>COTTON</u> Stripper or Spindle Harvested</p> <p>Harvest aid for defoliation and boll opening</p>	Broadcast	3 - 11 fl oz	Ground: 10 gal Air: 5 gal	3	<ul style="list-style-type: none"> • IT IS ADVISABLE, BECAUSE OF EXTREMES IN ENVIRONMENTAL AND PLANT CONDITIONS, TO APPLY THE RANGE OF RATES ON A SMALL BLOCK OF COTTON TO DETERMINE THE RATE THAT BEST FITS YOUR NEEDS. • Apply when 75% of the bolls are open and the remaining bolls to be harvested are mature. • DEVELOPMENT OF IMMATURE BOLLS WILL BE INHIBITED. SLICE BOLLS AND INSPECT THE SEED FOR MATURITY. • CYCLONE may be applied alone or tank mixed with the following cotton harvest aids: <table style="margin-left: 20px; border: none;"> <tr> <td>ACCELERATE defoliant</td> <td>FOLEX defoliant</td> </tr> <tr> <td>DEF defoliant</td> <td>HARVADE harvest growth regulant</td> </tr> <tr> <td>DROPP defoliant</td> <td>PREP PGR</td> </tr> <tr> <td>ETHEPHON plant growth regulant</td> <td></td> </tr> </table> • May be applied as a split application. Do not exceed a total of 2 pts/Acre. • To avoid leaf sticking, apply CYCLONE as a desiccant approximately 3-7 days after defoliation or a conditioning application and 7-14 days before harvest. • Cooler temperatures may cause a longer waiting period between application of CYCLONE as a desiccant and defoliation/condition. • Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation. 	ACCELERATE defoliant	FOLEX defoliant	DEF defoliant	HARVADE harvest growth regulant	DROPP defoliant	PREP PGR	ETHEPHON plant growth regulant	
ACCELERATE defoliant	FOLEX defoliant												
DEF defoliant	HARVADE harvest growth regulant												
DROPP defoliant	PREP PGR												
ETHEPHON plant growth regulant													

34/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>COTTON</u> Late season desiccation	Broadcast	1 - 2 pints	Ground: 10 gal Air: 5 gal	3	<ul style="list-style-type: none"> • IT IS ADVISABLE, BECAUSE OF EXTREMES IN ENVIRONMENTAL AND PLANT CONDITIONS, TO APPLY THE RANGE OF RATES ON A SMALL BLOCK TO DETERMINE THE RATE THAT BEST FITS YOUR NEEDS. • May be applied as a split application. Do not exceed a total of 2 pts/Acre. • Apply when 85% of the bolls are open and the remaining bolls to be harvested are mature (approximately 0 NACB). • Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity. • Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation. • If a defoliation or conditioning application has been made, delay desiccation application of CYCLONE approximately 3-7 days to minimize leaf sticking. • May be tankmixed with other harvest aid materials known to the local expert to be effective.
<u>COTTON</u> Desiccation of Regrowth	Broadcast	1 - 2 pints	Ground: 10 gal Air: 5 gal	3	<ul style="list-style-type: none"> • Use to desiccate regrowth occurring after defoliation or desiccation. • Regrowth is difficult to control, therefore, thorough coverage with the full recommended rate is necessary. • Control is dependent on growing conditions and desiccation of small new regrowth may not always be complete. • Use higher rate if regrowth is excessive.
<u>EASTER LILIES</u> (Field grown)	Preemergence	2 - 4 pints (use a minimum of 3 pts. in California)	Ground: 10 gal	-	<ul style="list-style-type: none"> • Do not apply more than twice per season.

35/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>FALLOW LAND*</u> Prior to planting of any crops approved on this labelling for preplant/ preemergence use	Preplant Broadcast to Fallow Land	2 - 4 pints	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Fallow land may be between operations such as disking, ripping, plowing, leveling, irrigating or listing for ground preparation purposes. • Use for the control of weeds such as bluegrass, chickweed, henbit, downy brome, ryegrass, cheatgrass, dog fennel, tansy mustard, London rocket, sowthistle, rescue brome, wild oats, volunteer cereals and other winter annuals and for suppression of perennial weeds or sedges. • Use the higher rate for weeds approaching the maximum size of 6". • Do not make more than two applications during the fallow period. • Allow maximum weed emergence prior to application to maximize the benefit of this use. • Adhere to the preharvest intervals and other crop specific restrictions for planted crops elsewhere on this label.
<u>GRASSES</u> (For Seed) (For Use in Seedbed Preparation)	Preplant, At Planting, or Preemergence	2 - 4 pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • Prepare the seedbeds and allow weeds to germinate. • Apply CYCLONE when weeds are at the 3 to 5 leaf stage. • Repeat applications as necessary prior to grass emergence. • Do not graze treated areas or use the seed or straw from treated areas for animal feed or bedding.
<u>GUAR</u> (Preharvest desiccation)	Preharvest	2 pints	Ground: 10 gal	4	<ul style="list-style-type: none"> • Apply after the pods are fully mature. • Do not graze treated areas or use the treated forage for animal feed.
<u>GUAVA</u>	Directed Spray	4 pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • Do not allow spray to contact green stems, fruit or foliage. • Do not graze treated areas. • Do not feed cover crops grown in treated areas to livestock. • For mature woody weeds, late-germinating weeds and grasses, and perennials, retreatment or spot spraying may be necessary.

36/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>HOPS</u> (ID, OR, & WA only)	Directed Spray and/or Suckering and Stripping	2 pints	Ground: 10 gal	14	<ul style="list-style-type: none"> • Retreatment or spot treatment may be necessary. • Do not apply more than 3 times per season. • Do not allow spray to contact green stems, foliage, flowers, or cones as injury may result. • Do not allow animals to graze in treated hopyards. • Hop vine refuse and silage may be fed to livestock. • For suckering and stripping, spray only the basal 2 ft. of the vines. Repeat as necessary. • Experience with varieties other than CASCADE, YAKIMA CLUSTER, and BULLION is limited. If using CYCLONE on other varieties than these, test the use pattern on a small number of vines of each variety to determine sensitivity to injury. Do not use on unlisted varieties if unacceptable crop injury occurs. • Chemical Pruning: To burn back existing vines and obtain even emergence of subsequent vines, spray when vines are less than 3 ft tall. • APPLICATION TO HOP VINES LESS THAN 6 FT TALL MAY CAUSE UNACCEPTABLE INJURY.
<u>MINT</u> (Peppermint, Spearmint)	Dormant Season	1½ - 3 pints	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • For suppression of weeds such as Italian ryegrass, prickly lettuce, groundsel, chickweed, downy brome and bluegrass. • Apply when crop is dormant before spring growth begins and when weeds are less than 6 inches tall. • Do not apply more than 2.4 pts per acre per dormant season. • May be tank mixed with Sinbar® herbicide (terbacil) weed killer for improved contact activity and residual control of Italian ryegrass, prickly lettuce and groundsel. Apply this tank mixture no more than once per season. Refer to the Sinbar label for rates, directions, and cautions and for a list of weeds controlled.
<u>ONIONS (seeded)</u> <u>AND GARLIC</u>	Preplant or Preemergence	2 - 4 pints (use a minimum of 3 pts. in California)	Ground: 10 gal	60 200 (CA only)	<ul style="list-style-type: none"> • Use the higher rate for heavy weed infestations or wild oat control. Apply only one application per season at the 5 pt per acre dosage. • Allow maximum weed and grass emergence prior to treatment but apply prior to crop emergence. • Apply a maximum of 5 pt per acre per season.

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>PASSION FRUIT</u>	Directed Spray	4 pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • Use a shield or wrap vine if bark is still green at application time. • If application is to be made during harvest season, pick all fruit off the ground prior to application. • Do not allow animals to graze on treated areas. • Retreatment or spot treatment may be necessary.
<u>PEANUTS</u>	Broadcast At Ground Crack Postemergence	8 - 16 fl oz	Ground: 10 gal	-	<ul style="list-style-type: none"> • To control or suppress small (1-6") emerged annual grass and broadleaf weeds in peanuts at ground crack. A second application may be made up to 28 days after ground crack. • For at ground crack use, CYCLONE can be tank mixed with Pursuit® herbicide or Dual for residual weed control. Consult the Pursuit or Dual label for a list of weeds controlled, rates of application, and precautions. • Make no more than two applications per season and do not apply a total of more than 16 fl oz of product per acre per season. • Crop foliage sprayed will be injured in the form of bronzing and crinkling but the crop will recover and develop normally. • Do not apply by air.

38/
61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
PEANUTS Basagran® herbicide tankmix	Broadcast At Ground Crack Postemergence	8 - 16 fl oz	Ground: 10 gal	-	<ul style="list-style-type: none"> • For improved control of weeds such as cocklebur, bristly starbur, smartweed and prickly sida, tank mix CYCLONE with Basagran at 1 pint/acre. • This tankmix can be applied at the ground crack stage of peanuts. A second application may be made up to 28 days after ground crack. • Make no more than two applications per season and do not apply a total of more than 16 fl oz of product per acre per season. • Crop foliage sprayed will be injured in the form of bronzing and crinkling but the crop will recover and develop normally. • Refer to the Basagran label for specific use directions, limitations, cautions and for a list of weeds controlled. • Do not apply this tankmix if peanuts show injury (leaf phytotoxicity and/or plant stunting) produced by any other herbicide treatment as injury may be enhanced and/or prolonged. • Do not apply this tankmix during prolonged periods of drought or unseasonably cold weather as unsatisfactory weed control may result. • Do not apply by air.
PEANUTS Butyrac® herbicide or Butoxone™ herbicide 200 tankmix	Broadcast Postemergence	8 - 16 fl oz	Ground: 10 gal	-	<ul style="list-style-type: none"> • For improved control of weeds such as cocklebur, sicklepod and morningglory tank mix CYCLONE with 8-16 oz (0.125-0.25 lbs) per acre of Butyrac or Butoxone 200. • Make no more than two applications per season and do not apply a total of more than 16 fl oz of product per season. • Crop foliage sprayed will be injured in the form of bronzing and crinkling but the crop will recover and develop normally. • Refer to the complete Butyrac or Butoxone 200 label for specific use directions, limitations, cautions and for a list of weeds controlled. • Do not apply by air.
PIGEON PEAS (Puerto Rico only)	Directed Spray	2 pints	Ground: 10 gal	60	<ul style="list-style-type: none"> • Avoid contact with pigeon pea foliage. • Do not make more than 1 application per season. • Do not graze treated areas or feed treated forage to livestock. • Cannery waste can be fed to livestock.

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>PINEAPPLE</u>	Directed Spray	2 - 4 pints	Ground: 10 gal	20	<ul style="list-style-type: none"> • Retreatment may be necessary on more mature weeds. • Do not exceed 3 applications per season.
<u>POTATO</u>	Preplant or Preemergence Broadcast	1 - 2 pints	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Apply up to ground cracking, before potatoes have emerged.
<u>POTATO</u> (California, Washington, Oregon, Idaho only) (Used alone)	Preplant Broadcast	8 - 16 fl oz	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • For control of volunteer barley in preformed seedbeds.
<u>POTATO</u> Preharvest vine killing and weed desiccation (Not for use in the states of Alabama, Arkansas, Georgia, Louisiana, Mississippi, New Mexico, North Carolina, Oklahoma, South Carolina, Tennessee and Texas)	Broadcast	1 - 2 pints	Ground: 20 gal	3	<ul style="list-style-type: none"> • Begin application when leaves begin to turn yellow. • Immature or drought-stressed potato foliage are tolerant to CYCLONE. Desiccation will not be complete under these conditions. • Use 2½ pts/acre rate where quick vine kill is desired. • Use two applications of 1½ pts/acre when vine growth is dense. Split applications must be applied a minimum of five days apart. • Do not pasture livestock in treated potato fields. • Do not use on muck or peat soils to avoid injury to subsequent crops. • DO NOT use to desiccate potato vines when potatoes are to be stored or used for seed as this use may result in tuber decomposition or failure of seed pieces to germinate and grow normally. • Do not exceed 5 pts/acre per season.
<u>RESIN SOAKING</u> (Pine)	See Page _____				

10/1

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>SAFFLOWER</u>	Preplant or Preemergence Broadcast or Banded Over Row	2 - 4 pints (use a minimum of 3 pts. in California)	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Apply before, during, or after planting but before crop emergence.
<u>SMALL FRUITS</u> Blackberries Boysenberries Raspberries Blueberries Strawberries	Postemergence Directed Spray Postemergence Directed Spray	2 - 4 pints 2 pints	Ground: 50 gal Ground: 20 gal	- 21	<ul style="list-style-type: none"> • Apply before emergence of new canes or shoots as injury to those canes or shoots can occur. • Apply as a coarse spray to avoid crop injury from fine spray mist. • Apply by directing spray between the rows and using shields to prevent spray contact with crop plants. • Do not allow spray to contact strawberry plants as injury or excessive residues may result. • Do not apply more than 3 times per season. • Do not graze livestock in treated areas.
<u>SMALL GRAINS</u> (Barley, wheat)	Preplant or Preemergence	Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts	Ground: 10 gal Air: 5 gal	-	

9/1/96

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
SMALL GRAINS (Wheat Only) Hoelon 3EC Tankmix	Preplant or Preemergence	Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4pts	Ground: 5 gal Air: 5 gal	-	<ul style="list-style-type: none"> • A tankmix with Hoelon 3EC will improve grass control. • Apply when weeds are actively growing and 1-8 inches in height. Weeds 6 inches or taller may not be controlled. • Do not apply this tankmix to Barley as crop injury may result.
SORGHUM (Grain)	Preplant or Preemergence Broadcast or Band	Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Seedbeds should be formed as far ahead of planting as possible to allow maximum weed and grass emergence. • Seeding should be done with a minimum amount of soil disturbance.
SORGHUM (Grain) Atrazine & 2,4-D ester [Low Volatile] tankmix	Preplant or Preemergence	Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • CYCLONE may be tank mixed with Atrazine for improved preemergence or residual weed control. The addition of 2,4-D ester (Low Volatile) may aid in the suppression of perennial and annual broadleaf weeds emerged at the time of application. Refer to the specific tankmix herbicide label(s) for rates, directions, limitations, and cautions and a list of weeds controlled.
SORGHUM (Grain) Harmony Extra tankmix	Preplant or Preemergence	1 - 4 pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • CYCLONE may be tank mixed with Harmony Extra for improved preemergence or residual weed control. • Refer to the Harmony Extra label for rates, directions, limitations, and cautions and for a list of weeds controlled.

42/174

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
SORGHUM (Grain)	Postemergence Directed Spray (Includes Hooded or Shielded)	1 - 2 pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • Apply when weeds are actively growing. • Use higher rate on larger or hard to control weeds. Weeds 6 inches or taller may not be controlled. • Severe damage and/or complete kill can occur if spray contacts sorghum plants. • Do not exceed 2 postemergence-directed applications per season or exceed a total of 8 pints CYCLONE per season. <p><u>HOODED OR SHIELDED SPRAYERS</u></p> <ul style="list-style-type: none"> • To avoid excessive crop phytotoxicity, use a hooded or shielded sprayer with skids or wheels on the spray boom to maintain spray height. • Apply by directing spray between the rows and using hooded or shielded sprayers to prevent spray contact with crop plants. <p><u>DIRECTED SPRAY WITHOUT HOODED OR SHIELDED SPRAYERS</u></p> <ul style="list-style-type: none"> • Apply when sorghum is at least 12 inches tall when naturally standing. • Do not exceed 30 psi nozzle pressure or spray under conditions which may cause excessive drift. • Use precision directed-spray application equipment adjusted so that no more than the lower 3 inches of the sorghum stalk is contacted by the application spray. • Some crop injury will occur. The degree of injury is related to the precision of application and spraying conditions. • Do not harvest forage, fodder, silage, or hay before the soft dough stage of the grain.

43/
/61

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments																								
<u>SOYBEANS</u>	Preplantor Preemergence	Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Do not exceed a total of 6 pints of CYCLONE per season. • Apply as a broadcast spray before, during or after planting, but before crop emergence. • For improved burndown or residual control, CYCLONE may be tank mixed with the following herbicides: <table style="margin-left: 20px; border: none;"> <tr> <td>2,4-DB</td> <td>Linex</td> <td></td> </tr> <tr> <td>Canopy</td> <td>Lorox</td> <td>Sencor</td> </tr> <tr> <td>Dual</td> <td>Lorox Plus</td> <td>Surflan® herbicide</td> </tr> <tr> <td>Gemini</td> <td>Modown</td> <td>Turbo® herbicide</td> </tr> <tr> <td>Goal</td> <td>Preview</td> <td></td> </tr> <tr> <td>Harmony Extra</td> <td>Prowl</td> <td></td> </tr> <tr> <td>Lasso</td> <td>Pursuit herbicide</td> <td></td> </tr> <tr> <td>Lexone</td> <td>Scepter® herbicide</td> <td></td> </tr> </table> • The rate of CYCLONE to be used in these tank mixtures is dependent on weed height and growing conditions. Use the highest recommended rate of CYCLONE under dry conditions or where the weed canopy is dense. Refer to the specific tank-mix herbicide label(s) for rates, directions, limitations, and cautions and for a list of weeds controlled. • The lower rate may be used when weeds are less than 4" tall and a selective post-emergence spray or cultivation will be made within 3 weeks after planting. • Seeding should be done with a minimum amount of soil disturbance. • Do not graze or harvest for forage or hay before the R3 stage of soybean development (early pod) 	2,4-DB	Linex		Canopy	Lorox	Sencor	Dual	Lorox Plus	Surflan® herbicide	Gemini	Modown	Turbo® herbicide	Goal	Preview		Harmony Extra	Prowl		Lasso	Pursuit herbicide		Lexone	Scepter® herbicide	
2,4-DB	Linex																												
Canopy	Lorox	Sencor																											
Dual	Lorox Plus	Surflan® herbicide																											
Gemini	Modown	Turbo® herbicide																											
Goal	Preview																												
Harmony Extra	Prowl																												
Lasso	Pursuit herbicide																												
Lexone	Scepter® herbicide																												

19/4/97

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
SOYBEANS 2,4-D ester (Low Volatile)* tankmix	Preplant or Preemergence	Weeds 1-3": 1 - 2½ pts Weeds 3-6": 2½ - 3 pts Weeds 6": 3 - 4 pts	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Apply 2,4-D ester (Low Volatile) at 0.35-0.475 lbs ai/A at least 7 days prior to planting. • Apply 2,4-D ester (Low Volatile) at 0.475-0.95 lbs ai/A at least 30 days prior to planting. • Do not apply 2,4-D ester (Low Volatile) prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield. • Do not use the amine formulation as CYCLONE activity may be reduced. • May be tank mixed with residual herbicides listed above. • Refer to the 2,4-D ester (Low Volatile) label for a list of rates, directions, limitations and cautions and for a list of weeds controlled.

17/54

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>SOYBEANS</u>	Postemergence, Directed Spray (Includes Hooded or Shielded)	1/3 to 2/3 pts	Ground: 10 gal	-	<ul style="list-style-type: none"> • Apply when weeds are actively growing. • For control of seedling johnsongrass, crabgrass, goosegrass, brachiaria, Texas millet and pigweed less than 2 inches tall, use the lower rate of CYCLONE. • For control of 2-4 inch red rice, <i>Brachiaria</i>, banyard grass, crabgrass, goosegrass, seedling johnsongrass, giant foxtail, and fall panicum, use 8 fl oz of CYCLONE. • For control of 2-3 inch sicklepod, purslane, pigweed, cutleaf ground cherry, and common ragweed, use 8 fl oz of CYCLONE. • For control of 2-4 inch grasses in mixture with common cocklebur, morningglory, and red rice, apply CYCLONE at 8 fl oz per acre plus 0.2 lb active ingredient per acre of a 2,4-DB formulation. • Refer to the 2,4-DB label for directions, limitations, and cautions. • Do not graze or harvest for forage or hay. • If needed make a second and final application 7-14 days later. <p><u>HOODED OR SHIELDED SPRAYERS</u></p> <ul style="list-style-type: none"> • Apply by directing spray between the rows and using hooded or shielded sprayers to prevent spray contact with crop plants. • Use higher rate on larger (> 6 inches) or hard to control weeds. Weeds 6 inches or taller may not be controlled. • Severe damage and/or complete kill can occur if spray intentionally or accidentally (including drift of fine droplets) contacts the plants. <p><u>DIRECTED SPRAY WITHOUT HOODED OR SHIELDED SPRAYERS</u></p> <ul style="list-style-type: none"> • Do not treat if soybeans are less than 8 inches tall. • Use precision directed spray application equipment adjusted so that no more than the lower 3 inches of the soybean plant is contacted by the application spray. • Do not exceed 30 psi nozzle pressure or spray under conditions which may cause excessive drift. • Some crop injury will occur. The degree of injury is related to the precision of application and spraying conditions.

19/96

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>SOYBEANS</u>	Harvest Aid	½ - 1 pint	Ground: 20 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Indeterminant varieties: Apply when at least 65% of the seed podshave reached a mature brown color or when seed moisture is 30% or less. • Determinant varieties: Apply when plants are mature, i.e. beans are fully developed, 1/2 of leaves have dropped, and remaining leaves are yellowing. • Immature soybeans will be injured. • Mature cocklebur, especially drought-stressed plants, are tolerant to CYCLONE and desiccation will not be complete. Always use the higher rate for cocklebur. • Do not apply within 15 days of harvest. • Do not graze or harvest for forage or hay.
<u>SUGAR BEETS</u>	Preplant or Preemergence	2 - 4 pints	Ground: 10 gal Air: 5 Gal	-	<ul style="list-style-type: none"> • Use the higher rate for heavier weed infestations. • Seeding or transplanting should be done with a minimum amount of soil disturbance. • Crop plants emerged at time of application will be killed. • Can be used in fallow bed/stale seedbed for weed control. • Seedbeds or plantbeds should be formed as far ahead of treatment as possible to permit maximum weed emergence.
<u>SUGARCANE</u>	Postemergence Directed Spray (Includes Hooded or Shielded)				<p><u>General Comments</u></p> <ul style="list-style-type: none"> • Apply as a hooded, shielded or directed spray to avoid contact with cane foliage to prevent leaf burn and yield reduction. • Make a second and final application, if necessary, when new weed growth is 2-8 inches high. • Do not graze treated areas or feed treated forage to livestock.
-- Florida --		2 pints	Ground: 50 gal	-	<ul style="list-style-type: none"> • For optimum results, apply in early spring (March-April) when weeds are small. • Do not apply after June 1 as cane growth may be stunted and yields reduced.
-- Hawaii --		2 pints	Ground: 20 gal	-	<ul style="list-style-type: none"> • Do not apply after cane rows have closed in.
-- Louisiana --		1 - 2 pints	Ground: 20 gal	30	<ul style="list-style-type: none"> • For tiller control, apply when tillers are less than 18 inches high. • Use the higher rate for heavier weed infestations or tiller growth.

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
-- Florida & Texas --	Harvest Aid	8 - 16 fl oz	Air: 5 gal	-	<ul style="list-style-type: none"> • Use higher rate under cool, cloudy weather conditions. • Apply 3-14 days before burning and harvest.
<u>SUNFLOWER</u>	Preplant or Preemergence Broadcast or Banded Over Row	2 - 4 pints (use a minimum of 3 pts. in California)	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Apply before, during, or after planting but before crop emergence.
<u>SUNFLOWER</u>	Preharvest Desiccation Broadcast	1 - 2 pts	Ground: 10 gal Air: 5 gal	7	<ul style="list-style-type: none"> • Apply when sunflower seeds reach physiological maturity (when seed moisture is 35% or lower). For many varieties, this corresponds to the time when the back of the heads are yellow and the bracts are turning brown. • Do not graze treated areas or feed treated forage to livestock. • Use the higher rate when crop stands or weed infestations are heavy.
<u>TARO, DRYLAND</u> (Hawaii Only)	Postemergence Directed Spray	1½ - 2½ pts	Ground: 10 gal	180	<ul style="list-style-type: none"> • Do not allow spray to contact the taro plants as injury may result. • Make the first application when weed growth is 1 to 4 inches high. • Weeds emerging after the application will not be controlled. • A single re-treatment may be made; however, do not harvest dryland taro within 6 months of the last application.

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
TREES AND VINES Orchards, Vineyards, Windbreak, Shade & Ornamental Trees Acerola Apples Apricots Avocados Bananas Beechnut Brazil Nut Butternut Cashew Cherries Chestnut Chinquapin Citrus Coffee Figs Filberts Grapes Hickory Nut Kiwi Fruit Macadamia Nuts Nectarines Olives Papayas Peaches Pears Pecans Pistachios Plums	Directed Spray	2 - 4 pints (use a minimum of 3 pts. in California)	Ground: 10 gal	Kiwi Fruit 14 Pistachios 7 Plums 28 Olives 13 Figs 13	<ul style="list-style-type: none"> • Do not allow spray to contact green stems (except suckers), fruit or foliage. • Use a shield or wrap plant when spraying around young trees or vines. • Do not graze treated areas. • Do not feed cover crops grown in treated areas to livestock. • Do not apply when figs, nuts or olives to be harvested are on the ground. • For grapes - treat when sucker growth is no more than 8" long. Late season applications to weeds should be made to avoid contact with desirable foliage. • For kiwi fruit - do not treat more than 3 times per year. • For pistachios - do not exceed two applications after shells split. • For mature woody weeds, perennial weeds, late germinating weeds and green suckers, retreatment or spot treatment may be necessary. • For plums - do not harvest within 28 days after application and do not exceed 3 applications per season. • For figs - do not harvest within 13 days after application and do not exceed 5 applications per season. • For olives - do not harvest within 13 days after application and do not exceed 4 applications per season.

10/1/96

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments								
<u>TREES AND VINES</u> Prunes Walnuts Other shade and ornamental trees such as arborvitae, ash, elm, fir, oak, pine, etc.													
<u>TREES AND VINES</u> Tankmixes	Directed Spray	2½ - 5 pts (use a minimum of 3 pts. in California)	Ground: 10 gal	Refer to other TM labels	<ul style="list-style-type: none"> • CYCLONE may be tank mixed with registered residual herbicides listed below for combined emerged and residual weed control. Always refer to other herbicide label(s) for respective precautions, limitations, restriction, dates and directions for use and weeds controlled. • CYCLONE may be tank mixed with the following herbicides: <table data-bbox="1336 839 1853 954"> <tr> <td>Goal</td> <td>Surflan</td> </tr> <tr> <td>Devrinol® herbicide</td> <td>Solicam® herbicide</td> </tr> <tr> <td>Princep</td> <td>Karmex</td> </tr> <tr> <td>Sinbar</td> <td>Krovar® I and II herbicides</td> </tr> </table> 	Goal	Surflan	Devrinol® herbicide	Solicam® herbicide	Princep	Karmex	Sinbar	Krovar® I and II herbicides
Goal	Surflan												
Devrinol® herbicide	Solicam® herbicide												
Princep	Karmex												
Sinbar	Krovar® I and II herbicides												
<u>TYFON</u> (New Hampshire only)	Preplant Preemergence	2½ - 5 pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • Seeding should be done with a minimum of soil disturbance. • Weeds and grasses emerging after treatment will not be controlled. • Crop plants emerged at time of application will be injured. 								

10/03

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>VEGETABLES</u> (Seeded or Transplanted) Beans (Lima, Snap) Broccoli Cabbage Cantaloupe Carrots Cauliflower Chinese Cabbage Collards Cucumber Eggplant Lettuce Musk Melons Peas Peppers Pumpkin Squash Sweet Corn Turnips Tomatoes Watermelons	Preplant Preemergence	2 - 4 pints (use a minimum of 3 pts. in California)	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • Seedbeds or plantbeds should be formed as far ahead of treatment as possible to permit maximum weed emergence. • Applications can be made as a banded or broadcast treatment before, during or after planting but prior to the crop emergence. • Use the higher rate for heavier weed infestations. • Seeding or transplanting should be done with a minimum amount of soil disturbance. • Crop plants emerged at time of application will be killed. • Can be used in fallow bed/stale seedbed for weed control alone or tank mixed with Goal. Consult the Goal label for a list of weeds controlled, rates of application and precautions. • Do not harvest tomatoes within 30 days after application.
<u>VEGETABLES</u> Tomatoes Peppers	Directed Spray	2 pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • For control or suppression of emerged weeds between rows after crop establishment. • Use precision directed spray application equipment adjusted to prevent spray contact with crop plants. Do not exceed 30 psi nozzle pressure or spray under conditions which may cause excessive drift. • Apply when weeds are succulent and weed growth is less than 6 inches. • Do not apply more than 3 applications per season. • Do not allow animals to graze in treated areas. • Do not harvest tomatoes within 30 days after application.

5/1/91

Crop	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>VEGETABLES</u> Lettuce Melon Sugar Beets Tomatoes (California, Washington, Oregon, Idaho only)	Broadcast	8 - 16 fl oz	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> • For control of volunteer barley in preformed seedbeds. • Do not harvest tomatoes within 30 days after application.
<u>VEGETABLES</u> Rhubarb	Dormant	2 - 4 pints (use a minimum of 3 pts. In California)	Ground: 10 gal	-	<ul style="list-style-type: none"> • Apply during dormant season before buds in crown begin to grow. • Do not make more than 2 applications per season.

52/
/61

ALFALFA

New Seedlings - Suppression and control of broadleaf weeds and grasses in new alfalfa seedlings grown for hay (California only).

For Control of:	Rate/Acre	
	For Suppression	For Control
Spikeweed (4 in. tall or less)	½ pt.	1 - 2 pts.
Volunteer Small Grain (8 in. tall or less)	½ to 1 pt.	2 pts
Fiddleneck (6 in. tall or less)	½ to 1 pt.	2 pts.
Shepherd's-purse	1 - 2 pts.	-
Annual Bluegrass	-	1 to 2 pts.
Chickweed	-	1 to 2 pts.
Red Maids (6 in. tall or less)	-	1 to 2 pts.

Do not use the ½ pt rate unless the alfalfa has at least 3 trifoliate leaves; the 1 pt rate unless the alfalfa has 6 trifoliate leaves; or rates over 1 pt unless there are 9 trifoliate leaves.

RESIN SOAKING

Pines (Loblolly, Shortleaf, Longleaf, Slash, Virginia, Pond, Pitch, and Spruce Pines).

Tree Selection - Select trees to be treated from stands on sites not subject to periods of extreme drought stress as the desiccating effect of CYCLONE herbicide to pines is accentuated during such periods, causing a reduction in the amount of oleoresin deposited in the xylem. Select trees to be treated from vigorous, non-stagnated stands, either natural or planted. In stagnated stands or commercial timber stands, plan treating with CYCLONE herbicide not sooner than three years after a commercial thinning.

Application Directions - Apply CYCLONE herbicide diluted in water (~~0.2% to 5% cation, wt./wt. basis, depending on the type of treatment method~~) to a suitable wound in the tree trunk to bring the treatment into contact with the xylem (sapwood). ~~One of the following methods may be used:~~

Bark Streaks or Cuts: This type of wound is made using a standard or rotary bark hack or a chainsaw chipping tool employed in naval stores work to remove a single one-inch wide streak of bark about 1 to 2 feet from ground level. The total length should not exceed 1/3 of the tree circumference. Multiple streaks or cuts can result in serious girdling of the trunk and premature death of the tree. A coarse spray (about 2 to 6 ml.) of CYCLONE herbicide solution (1% to 5% cation, wt./wt. basis) should be applied to runoff to the exposed xylem, using a plastic acid bottle or a low-pressure sprayer. ~~Wear rubber gloves when using the acid bottle.~~ The amount of spray required per cut depends on tree circumference and the length of cut or

streak (1/3 of circumference). For a 9-inch diameter tree, 3 ml. of spray will cover the 1-inch wide streak. Using 3 ml. of a 2% or 4% CYCLONE herbicide solution will result in application of 60 or 120 mg. CYCLONE herbicide per streak.

~~Tree Injection: Pressure injections of 1% to 4% CYCLONE herbicide solution (cation, wt./wt. basis) should be applied beneath the bark and just beneath the cambium layer. Injections should be made 1 to 2 inches apart around 1/3 of the tree circumference or no closer than 3 inches apart all around the tree. From 0.2 ml to 0.4 ml of CYCLONE herbicide solution is applied per injection. Using 2% to 4% CYCLONE herbicide solution, this results in 36 to 144 mg CYCLONE herbicide applied per 9-inch diameter tree, for holes spaced 1 inch apart around 1/3 of the tree or 3 inches apart all around the tree.~~

~~Bore Holes: 3/8 to 5/8 inch diameter holes 2 to 6 inches deep, depending on tree diameter are bored sloping slightly downward and tangentially inward rather than directly towards the center of the tree. Solutions of 0.3% to 5% CYCLONE herbicide can be used. Lower concentrations* (0.3%) are usually applied at higher volumes of 15 to 35 ml per tree, resulting in application of 45 to approximately 100 mg CYCLON herbicide per tree. Using 2% to 4% solutions and applying 5 ml per single hole per tree results in application of 100 or 200 mg CYCLONE herbicide per tree. Bore hole applications in the range of 45 to 100 mg CYCLONE herbicide per tree have resulted in effective oleoresin induction.~~

~~Time of Treatment: Resin soaking can occur from treatments made any time of the year; however, cool season treatments under nondrought conditions usually result in less severe pine beetle infestations and longer tree life.~~

~~Interval Between Treatment and Tree Harvest: The interval between application of CYCLONE herbicide and tree harvest should be a minimum of 6 months and preferably from 12 to 24 months. Intervals of over 6 months may not be possible under conditions of drought or serious pine beetle attacks, which may make early harvest necessary. The CYCLONE herbicide treatment may encourage beetle attack, or may cause premature death of the tree. Desiccation of the xylem tissue, rather than the desired resin soaking, may occur, and is more likely at higher dosage rates.~~

Effect on Stem Growth:

CYCLONE herbicide treatment can result in reduced stem growth during the interval between treatment and tree harvest.

Dilution Table for CYCLONE Herbicide (2.0 lbs. cation per gallon):	
Concentration of Cation Desired (Wt./Wt. Basis)	To 1½ Gallon of CYCLONE Herbicide add the following No. Gals. of Water:
0.2%	118.8
0.5%	46.8
1.0%	22.9
2.0%	10.9
3.0%	6.9
4.0%	4.9
5.0%	3.7

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

~~NOTE: DO NOT USE AROUND HOME GARDENS, SCHOOLS, RECREATIONAL PARKS, GOLF COURSES, OR PLAYGROUNDS.~~

~~DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.~~

~~DO NOT enter or allow others to enter the treated area until sprays have dried. AVOID working in spray mist. If there is risk of exposure, wear goggles and approved face mask capable of filtering spray droplets.~~

~~KEEP all unprotected persons out of operating areas or vicinity where there may be danger of drift.~~

~~Certain states may require more restrictive reentry intervals; consult your State Department of Agriculture for further information.~~

~~Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.~~

~~IMPORTANT: The effect of swallowing CYCLONE is far more severe than that which usually results from skin contact or inhalation of spray mist. This product should not cause injury if used according to label instructions. Follow the label as if your life depends on it.~~

Other Uses	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
<u>CONSERVATION RESERVE, FEDERAL SET-ASIDE, CONSERVATION COMPLIANCE PROGRAMS</u> (For use in compliance with the Federal Conservation Reserve Program or Federal set-aside programs).	Broadcast	2 - 4 pints	Ground: 10 gal Air: 5 gal	-	<ul style="list-style-type: none"> For improved emerged weed control or extended weed control, CYCLONE may be tank mixed with other herbicides registered for this use. Refer to tank mix herbicide labels for specific directions, limitations, cautions and for a list of weeds controlled.
<u>NONCROP USES</u>	Broadcast or Spot Treatment	2 - 4 pints (use a minimum of 3 pts. in California)	Ground: 10 gal	-	<ul style="list-style-type: none"> For use in noncrop areas such as public airports, electric transformer stations, pipeline pumping stations, around commercial buildings, storage yards and other installations, fence lines or similar noncrop areas. Avoid contact with the foliage of ornamentals or desired plants. Repeat as necessary.

56/61

Other Uses	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
PASTURE RESEEDING For suppression of existing sod and undesirable emerged broadleaf weeds and grasses prior to or at time of planting grasses or forage legumes	Broadcast	1 - 2 pints	Ground: 10 gal Air: 5 gal	See specific geographic recommendation	<u>West of Cascade and Sierra Nevada Mountains:</u> <ul style="list-style-type: none"> • Apply in October through December after first fall rains and after weeds have emerged and sod has started new growth. • For best seeding results, apply on moderately to heavily grazed areas. • Do not use in areas with heavy sod and weed growth. <u>East of Rocky Mountains</u> <ul style="list-style-type: none"> • Use the 2 pint rate on vigorous or coarse sod species such as bromegrass. • Apply prior to, or at time of seeding grasses or forage legumes. • Apply only to grazed or mowed pastures not more than 2 inches in height at time of treatment.
		1 - 2 pints	Ground: 10 gal Air: 5 gal		<u>Bermudagrass or Bahiagrass Sods:</u> <ul style="list-style-type: none"> • Apply in late summer or early fall to sod not exceeding 2 inches in height. • For control of emerged Little Barley, apply in February or March before the midboot stage of Little Barley.
		1 - 2 pints	Ground: 10 gal Air: 5 gal		<u>Bermudagrass and Coastal Bermudagrass Pastures</u> <ul style="list-style-type: none"> • Apply when bermudagrass is dormant. • For control of little barley, apply before the mid-boot stage. • Do not mow for hay until 40 days after treatment.

57/61

Other Uses	Use Pattern	CYCLONE Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
For Control of Endophyte-Fungus-Infected Fescue Forage Legume/Grass Mixture and Other Grass Pastures	Broadcast (Split Application)	1½ - 2½ pints followed by 1½ - 2½ pints	Ground: 10 gal	-	<ul style="list-style-type: none"> • Use split applications of 10 to 21 days apart if necessary. • Do not exceed 5 pints per acre total in preparation for reseeding. • For spring plantings, the initial application of 1½ - 2½ pts may be made the previous fall. • Apply when fescue is actively growing and no more than 4 inches high. • To reduce the infestation of endophyte-infested grass, do not allow fescue to go to seed starting with the preceding year's crop.

5/6/95

Conversion Table CYCLONE Herbicide to be Applied			
Pints/Ac	Fl Oz/Ac	Lb a/Ac	Ac/Gal
-	7	.08	18.3
-	11	.13	11.6
-	21	.25	6.1
1½	24	.28	5.3
2	32	.38	4.0
2½	40	.47	3.2
3	48	.56	2.7
4	64	.75	2.0
5	80	.94	1.6

STORAGE AND DISPOSAL

PROHIBITIONS: ~~Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.~~

PESTICIDE STORAGE: ~~Do not contaminate water, food, or feed by storage or disposal.~~ Store at temperatures above 32°F. ~~Do not contaminate feed, foodstuffs or drinking water. Do not store next to feed or food, or transport in or on vehicles containing foodstuffs or feeds.~~ For help with any spill, leak ~~or~~ fire ~~or~~ exposure involving this material, call CHEMTREC 1-800-424-9300.

PESTICIDE DISPOSAL: This product is ~~Pesticides wastes are~~ acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). ~~Do not reuse container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning, or dispose of by other procedures if burned, stay out of smoke allowed by state and local authorities.~~

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER!

FOR BULK/MINI-BULK AND E-Z HANDLER® RECYCLABLE/REFILLABLE CONTAINERS: Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS:

REFILL ONLY WITH CYCLONE HERBICIDE. ~~The contents of this container cannot be completely removed by cleaning. Refilling with materials other than CYCLONE will result in contamination and may weaken container.~~

Container is not safe for food, feed or drinking water!

AMBUSH®, CYCLONE®, DEVRINOL®, E-Z HANDLER®, KARATE®, SURPASS® and TOPNOTCH® are trademarks of ZENECA Group Companies.

Aatrex®, Bicep®, Bicep Lite®, Caparol®, Cotoran®, Dual®, and Princep® are trademarks of CIBA-GEIGY Corporation.

Accelerate® is a trademark of Atochem North America.

Banvel®, Frontier®, Guardsman®, Marksman®, Solicam®, and Zorial® are trademarks of Sandoz Crop Protection Company.

Basagran® is a trademark of BASF Corporation.

Bladex®, Canopy®, Extrazine®, Gemini®, Harmony® Extra, Krovar®, Lexone®, Lorox®, Lorox Plus™, Preview®, Sinbar® and Velpar® are trademarks of E. I. DuPont de Nemours & Company.

Butoxone™ is a trademark of Cedar Chemical Corporation.

Butyrac®, Folex®, Modown®, and Prep™ are trademarks of Rhone-Poulenc, Inc.

Command® is a trademark of FMC Corporation.

Cotton-Pro®, Griffex®, Linex® and Meturon® are trademarks of Griffin Corporation.

DEF®, Sencor® and Turbo® are trademarks of Miles Inc.

Dropp® is a trademark of NOR-AM Chemical Company.

Goal® is a trademark of Rohm and Haas Company.

Harvade® is a trademark of Uniroyal Chemical Company.

Harness®, Lariat® and Lasso® are trademarks of Monsanto Agricultural Company.

Prowl®, Pursuit®, and Scepter® are trademarks of American Cyanamid Company.

Surflan® is a trademark of DowElanco.

Please consult each appropriate label for specific recommendations.

Made in U.S.A.
ZENECA Ag Products
ZENECA Inc.
Wilmington, DE 19807