

Reg # 10182-141

PM-23

197

DIRECTIONS FOR USE

Annual Grass and
Broadleaf Herbicide
for SoybeansFOR USE IN
ILLINOIS, INDIANA, IOWA,
KENTUCKY, MISSOURI, OHIO,
AND TENNESSEE* ONLY

*See inside cover



Agricultural Products

ICI Americas Inc.
Wilmington, Delaware 19807
A Business of ICI Americas Inc.03 0804 003P*
DE 1 90EPA Reg. No. 10182-141
A-1RPANTONE
540PANTONE
471CPANTONE
356CInk
PMS 540 (Blue)
PMS 471C (Brown)
PMS 356 (Green)*All counties except Benton, Carroll, Chesser,
Crockett, Decatur, Dyer, Fayette, Gibson,
Hardeman, Harlan, Howard, Henderson,
Henry, Lake, Lauderdale, McIntosh, Madison,
Ozon, Shelby, Tipton, and Wabash

INDEX

Precautionary Statements	2
Statement of Practical Treatment	3
Directions for Use	3
General Information	3
Application Directions	4
General Use Precautions	7
Rotational Crop Restrictions	9
TORNADO Herbicide	11
Basagran Herbicide	11
Appendix	13
Storage and Disposal	14

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions or under abnormal use conditions or under conditions not reasonably foreseeable to Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Basagran - Reg. TM of BASF Corporation

NOT REVIEWED
In Accordance with PR Notice 82-2.
Based on Draft Labeling Dated

5/24/90

BEST AVAILABLE COPY

FIRENZE & COMPANY, INC.		(302) 571-TYPE	
300 Cornett Dr., Suite A-2, Wilmington, DE 19801		FAX (302) 571-0897	
UNIT WD2	VOLUME GROM	GP XIP ICI	
FILE F8826-01	DATE 2 2 90	TIME 13 35	OPERATOR pm
DESCRIPTION MARISA DEPAULIS - ICI AMERICAS - PO 0804 - A-1R			

BEST AVAILABLE COPY

ACTIVE INGREDIENTS:

Fluazifop-P-butyl
Butyl (R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate 8.30%
Sodium salt of fomesafen
S-[2-chloro-4-[[trifluoromethyl]phenoxy]-N-(methylsulfonyl)-2-nitrobenzamide 11.50%
INERT INGREDIENTS 80.11%
TOTAL 100.0%

Contains 0.75 lbs (+) isomer of fluazifop-P-butyl and the equivalent of 11.0% fomesafen or 1 lb fomesafen active ingredient per gallon

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

This product contains fomesafen which has been determined to cause tumors in laboratory animals (mice). Risks can be reduced by closely following use directions and precautions and by wearing the protective clothing specified elsewhere on this label.

CAUSES SUBSTANTIAL, BUT TEMPORARY EYE INJURY. HARMFUL IF ABSORBED THROUGH SKIN OR INHALED. MAY CAUSE ALLERGIC SKIN RESPONSE.

Do not get in eyes or skin or clothing. Wear protective clothing, goggles or full face shield, rubber gloves and rubber apron when handling or mixing concentrate. Wear protective clothing and rubber gloves when spraying. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633)
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident Call CHEMTREC 1-800-424-9300

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

Read all label directions before using.

TORNADO is a selective early postemergence herbicide for control of annual grass and broadleaf weeds in soybeans.

TORNADO has systemic activity moving from the treated foliage into the shoots, roots, and growing points (meristematic regions) of treated grass weeds and contact activity for broadleaf weeds. Thorough coverage of all weed plant foliage is important for good activity. Optimum weed control is achieved when young actively growing weeds are treated that are not under stress

from moisture, temperature, low soil fertility, mechanical or chemical injury.

Control Symptoms: Growth of treated grass weeds stops soon after application. Symptoms include loss of vigor, yellowing and/or reddening, and eventual death to the treated grass plant. Symptoms on grass weeds are generally observed within one to three weeks, depending on species and environmental conditions.

Symptoms on broadleaf weeds occur within 3 days and appear as browning and crinkling.

Soybean plants are tolerant to TORNADO when it is applied at the recommended rate. There may be slight bronzing, crinkling or spotting of soybean leaves but soybeans soon outgrow these effects and develop normally.

APPLICATION DIRECTIONS

Timing: Best control of susceptible weeds is obtained when TORNADO is applied to actively growing young weeds before they exceed the recommended growth stages shown on this label. Refer to the weed tables for specific recommendations on weed growth stages. Generally, the application should occur 10 to 14 days after soybean emergence.

Spray Additives: Only crop oil concentrate and nonionic surfactants cleared for use on growing crops under 40 CFR 180.1001 may be used in spray mixture.

ALWAYS ADD ONE OF THE FOLLOWING:

• **Crop Oil Concentrate:** Add a nonphytotoxic crop oil concentrate at 0.5-1% (1 quart per 25 gallons) of the finished spray volume. Crop oil concentrate can improve weed control but may slightly reduce crop tolerance.

• **Nonionic Surfactant:** Add nonionic surfactant containing at least 75% surface active agent, at 0.25-0.5% (1/4-1 pint per 25 gallons) of the finished spray volume for ground sprays.

In addition to crop oil concentrate or nonionic surfactant, diammonium phosphate (aqueous ammonium polyphosphate) commonly sold as a solution (10-34-0) can be added to the spray mixture. This water soluble material should be used at a rate of 2 pts. per acre. Diammonium phosphate should not be used as a substitute for crop oil concentrate or nonionic surfactant in the spray mixture. Liquid nitrogen fertilizer (28% or similar) can also be added to spray mixture. Liquid nitrogen fertilizers are completely water soluble and should be used at a rate of one gallon per acre. Liquid nitrogen fertilizers should not be used as a substitute for crop oil concentrate or nonionic surfactant in the spray mixture.

Mixing: Fill a clean sprayer tank 1/2 full with clean water. Begin agitation and add the recommended rate of TORNADO. Add the appropriate amount of nonionic surfactant or crop oil concentrate to be used. Finish filling the tank with water to the needed volume. Allow the spray mixture to agitate and recycle 5-10 minutes before application.

Ground Application: Use sufficient spray volume and pressure to ensure complete coverage of the target weeds. A minimum of 15 gallons per acre of spray mixture should be applied with spray pressures of 40 to 60 psi at the nozzle tip. A sufficient spray volume and pressure should be used to provide complete coverage of the treated weed species. When foliage is dense, use 60 psi and a minimum of 20 gallons per acre to ensure coverage of weed foliage.

Use only hollow cone or flat fan nozzles. The sprayer must be calibrated to provide the proper volume and rate per acre. In addition, the boom and nozzle height must be adjusted to provide complete coverage of all weeds.

DO NOT USE FLOOD TYPE OR OTHER SPRAY NOZZLE TIPS WHICH DELIVER COARSE, LARGE DROPLET SPRAYS

DO NOT APPLY TORNADO WITH RECIRCULATING SPRAYERS, ROPEWICK APPLICATORS, CONTROLLED DROPLET APPLICATORS (CDA) OR ANY SIMILAR DEVICE

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM

Band Applications - Adjust band application equipment to provide thorough weed coverage. Best coverage is obtained by using a minimum of two nozzles, one directed to each side of the planted row. A single nozzle directed over the top of the row will not provide adequate coverage and is not recommended. Cultivation of untreated areas may be needed following band applications. When making band applications and cultivating in the same operation, position nozzles in advance of the cultivation device. This will reduce dust in the spray area. Dust can intercept the spray, reducing weed coverage resulting in less than adequate weed control.

Calculate the amount of herbicide and water volume needed for band treatment by the following formulas:

$$\frac{\text{Band width in inches} \times \text{broadcast rate}}{\text{row width in inches}} = \text{band herbicide rate per acre}$$
$$\frac{\text{Band width in inches} \times \text{broadcast volume}}{\text{row width in inches}} = \text{band water volume per acre}$$

Cultivation - Cultivation within 7 days prior to or within 7 days after application is not recommended as weeds may be put under stress reducing weed control. Timely cultivation 2-3 weeks after applying TORNADO may assist weed control.

GENERAL USE PRECAUTIONS

TORNADO can be applied only in the following states: Illinois, Indiana, Iowa, Kentucky, Missouri, Ohio, and Tennessee.

Applications of TORNADO should not be made beyond 3 weeks after the soybeans have emerged. A maximum of 1 quart (0.25 lb. active fomesafen + 0.188 lb. active fluazifop-P-butyl) per acre of TORNADO herbicide may be applied in alternate years. In the year of application, however, make only one application per growing season.

DO NOT APPLY FORMULATIONS CONTAINING FOMESAFEN TO ANY FIELD MORE THAN ONCE EVERY TWO YEARS

Applications of TORNADO herbicide when crop and weeds are under stress from cold temperature, when maximum day temperature is below 70°F, or when soil temperature is below 60°F, or when weeds are stressed due to low soil fertility, moisture stress, or mechanical, or chemical injury may result in reduced weed control since the weeds may not be actively growing.

TORNADO should not be applied to soybeans which have been under stress conditions such as drought, hail damage, flooding or herbicide injury as increased crop injury may result.

For mixed weed populations, apply when the first weed species reaches the recommended growth stage for treatment.

Where irrigation is used as part of normal cropping practice, best results are usually obtained when TORNADO is applied within 7 days after irrigation.

Thoroughly clean spray tank with water and a commercial tank cleaner before and after each use.

TORNADO herbicide should not be mixed with fertilizers or pesticides unless specifically recommended on this label or on other approved ICI Americas supplemental labeling.

TORNADO herbicide requires a 4-hour rain-free period for best results. Do not apply if rain is threatening.

Following soybean harvest, plow or till (moldboard or disk-plow) the soil in the fall or spring to minimize the possibility of injury to rotational crops.

Avoid overlapping spray swaths.

Avoid drift to all other crops and nontarget areas. Crops other than soybeans may be severely injured by drift.

Do not graze treated areas or harvest for forage or hay.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying TORNADO at recommended rates in soybeans.

Crop to Be Planted	Minimum Rotation Interval (Months After Last TORNADO Application)
Small grains such as wheat, barley, rye	4
Corn, cotton, peanuts, rice	10
To avoid crop injury do not plant sunflowers, sugar beets, sorghum or any other crop within	18

Do not graze rotated small grain crops or harvest for livestock forage or straw. In the event of a crop loss due to weather conditions soybeans can be replanted.

BROADLEAF WEEDS CONTROLLED WITH TORNADO IN SOYBEANS AT 1 QT/ACRE^{1,2}

Weed Growth Stages

Weeds Controlled	Maximum Growth Stage at 1 QT/Acre (No. of Leaves)
Caperweed	6" diameter
Croton, showy	4
Cucumber volunteer	4
Jimsonweed	4
Ladysthumb	2
Lambquarters, common	2: Suppression Only
Morningglory	
Entireleaf	
Integrifolia	2: Suppression Only
Lythrum, hederacea	2: Suppression Only
Purple moonflower	2
Smothering vine	2
Tan (common)	2: Suppression Only
Mustard seed	4
Nightshade, black	4
Pigweed, redroot	4
Pigweed, smooth	4
Purslane, common	6" diameter
Ragweed, common	4
Smartweed, pennsylvanian	4
Vernice, mallow	4
Yarrow, rosette	4

¹Do not apply TORNADO or other fomesafen containing compounds to any field more than once every two years.

²It is necessary to use 0.25-0.5% nonionic surfactant or 5% crop oil concentrate.

³Scientific names for weeds are listed in the Appendix (Page 12).

BEST AVAILABLE COPY

BEST AVAILABLE COPY

ANNUAL GRASS WEEDS CONTROLLED WITH TORNADO IN SOYBEANS AT 1 QT/ACRE

Weed Growth Stages		
Weeds Controlled	Maximum Height at 1 Qt/Acre (In.)	Minimum Growth Stage at 1 Qt/Acre (No. of Leaves)
Barnyardgrass	2-3	3
Broadleaf signalgrass	2-4	5
Crabgrass		
Large crabgrass	1-2	4
Smooth crabgrass	1-2	4
Southern crabgrass	1-2	4
Tropical crabgrass	1-2	4
Downy brome	2-6	4
Fall panicum	2-6	6
Fall sandbur	2-4	4
Forbes		
Giant foxtail	2-4	4
Green foxtail	2-4	4
Yellow foxtail	2-4	4
Goosegrass	2-4	6
Italian ryegrass	2-4	4
Lichgrass	4-24	6
Johnsongrass seedling	2-8	4
Jungle rice	2-3	3
Shattercane	8-12	8
Sorghum	8-12	8
Southern sandbur	2-6	6
Teas panicum	8	6-12
Volunteer cereals		
V. barley	2-6	6
V. corn	12-24	10
V. milo	6-12	4
V. oats	2-6	6
V. rye	2-6	6
V. wheat	2-6	6
Wild proso millet	4-8	6
Witch grass	2-4	6
Wild oats	2-6	6
Woolly cupgrass	2-4	6

TORNADO HERBICIDE/ BASAGRAN HERBICIDE

Postemergence Control of Grass and Broadleaf Weeds

TORNADO herbicide and Basagran herbicide may be used together in a postemergence program for broad-spectrum weed control in soybeans.

TORNADO herbicide and Basagran herbicide may be applied sequentially or in tank-mix. The growth stage of weeds at the time of application will determine which method of application will provide the most satisfactory results. Both TORNADO and Basagran herbicide should be applied to actively growing weeds. Read and observe all applicable label directions and limitations for both TORNADO herbicide and Basagran herbicide before using.

METHOD 1: Sequential Application TORNADO Herbicide Followed by Basagran Herbicide

Apply TORNADO herbicide and an approved spray additive at the recommended rate and growth stage for the grass and broadleaf weeds being treated according to label recommendations. To allow adequate time for the systemic activity of TORNADO to translocate to the meristematic regions of the grass plants, delay application of Basagran for at least 24 hours following the TORNADO application. Basagran can then be applied to susceptible broadleaf weeds following the Basagran label recommendations.

METHOD 2: Sequential Application Basagran Herbicide Followed by TORNADO Herbicide

Apply Basagran herbicide at the recommended rate and growth stage for susceptible broadleaf weeds following the Basagran

label recommendations. Delay application of TORNADO for at least 24 hours following the Basagran application. TORNADO must be applied to actively growing weeds for best results. Apply TORNADO herbicide and an approved adjuvant at the recommended rate and growth stage for the grass and broadleaf weeds being treated according to label recommendations.

METHOD 3: Tank Mix Applications TORNADO Herbicide and Basagran Herbicide

A tank mix of TORNADO herbicide and Basagran herbicide may be applied using a 1 Qt/Acre rate of TORNADO. Basagran should be added to the mixture at the recommended rates for growth stages described on the Basagran label. Use a non-ionic surfactant or crop oil concentrate in the spray mixture.

A minimum of 15 gallons per acre of spray mixture should be used with spray pressures of 40 to 60 psi at the nozzle tip. When foliage is dense, use 60 psi and a minimum of 20 gallons per acre to ensure coverage of weed foliage. Load the spray tank with half the amount of required water and add the recommended amounts of TORNADO herbicide, Basagran herbicide and crop oil concentrate while the agitator is running and then add the remaining quantity of water. Do not make more than one application of this tank mix per season.

Note: Tank mix applications sometimes have resulted in reduced grass weed control. A tank mix application is not recommended if perennial grass weeds are the predominant grasses to be controlled. If grass regrowth occurs following an application of the tank mix or an additional flush of grasses emerges, make an application of FUSILADE® 2000 herbicide to actively growing grass weeds, according to label recommendation.

APPENDIX

Scientific names are listed for annual grass and broadleaf weeds referred to in the TORNADO label. Specific weed control recommendations can be found in specific rate tables.

COMMON NAME	SCIENTIFIC NAME
Broadleaf Weeds	
Carpetweed	Mollugo verticillata
Crotalaria showy	Crotalaria spectabilis
Cucumber volunteer	Cucumis sativus
Jimsonweed	Datura stramonium
Ladythumb	Polygonum persicaria
Lambsquarters common	Chenopodium album
Morning glory	
Entireleaf	Ipomoea hederacea var. integrifolia
Ivyleaf	Ipomoea hederacea var. hederacea
Purple moonflower	Ipomoea tubiflora
Small white (pitted)	Ipomoea acuminata
Tail (common)	Ipomoea purpurea
Mustard wild	Brassica kaber
Nightshade black	Solanum nigrum
Pigweed redroot	Amaranthus retrofractus
Pigweed smooth	Amaranthus hybridus
Purslane common	Portulaca oleraceae
Ragweed common	Ambrosia artemisiifolia
Smartweed pennsylvanian	Polygonum pennsylvanicum
Vernice mallow	Hibiscus trionum
Yellow rocket	Barbarea vulgaris
Annual Grass Weeds	
Barnyardgrass	Echinochloa crusgali
Broadleaf signalgrass	Bracharia distachya
Crabgrass	
Large	Digitaria sanguinalis
Smooth	Digitaria ischaemum
Southern	Digitaria ciliaris
Tropical	Digitaria bicornis
Downy brome	Bromus tectorum

BEST AVAILABLE COPY

COMMON NAME	SCIENTIFIC NAME
Annual Grass Weeds (Continued)	
Fall panicum	Panicum dichotomiflorum
Field sandbur	Cenchrus incertus
Forbale	
Green	Setaria faberii
Green	Setaria viridis
Yellow	Setaria italica
Goosegrass	Elymus indica
Italian ryegrass	Lolium multiflorum
Itchgrass	Rottboellia exaltata
Johnsongrass seedling	Sorghum halepense
Jungle rice	Echinochloa colonum
Shallfence	Sorghum bicolor
Sorghum	Sorghum alatum
Southern sandbur	Cenchrus echinatus
Texas panicum	Panicum texanum
Volunteer cereals	
V. barley	Hordeum vulgare
V. corn	Zea mays
V. milo	Sorghum bicolor
V. oats	Avena sativa
V. rye	Secale cereale
V. wheat	Triticum aestivum
Wild proso millet	Panicum miliaceum
Witchgrass	Panicum capillare
Wild oats	Avena fatua
Woody cupgrass	Echinochloa villosa

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. **STORAGE:** Store above 32°F in original containers only. If product freezes, return to room temperature and agitate to reconstitute. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL: Metal Containers:** Triple-rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. **Plastic Containers:** Triple-rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Basagran—Reg. TM of BASF Corporation
Tornado—Reg. TM of an ICI Group Company

*A. Counties except Benton, Carroll, Chester, Chickasaw, Decatur, Dyer, Fayette, Gibson, Hardeman, Hardin, Haywood, Henderson, Henry, Lake, Lauderdale, McNairy, Madison, Marion, Shelby, * Pike and Weakley.

697

NOT REVIEWED
In Accordance with FR Notice 82-2.
Based on Draft Labeling Dated

5/24/90



ANNUAL GRASS AND BROADLEAF HERBICIDE FOR SOYBEANS

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions or under abnormal use conditions or under conditions not reasonably foreseeable to Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

This product contains fomesafen which has been determined to cause tumors in laboratory animals (mice). Risks can be reduced by closely following use directions and precautions and by wearing the protective clothing specified elsewhere on this label.

CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY. HARMFUL IF ABSORBED THROUGH SKIN OR INHALED. MAY CAUSE ALLERGIC SKIN RESPONSE.

Do not get in eyes or skin or clothing. Wear protective clothing, goggles or full face shield, rubber gloves and rubber apron when handling or mixing concentrate. Wear protective clothing and rubber gloves when spraying. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-80-E-8 (327-0633)
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident Call CHEMTREC 1-800-424-9300

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to Supplemental Label for directions for use. This labeling must be in the possession of the user at the time of pesticide application.

Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. **STORAGE:** Store above 32°F in original containers only. If product freezes, return to room temperature and agitate to reconstitute. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or residue is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL: Metal Containers:** Triple-rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. **Plastic Containers:** Triple-rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke. **Glass Containers:** Triple-rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.



**KEEP OUT
OF REACH
OF CHILDREN
WARNING**

See Side Panel for Additional
Precautionary Statements and
Statement of Practical Treatment

EPA Reg. No. 10182-141
EPA Est. No. 34704-MS-1
U.S. Patent No. 4,285,723

Net Contents:
2½ U.S. GALLONS
(9.46 LITERS)

**ANNUAL GRASS AND BROADLEAF
HERBICIDE FOR SOYBEANS**

ACTIVE INGREDIENT:

Fluazifop-P-butyl
butyl (R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyloxy]
phenoxy]propanoate
Sodium salt of fomesafen
5-[2-chloro-4-(trifluoromethyl)phenoxy]N-(methylsulfonyl)-2-nitrobenzamide

8.30%

11.59%

INERT INGREDIENTS

80.11%

TOTAL

100.0%

Contains 0.75 lbs (+) isomer of fluazifop-P-butyl and the equivalent of 11.07- fomesafen or
1 lb fomesafen active ingredient per gallon



Agricultural Products

ICI Americas Inc.
Wilmington, Delaware 19897

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to
Supplemental Label for directions for use. This labeling must be in the possession of the user at the time
of pesticide application.
Do not apply this product through any type of irrigation system.

