

JUN 1993

Ms. Becky Rhodes
SENECA Ag Products
P.O. Box 751
Wilmington, DE 19897

Dear Ms. Rhodes:

Subject: CAPTAN 50 WP
EPA Registration No. 10182-145
Application Dated May 18, 1993, Request To
Amend Labeling To Reflect Residue Chemistry
Data Supporting Food Uses of Product

The subject labeling submitted in connection with registratin under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) as amended is acceptable provided that you:

1. Either repeat the entire definition of crop cycle or delete the phrase "including postharvest sprays". on pages 8, 9 and 10. The redundancy could confuse the reader in regard to the term "crop cycle".
2. On page 7, line 3 add the word "in" to complete the sense of "equipment or in 5 to 20 gallons of water by air". Under "Apricots", line 6 add the word "per" to complete the sense of "...acre per crop cycle".
3. Revise the Environmental Hazards statements in compliance with PR Notice 93-3.
4. Submit five (5) printed copies of the final printed label before releasing the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section 5(4). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your record.

Sincerely yours,

BEST AVAILABLE COPY

Joanne L. Miller
Product Manager (23)
Fungicide-Herbicide Branch

2815

Captan 50-WP

A Fungicide For Plant Disease Control

ACTIVE INGREDIENTS

*Captan	48.9%
Related Derivatives	1.1%

INERT INGREDIENTS	<u>50.0%</u>
TOTAL	100.0%

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

Captan 50-WP is a microfine wettable powder for use in water as a spray for the control of certain fungal diseases of fruit, and ornamental crops, and as a soil treatment for the control of certain seed rots and damping-off diseases.

EPA Reg. No. 10182-145
EPA Est. No.

___ LBS. NET ___ KG

KEEP OUT OF REACH OF CHILDREN

**D A N G E R
P E L I G R O**

PRECAUCION AL USUARIO:

Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

CAUSES IRREVERSIBLE EYE DAMAGE

Made in U.S.A.
ZENECA Ag Products
ZENECA Inc.
Wilmington, DE 19897

ACCEPTED
with COMMENTS
EPA Letter Dated:

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this pesticide
registered under EPA Reg. No.

10182-145

2815

STATEMENT OF PRACTICAL TREATMENT

FIRST AID

If a known exposure occurs or is suspected, immediately start the procedures given below and contact a POISON CENTER, PHYSICIAN, OR THE NEAREST HOSPITAL. Describe the type and extent of exposure, the victim's symptoms and follow the advice given.

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if abnormal reaction occurs.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL 1-800-F-A-S-T-M-E-D (327-8633)

FJR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

For terrestrial uses,
This pesticide is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources.

X
Direct
93-3

STORE IN COOL, DRY PLACE -- PROTECT FROM EXCESSIVE HEAT

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States

5/15

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed federal tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to adjoining food, fiber or pasture crops. Drift of Captan onto sensitive crops (e.g., D'Anjou Pears) can cause severe phytotoxicity and crop loss.

Consult State Agricultural Experiment stations or State Agricultural Extension Service for additional information, as the time of applications needed will vary with the local conditions.

COMPATIBILITY AND PLANT SAFETY: Captan 50-WP can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and bordeaux mixture will reduce the fungicidal activity of Captan. Do not apply Captan in combination with or immediately before or closely following oil sprays. Do not allow oil sprays on adjacent crops to drift onto crops which have been or will shortly be treated with Captan. The time factor governing the safe interval between Captan and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of Captan and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, Captan may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums and cherries. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use Captan and other spray materials at lowest recommended rates and avoid drenching trees.

Applications can be made by aircraft or ground power equipment (including concentrate and semi-concentrate equipment). Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read compatibility and plant safety information).

For aerial or concentrate spray applications, apply the same amount of Captan 50-WP per acre as would normally be applied for dilute spray applications. Apply aerial or concentrate sprays in sufficient water for coverage. Do not apply this product through any type of irrigation system.

BEST AVAILABLE COPY

6/15

USE PRECAUTIONS

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

A NIOSH- or MSHA-approved dust mask must be worn when mixing and loading this product. When mixing and loading this product, handlers must wear a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Field workers and harvesters must wear chemical resistant gloves if entering treated areas within 4 days following applications. Leather or fabric gloves are not acceptable.

During application, even from a tractor with a completely enclosed cab or from an aircraft with an enclosed cockpit, a long-sleeved shirt and long-legged pants must be worn. Chemically resistant gloves must be available in the cab or cockpit and worn while exiting.

Only the applicator is permitted to be in the greenhouse during application of Captan. Open vents to greenhouse during application and for at least 1 hour after application.

Workers planting in Captan-treated soil must wear chemical resistant gloves.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with Captan must be destroyed according to state and local regulations.

REENTRY

- Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.
- Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn. Conspicuously post reentry information at site of application.
- Certain states may require more restrictive reentry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information.
- Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

BEST AVAILABLE COPY

GENERAL USE PRECAUTIONS

Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. ~~Maximum application is for a crop cycle. Crop cycle is defined as prebloom through postharvest.~~ Apply the high rate ~~and/or spray at shorter intervals~~ when climatic conditions most favor disease(s). Apply the low rate ~~and/or spray at larger intervals~~ when climatic conditions least favor disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions. ✓

IMPORTANT: Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the eastern or western U.S. Follow those directions for your growing area where specified.

FRUIT AND NUT CROPS

ALMONDS: Brown rot twig and blossom blight, shothole, scab, leaf blight -- Apply 4 to 9 pounds Captan 50-WP per acre in 20 to 300 gallons of water using ground equipment or 5 to 20 gallons of water by air. Use 6 to 9 pounds per acre when Captan is used alone. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, Captan 50-WP may be used in a tankmix at a rate of 4 to 6 pounds per acre. Apply at popcorn, bloom and petal fall stages and up to 5 weeks after petal fall. Do not apply within 30 days of harvest. Hulls may be fed to livestock. ~~Subsequent applications may be made up to 12 days of harvest in accordance with pest infestations and state agricultural extension service recommendations, provided the hulls are not fed to dairy animals or animals being finished for slaughter.~~ Do not apply more than 45 pounds per acre per crop cycle (see GENERAL USE PRECAUTIONS section on Page ____ for definition of crop cycle). ✓

APPLES (Eastern U.S.): Primary scab, black rot (frog-eye), botrytis blossom-end-rot -- Apply 6 pounds Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment or 5 to 20 gallons of water by air. Apply at 5 to 7 day intervals as needed to maintain control in prebloom, bloom, petal fall and first cover sprays.

Secondary scab, Brooks fruit spot, sooty blotch, fly speck, black rot, black pox, botryosphaeria rot, bitter rot -- Apply 4 to 6 pounds Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment or 5 to 20 gallons of water by air. Apply at 10 to 14 day intervals in second and later cover sprays. Do not apply within 14 days of harvest. Do not apply more than 64 pounds of Captan 50-WP per acre per crop cycle.

Powdery mildew -- If powdery mildew is a problem, add 6 to 12 pounds sulfur per acre to all postbloom sprays until foliage matures. CAUTION -- Do not use Captan 50-WP in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin, King, etc., as severe injury and defoliation may occur.

APPLES (Western U.S., except CA): Primary scab -- Apply 6 pounds Captan 50-WP per acre in a high volume application for preblossom sprays. In mid-summer, the dosage may be reduced

BEST AVAILABLE COPY

X to 4 pounds per acre in high volume applications. (Pacific northwest): Bull's eye rot, Botrytis rot -- Apply 6 pounds Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment or 5 to 20 gallons of water by air. Make 1 or 2 applications with late cover sprays and 1 final spray prior to harvest. Do not apply within 14 days of harvest. Do not apply more than 64 pounds of Captan 50-WP per acre per crop cycle.

APPLES (California): Primary scab -- for prebloom through petal fall sprays, apply 4 to 6 pounds Captan 50-WP in 20 to 400 gallons of water per acre using ground equipment or 5 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, the lower rate may be used in tank mixtures.

Secondary scab -- In mid-summer cover sprays, the dosage may be reduced to 4 pounds per acre. Do not apply within 21 days prior to harvest. Do not apply more than 30 pounds of Captan 50-WP per acre per crop cycle.

) **APPLES (SEVERE SUMMER DISEASE DEVELOPMENT) (Arkansas, Louisiana, Missouri, Texas and states east of the Mississippi River Only):** Upon recommendation of the local cooperative extension office, during periods highly favorable for severe summer disease development (black rot, bitter rot, sooty blotch, flyspeck, etc.), apply 8 pounds of Captan 50-WP per acre on a 7 to 10 day schedule. Do not apply more than 64 pounds of Captan 50-WP per acre per crop cycle. May be applied up to day of harvest.

APRICOTS: Brown rot (twig blight), jacket rot -- Apply 3 to 5 pounds Captan 50-WP per acre in 20 to 250 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in red bud, bloom 75% petal fall, and cover sprays.

To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rate in tank mixtures. Do not apply within 7 days prior to harvest. Do not apply more than 25 pounds per acre per crop cycle.

) **BLUEBERRIES (Eastern U.S.):** Botrytis gray mold or berry rot, mummy berry - Apply 5 pounds Captan 50-WP per acre in sufficient water for thorough coverage or a minimum of 5 gallons of water by air. Start spray program when buds swell or when buds have loose scales. Repeat at 7 day intervals through blossom period. Repeat at 7 to 10 day intervals from late bloom. Do not apply more than 70 pounds per acre per crop cycle.

BLUEBERRIES (Western U.S., except CA): Botrytis gray mold or berry rot, mummy berry - Apply 2 to 5 pounds Captan 50-WP per acre in sufficient water for thorough coverage or a minimum of 5 gallons of water by air. Begin at mid-bloom, repeat at 7 to 10 day intervals until maturity. Do not apply more than 70 pounds per acre per crop cycle.

BLUEBERRIES (California): Botrytis gray mold or berry rot, mummy berry -- Apply 2 to 5 pounds Captan 50-WP per acre in 20 to 200 gallons of water by ground or 5 to 20 gallons of water by air. Begin at mid-bloom, repeat at 7 to 10 day intervals until maturity. Do not apply within 4 days of harvest. Do not apply more than 25 pounds per acre per crop cycle.

CHERRIES (Eastern U.S.): Brown rot, leaf spot, Botrytis rot -- Apply 4 pounds Captan 50-WP per acre in 20 to 200 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in prebloom, bloom, petal fall, shuck, cover and pre-harvest sprays. Applications at 3 to

9315

4 day intervals may be necessary during bloom to control blossom blight. Repeat applications at 7 to 20 day intervals as needed to maintain control up to start of harvest. If powdery mildew is a problem, add 6 lbs. sulfur per acre to the petal fall, shuck or early cover sprays. If sulfur is added, Captan 50-WP may be reduced to 2 pounds per acre in these sprays. Do not apply more than 28 pounds per acre per crop cycle (including postharvest sprays).

Postharvest sprays: leaf spot -- Apply 4 pounds Captan 50-WP per acre in 20 to 200 gallons of water using ground equipment. Apply immediately after harvest and repeat application in 10 to 14 days.

thru to post harvest sprays
in leafy

CHERRIES (Western U.S., except CA): Brown rot blossom blight, brown rot (fruit), leaf spot -- Apply 4 pounds Captan 50-WP per acre in 20 to 200 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in prebloom, bloom, petal fall, shuck, cover and preharvest sprays. Do not apply more than 28 pounds per acre per crop cycle.

CHERRIES (California): Brown rot blossom blight, brown rot (fruit), leaf spot -- Apply 3 to 4 pounds Captan 50-WP per acre in 20 to 200 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in prebloom, bloom, petal fall, shuck, cover, and preharvest sprays. Do not apply more than two late cover preharvest sprays. Do not apply within 7 days of harvest. Do not apply more than 28 pounds per acre per crop cycle.

GRAPES (U.S., except CA): Phomopsis cane and leaf spot, downy mildew, suppression of black rot -- Apply 2 to 4 pounds Captan 50-WP per acre in 20 to 200 gallons water using ground equipment or 7 to 20 gallons water by air; when shoots are 1/2 to 1 1/2 inches long, when shoots are 3-5 inches long, and when shoots are 9-12 inches long. Repeat just before bloom, immediately after bloom, and continue at 10-14 day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible grape varieties and during periods of weather highly favorable for disease development. Do not apply more than 24 pounds of Captan 50-WP per acre per crop cycle. Do not apply within 14 days of harvest.

GRAPES (California): Bunch rot (Botrytis) -- Apply 4 pounds Captan 50-WP per acre in 20 to 200 gallons of water using ground equipment, or 7 to 20 gallons of water by air. Make 2 applications before bloom and 1 immediately after bloom. Repeat periodically, making 3 cover applications before the bunches close. Phomopsis cane and leaf spot (current season infection) - Apply 3 to 4 pounds Captan 50-WP per acre in 20 to 200 gallons of water using ground equipment or apply 4 pounds Captan 50-WP per acre in 7 to 20 gallons of water by air. Apply first spray when green tissue begins to show but before shoots are 1 inch long and repeat application when shoots are 6 to 8 inches long. Do not apply more than 24 pounds of Captan 50-WP per acre per crop cycle. Do not apply within 45 days of harvest. Do not conduct cultural practices which require intensive foliar contact (tipping, cluster thinning, cane throwing) within 7 days after application.

NECTARINES (U.S., except CA): Brown rot, scab -- Apply 8 pounds Captan 50-WP per acre in 20 to 250 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain control. Continue applications throughout harvest if conditions favor brown rot. If powdery mildew is a problem, add 7 1/2 pounds sulfur per acre to the

petal fall, shuck and early cover spray. If sulfur is added, Captan 50-WP may be reduced to 2½ pounds per acre in these sprays.

Coryneum blight (peach blight, shothole) -- Apply 8 pounds Captan 50-WP per acre in 20 to 250 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in pink bud, full bloom, petal fall and cover sprays as necessary and as a post harvest spray (but before leaves drop).

Do not apply more than 48 pounds per acre per crop cycle (including postharvest sprays).

NECTARINES (California): Brown rot, scab -- Apply 4 to 8 pounds Captan 50-WP per acre in 20 to 250 gallons of water using ground equipment or 10 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rate of Captan 50-WP in tank mixes. Apply in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as conditions favor brown rot. If powdery mildew is a problem, add 7½ pounds sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, Captan 50-WP may be reduced to 2½ pounds per acre in these sprays.

Coryneum blight (peach blight, shothole) -- Apply 4 to 8 pounds Captan 50-WP per acre in 20 to 250 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in pink bud, full bloom, petal fall and cover sprays as necessary and as a postharvest spray (but before leaves drop).

Do not apply within 7 days of harvest. Do not apply more than two late cover/preharvest sprays. Do not apply more than 40 pounds per are per crop cycle (including postharvest sprays).

PEACHES (U.S., except CA): Brown rot, scab -- Apply 8 pounds Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in full pink, bloom, petal fall, shuck stages and in cover and preharvest sprays. When conditions are favorable, make applications at 3 to 4 day intervals ~~may be necessary~~ during bloom to control blossom blight. Then repeat application at 7 to 14 day intervals as needed to maintain control (Refer to GENERAL USE PRECAUTIONS Section on Page ____ for guidance on intervals and rates). Continue applications through harvest if conditions favor brown rot. If powdery mildew is a problem, add 12 pounds sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, Captan 50-WP may be reduced to 4 pounds per acre in these sprays.

Coryneum blight (peach blight, shothole) -- Apply 8 pounds Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in pink bud, full bloom, petal fall stages and cover sprays as necessary and as a postharvest spray (but before leaves drop).

Do not apply more than 64 pounds per acre per crop cycle (including postharvest sprays).

PEACHES (California): Brown rot, scab -- Apply 4 to 8 pounds Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment or 10 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rates of Captan 50-WP in tank mixes. Apply in full pink, bloom, petal fall, shuck

11/3/15

stages and in cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. If powdery mildew is a problem, add 12 pounds sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, Captan 50-WP may be reduced to 4 pounds per acre in these sprays.

Coryneum blight (peach blight, shothole) -- Apply 8 pounds Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in pink bud, full bloom, petal fall stages and cover sprays as necessary and as a postharvest spray (but before leaves drop).

Do not apply within 7 days of harvest. Do not apply more than two late cover/preharvest sprays. Do not apply more than 40 pounds per acre per crop cycle (including postharvest sprays).

) **PLUMS, FRESH PRUNES (Eastern U.S.):** Brown rot -- Apply 6 pounds Captan 50-WP per acre in 20 to 300 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. The addition of a neutral spreader has improved coverage. Do not apply more than 54 pounds per acre per crop cycle.

PLUMS, FRESH PRUNES (Western U.S., except CA): Brown rot -- Apply 6 pounds Captan 50-WP per acre in 20 to 300 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply at green bud, popcorn, bloom and petal fall stages. Repeat in cover sprays as conditions warrant. ~~Prune russet scab (lacy scab) (in California only) -- Apply 6 pounds Captan 50-WP per acre in 20 to 300 gallons of water using ground equipment. Apply at full bloom. Do not apply more than 54 pounds per acre per crop cycle.~~

) **PLUMS, FRESH PRUNES (California):** Brown Rot -- Apply 4 to 6 pounds Captan 50-WP per acre in 20 to 300 gallons of water using ground equipment or 10 to 20 gallons of water by air. Use the lower rates when tank mixes with fungicides of similar spectrum of activity are used. Apply at green bud, popcorn, bloom and petal fall stages. Repeat in cover sprays as conditions warrant. ~~Prune russet scab (lacy scab) -- Apply 4 to 6 pounds Captan 50-WP per acre in 20 to 300 gallons of water using ground equipment. Apply at full bloom. Do not apply within 7 days of harvest. Do not apply more than two late cover/preharvest sprays. Do not apply more than 30 pounds per acre per crop cycle.~~

STRAWBERRIES (U.S., except CA): Botrytis (gray mold), leaf spot -- Apply 6 pounds Captan 50-WP per acre in sufficient water for thorough coverage by ground equipment or 10 to 20 gallons of water by air. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at weekly intervals. Under conditions favorable to fruit rot, continue applications through harvest period, treating immediately after each picking. Do not apply more than 48 pounds per acre per year.

STRAWBERRIES (California): Botrytis (gray mold), leaf spot -- Apply 3 to 5 pounds Captan 50-WP per acre in sufficient water for thorough coverage by ground equipment or 10 to 20 gallons of water by air. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 10 to 14 day intervals as disease conditions warrant. Under conditions

BEST AVAILABLE COPY

favorable to fruit rot, continue applications through harvest period treating immediately after picking. Do not apply within 48 hours prior to harvest. Do not apply more than 32 pounds per acre per year.

OK ~~ok~~
SPECIAL USES

PEACH (NURSERY STOCK) PREPLANT ROOT DIP (California): Preventative preplant dip treatment for crown gall. Use 4 pounds Captan 50-WP plus 3.2 pints diluted sodium hypochlorite (5.25% household bleach) per 100 gallons water. Wash nursery trees to remove soil from roots. Cut off all dormant buds and suckers in crown area and prune root system if necessary. Submerge the entire dormant tree for 5 minutes. ~~Recharge dip during operation as necessary to maintain a proper concentration of 200 ppm chlorine. Check the concentration frequently using a chlorine test kit.~~ Recharge dip during operation at a rate of 3.2 pints diluted sodium hypochlorite per 100 gallons of water.

) **POST HARVEST DIP FRUIT APPLICATION:** For control of various molds and storage rots (Botrytis, Gleosporium, Rhizopus). Use as a post harvest dip or spray wash on the following fruits: Apples, Cherries, Pears -- Use 2½ pounds Captan 50-WP per 100 gallons of water. Apply as a spray or in a dip tank. When used as a dip, recharge wash solution periodically when tank volume is reduced by 25%. Bring water back to volume and add 2½ pounds Captan 50-WP for each 100 gallons added. At end of every 8 hour shift, empty tank, flush and charge with fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operation.

DISPOSAL OF LEFTOVER POST HARVEST TREATMENT MIXTURE: Leftover dip or spray mixtures containing Captan may be used as a foliar spray for the same crop in case of apples and cherries (but not pears) as treated by the dip or spray mixture, or to registered turf and ornamentals sites; observing all restrictions such as maximum pounds applied per application and season.

) When calculating application rates, if analytical services are not available to determine the exact quantity of Captan remaining in the mixture, assume that all Captan which was added to the dip or spray tank during the treatment period still remains in the tank. If the dip or spray mixture contains other pesticides in addition to Captan, refer to the product label(s) for information regarding disposal.

Captan wastes are acutely hazardous to the eyes. Improper disposal of spray or dip tank mixtures is a violation of Federal Law. If the leftover dip or spray mixture cannot be disposed of in the manner prescribed above, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance concerning the disposal of spent or excess dip tank mixtures.

BEST AVAILABLE COPY

ORNAMENTALS

USE PRECAUTIONS

Do not apply spray to ornamental plants listed below beyond the point of drip from the leaf surface. When applying as a drench, apply only sufficient mixture to wet the surface of the soil except when the dose is specified in terms of volume of mixture per square foot of area.

AZALEAS: Damping-off of cuttings -- Use 4 pounds Captan 50-WP per 100 gallons of water. Dip cuttings before bedding. Petal blight -- Use 2 lbs. Captan 50-WP per 100 gallons of water. Apply to soil area around plants and spray flowers just before bloom. Repeat at 7 to 14 day intervals through bloom.

BEGONIAS (Tuberous): Damping-off, tuber rot -- Use 4 pounds Captan 50-WP per 100 gallons of water. Dip tubers 30 minutes, drain and plant.

CAMELLIAS: Petal blight -- Use 1 pound Captan 50-WP per 100 gallons of water. Apply to drench soil around plants beginning when flowers start to open. Repeat at 7 to 10 day intervals through bloom.

CARNATIONS: Alternaria leaf spot, rust -- Use 2 pounds Captan 50-WP per 100 gallons of water. Begin application at first sign of disease. Repeat at 7 to 10 day intervals. Shorten intervals during frequent rains and heavy dews. Damping-off of cuttings -- Use 4 pounds Captan 50-WP per 100 gallons of water. Dip cuttings before bedding.

CHRYSANTHEMUM: Botrytis flower blight, Septoria leaf spot -- Use 2 pounds Captan 50-WP per 100 gallons of water. Apply at first sign of disease. Repeat at 7 to 10 day intervals. Damping-off of cuttings -- Use 4 lbs. Captan 50-WP per 100 gallons of water. Dip cuttings before bedding.

DICHONDRA (California Only): White mold (*Sclerotium rolfsii*) -- Use 2 pounds Captan 50-WP per 100 gallons of water. Apply 1 gallon spray per 10 square feet. Make 2 to 3 applications at 7 day intervals.

GLADIOLUS (Corms): Corm rot and decay, damping-off -- Use ½ to 1½ pounds Captan 50-WP per 10 gallons of water, dip corms 20 to 30 minutes. Drain and plant.

GRASSES (Ornamental in Non-Pastured Areas Only): Leaf spot, damping-off, brown patch, melting out, seedling blights, and brown spot on St. Augustine grass -- Use 2 pounds Captan 50-WP per 100 gallons of water. Apply 10 gallons spray per 1,000 square feet. Begin when growth starts in spring. Repeat at 7 to 14 day intervals throughout season. Do not graze treated areas or feed clippings to livestock.

GRASSES (Lawn Seedbeds): Damping-off and other soil borne diseases -- Use 1/4 to 1/3 pound Captan 50-WP per 1,000 square feet or 2 pounds Captan 50-WP per 100 gallons of water, using 15 gallons spray per 1000 square feet. Cultivate into upper 3 to 4 inches before planting.

BEST AVAILABLE COPY

14515

ROSES: Black spot, Botrytis blossom blight -- Use 2 pounds Captan 50-WP per 100 gallons of water. Begin at first growth or first sign of disease. Repeat at 7 to 14 day intervals, and more frequently during frequent rains and heavy dews.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplant treatment for damping-off, root rot diseases on seedlings or transplants of roses (and other shrubs, trees, flowers) and lawn seedbeds -- Use 2 pounds Captan 50-WP per 100 gallons of water at a rate of 1½ gallons of spray per 1,000 square feet. Cultivate into upper 3 to 4 inches of soil before planting.

Only the applicator is permitted to be in the greenhouse during application of Captan. Open vents to greenhouse during application and for at least 1 hour after application.

Workers planting in Captan-treated soil must wear chemical resistant gloves.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep pesticide in original container. Keep container tightly closed when not in use. Protect from excessive heat. Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

) **CONTAINER DISPOSAL:** Do not reuse container. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

) **NOTICE TO BUYER AND USER:** Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.