Page 1 of 5 Captan 75 SP RS-082389

CAPTAN 75 SEED PROTECTANT

SLURRY APPLICATION FOR SEED DISEASE CONTROL

DYE ADDED

ACTIVE INGREDIENTS:

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

This product contains the toxic inert ingredient Rhodamine B.

KEEP OUT OF REACH OF CHILDREN

DANGER

CAUSES IRREVERSIBLE EYE DAMAGE

STATEMENT OF PRACTICAL TREATMENT

Immediately start the procedures given below and CONTACT AMPOISON CENTER, A PHYSICIAN OR THE NEAREST HOSPITAL. Describe the type and extent of exposure, the victim's symptoms, and follow the advice given.

If in eyes: Flush with plenty of water. Call a physician

If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water.

Avoid alcohol.

If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention if abnormal reaction occurs.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL (800)-F-A-ST-M-E-D(327-8633).
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, CALL CHEMITREC 1-800-424-9300.

EPA Reg. No. 10182-147

EPA Est. No.

50 LBS. WET 22.7 Kg

Hade In U.S.A. by
ICI Agricultural Products
ICI Americas Inc.
Wilmington, DE 19897

ACCEPTED SEP 7:8 1989

Under the Federal Insecticities Fungicials, and Rodentic to Administration of American arrange independent of the Parkers No. 10/122 - 147

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Avoid inhalation of vapor. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

BEST, AVAILABLE COPY

This product is toxic to fish. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CAPTAN 75 SEED PROTECTANT is a microfine wettable powder especially formulated for the treatment of seed, prior to storage and planting, to protect seed from molds and other fungi causing storage loss and to protect seed from seed-borne and soil-borne fungi which cause seed decay and seedling blights.

COMPATIBILITY: CAPTAN 75 SEED PROTECTANT can be used in combination with most commonly used seed protectants, both insecticides and fungicides, as well as fertilizers. Do not use with strongly alkaline materials.

PLANT SAFETY: At the recommended dosage CAPTAN 75 SEED PROTECTANT is non-injurious from the standpoint of seed injury or yield reduction. The germination of treated seed is frequently stimulated, giving a superior result as compared to results from other seed fungicides.

USE PRECAUTIONS:

Read all label instructions before using and use only as specified on this label.

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

A dust mask must be worn when mixing and loading this product.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Fieldworkers and harvesters must wear chemical resistant gloves. Leather or fabric gloves are not acceptable.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

SPECIAL USE PRECAUTIONS:

IMPORTANT: Treated seed must not be used for human consumption nor for animal feed.

The Federal Government and most States have seed laws or regulations requiring that an appropriate caution statement be affixed to bags containing treated seed, such as: "Treated Seed - Do Not Use for Feed, Food or Oil Purposes."

SLURRY TREATMENT: Pre-mix the recommended amount of this material with a little water in a bucket and make a thin paste. Stir this paste into the required amount of water for the slurry equipment to be used.

NOTE: Seed treated by the slurry method should not be bagged until such time as the seed has had an opportunity to dry. Store in well ventilated areas. If seed is bagged when wet, heat is developed and impairs seed germination.

RECOMMENDATIONS:

For control of many seed rots, seedling rots, root rots, fungi and kernel smut of sorghum.

AMOUNT OF CAPTAN 75 SEED PROTECTANT TO USE ON SEEDS

SEED	SLURRY
Alfalfa, Clover, Lespedeza, Trefoil	5-1/2 oz./100 lb.
Barley, Oats, Wheat	2/3 oz./bu.
Beans (dry, snap, lima), Cowpeas, Peas,	l oz./bu.
Beets (garden)	8 oz./100 ib.
Cantaloupe, Cucumber	2 oz./100 lb.
Cole Crops (Broccoli, Brussels Sprouts,	
Cabbage, Cauliflower)	1 oz./100 1b.

Corn - Field	2/3 oz./bu.
Sweet	2-1/2 oz./100 lb.
Cotton - Acid delinted	2 oz./100 lb.
Reginned	3 oz./100 lb.
Fuzzy	3-1/2 oz./100 lb.
Crucifers (Mustard, Radishes, Rape, Turnips)	1 oz./100 lb.
Flax	1-1/3 oz./bu.
Grasses	5-1/2 oz./100 lb.
Muskmelon, Pumpkin, Squash, Watermelon	1-1/3 oz./100 lb.
Onions (pelleting)	1 oz./1 lb.
Peanuts	4 oz./100 lb.
Soybeans	l oz./bu.
Spinach	4 oz./100 lb.
Sugar Beets - West	4 oz./100 lb.
East	0 - (100.11)
	8 oz./100 lb.

Further specific information of the subject of seed treatment may be obtained from State Agricultural Experimental Stations or the State Agricultural Extension Service.


STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep pesticide in original container. Protect from excessive heat. Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling, reconditioning, or puncture and dispose in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



NOTICE - READ CAREFULLY

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States of America

NA 9099 RQ-CAPTAN MIXTURE

ORM-E

