

PH 25 10182-158

1078

RECEIVED  
10-1-1992  
U.S. Department of Agriculture  
National Plant Disease Survey  
10182-158



**Agricultural Products**

# Tillam® 6-E


Selective Herbicide

ACTIVE INGREDIENT:	77.32%
S-propyl butylthiophosphorothioate	
INERT INGREDIENTS:	22.68%
<b>TOTAL</b>	<b>100.00%</b>

Contains 8 pounds active ingredient per gallon.  
EPA Reg. No. 10182-158  
EPA Est. No. 10182-AP-1

Keep Out  
Of Reach  
Of Children  
**CAUTION**

**liquid herbicide for weed control in  
tobacco, tomatoes, and sugar beets**



Contents: 2 1/2 gal (9.46L)

See next panel for  
Statement of Practical  
Treatment and  
Precautionary Statements

## Tillam® 6-E

Selective Herbicide

### STATEMENT OF PRACTICAL TREATMENT

**FIRST AID:** If a known exposure occurs or is suspected, immediately start the recommended procedures below. Simultaneously contact a Poison Control Center, a physician or the nearest hospital. Describe the situation and follow the advice given.

**IF SWALLOWED:** Immediately give several glasses of water but DO NOT induce vomiting. This product contains hydrocarbon solvent. If vomiting occurs, give fluids again. Have a physician determine if condition of patient will permit induction of vomiting or evacuation of stomach. Do not give anything by mouth to an unconscious or comatose person.

**IF ON SKIN:** Flush affected areas with large amounts of running water for at least 15 min. Use and get medical attention. Remove and clean contaminated clothing and shoes.

**IF IN EYES:** Hold eyes apart and flush eyes with large amounts of running water for 15 minutes. Get medical attention if irritation occurs.

**IF INHALED:** Remove to fresh air. Get medical attention if respiratory irritation occurs or if breathing becomes difficult.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-A-I-D (327-6633)  
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
**CAUTION**

**HARMFUL IF SWALLOWED:** Avoid contact with skin, eyes, and clothing. Avoid inhalation of spray mist. Wear rubber gloves and fresh clothing. Wear goggles to avoid contact with eyes. Wash with soap and water immediately after use or contact. Keep away from heat and flame. Do not consume food or feed.

Do not apply this product in such a manner as to drift or through drift, expose workers or other persons. When using the solid jet sprinker method of application, the area being treated must be vacated by unprotected persons.

**ENVIRONMENTAL HAZARDS:**  
Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

CROPS PROTECTED	TARGET WEEDS
Tobacco	Annual Grasses
Tomatoes	Barnyardgrass (Panicum)
Sugar beets	Crabgrass seedling
	Cyperus
	Annual Broadleaf Weeds
	Lambquarters
	Redroot Pigweed
	Ragweed Weeds
	Purple Nutsedge
	Water Hellebore

## Tillam® 6-E

Selective Herbicide

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to attached booklet for use directions.

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, feed, or fertilizer by storage or disposal. Open dumping is prohibited.

**STORAGE:** Keep containers closed when not in use. Do not store near seeds, fertilizers, or foodstuffs. Protect from temperatures below 20°F, but do not store near sources of heat and open flame.

**PESTICIDE DISPOSAL:** Pesticide spray equipment. Wastes resulting from the use of the product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent), then offer for recycling or reconditioning, or purchase and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

### CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of the product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, malperformance, or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims resulting to such factors.

Seller warrants that the product conforms to the chemical description on the label and is responsibly fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be claimed or receive notice of the release or limit below. The EXCLUSIVE REMEDY OF BUYER OR USER AND THE LIMIT OF LIABILITY of Seller will be, at the discretion of Seller, return of the purchase price paid for product bought or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES, THE SUITING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

This product is sold only for use as stated on its label. No express or implied warranty is granted to use it for the product under any circumstances in any country except as specified. Country: United States of America.

Tillam® 6-E is a trademark of an ICI Group Company.  
©1982 ICI Americas Inc.

Made in U.S.A.



**Agricultural Products**

ICI Americas Inc.  
Wilmington, Delaware 19897

0022 G

**BEST AVAILABLE COPY**

# Triallin O-L

## Selective Herbicide

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 0183-158

### MINI-BULK CONTAINERS

liquid herbicide for weed control in tobacco, tomatoes, and sugar beets

<b>ACTIVE INGREDIENT</b>	
Sulfonyl-tolyl-methyl-carbamate	77.32%
<b>INERT INGREDIENTS</b>	22.68%
<b>TOTAL</b>	<b>100.00%</b>

Contains 6 pounds active ingredient per gallon  
EPA Reg. No. 0182-158  
EPA Est. No. 0162-AR1

Keep Out of Reach of Children

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**FIRST AID:** If skin or eye exposure or inhalation is suspected, immediately start first aid. If eye exposure occurs, flush eyes continuously with water for 15 minutes. If eye exposure is severe, contact a Poison Control Center, a physician, or the nearest hospital. If skin exposure occurs, wash with soap and water.

**IF SWALLOWED:** Immediately give water to drink. **DO NOT** induce vomiting. This product is not to be used on food crops. If you are unsure of the amount of product ingested, contact a Poison Control Center, a physician, or the nearest hospital. Do not eat, drink, or smoke until you have been examined by a physician.

**IF ON SKIN:** Flush affected areas with large amounts of water. Wash with soap and water. If irritation occurs, consult a physician. Do not use on clothing or linens.

**IF IN EYES:** Hold eyelids apart and flush eyes with large amounts of water for 15 minutes. If irritation occurs, consult a physician.

**IF INHALED:** Remove to fresh air. Get medical attention if respiratory irritation occurs or if breathing becomes difficult.

**FOR 24 HOUR EMERGENCY MEDICAL ASSISTANCE CALL 1 800 F A S T M E D (327 6533)**  
**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident call CHEMTREC**  
**1 800 424 9300**

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

**HARMFUL IF SWALLOWED:** Avoid contact with skin, eyes, or clothing. Avoid contact with spray mist. Wear rubber gloves and flush clothing. Wear goggles to avoid contact with eyes. Wash with soap and water immediately after use. Keep away from food and feed. Do not eat, drink, or smoke until you have been examined by a physician.

Use only in the manner as directed on the label. Do not use on food crops. Do not use on lawns, lawns, or other areas where children or pets may be present. Do not use on areas where children or pets may be present.

#### ENVIRONMENTAL HAZARDS

This herbicide is toxic to fish, aquatic invertebrates, and birds. Do not use in or near water bodies. Do not use in or near areas where birds, bees, or other beneficial insects may be present.

#### CROPS PROTECTED

Tobacco  
Tomatoes  
Sugar Beets

#### TARGET WEEDS

Annual Grasses  
Annual Broadleaf Weeds  
Perennial Grasses  
Perennial Broadleaf Weeds

Annual Broadleaf Weeds  
Annual Grasses  
Perennial Grasses  
Perennial Broadleaf Weeds

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to attached booklet for use directions.

#### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, feed, or fertilizer by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Rinse spray equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**STORAGE:** Keep containers closed when not in use. Do not store near seeds, fertilizers, or foodstuffs. Protect from temperatures below 20°F, but do not store near sources of heat and open flame.

**CONTAINER DISPOSAL:** Reseal container and offer for reconditioning or recycle use, or equivalent. Offer for recycling or reconditioning or clean in accordance with manufacturer's instructions.

#### FOR MINI BULK CONTAINERS

**Container Precautions:** Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on the screw threads.

**Refill Only WITH TRIALLIN O-L.** The contents of this container are to be refilled only with the same product. Do not mix with other products.

**After filling and before transporting, check for leaks.**

**Do not use a damaged or leaking container.**

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Limited Warranty before using this product.

#### CONDITIONS OF SALE AND LIMITED WARRANTY

This product is sold as a selective herbicide for use on tobacco, tomatoes, and sugar beets. It is not intended for use on other crops. The user assumes all responsibility for the proper use of this product. The manufacturer warrants that the product is free from defects in material and workmanship at the time of sale. The manufacturer does not warrant that the product will perform as described on all soils, climates, or weed populations. The manufacturer does not warrant that the product will be effective against all weeds. The manufacturer does not warrant that the product will be effective against all weeds in all areas. The manufacturer does not warrant that the product will be effective against all weeds in all areas. The manufacturer does not warrant that the product will be effective against all weeds in all areas.

The user assumes all responsibility for the proper use of this product. The manufacturer does not warrant that the product will perform as described on all soils, climates, or weed populations. The manufacturer does not warrant that the product will be effective against all weeds. The manufacturer does not warrant that the product will be effective against all weeds in all areas. The manufacturer does not warrant that the product will be effective against all weeds in all areas.

ICI AMERICA, INC.



### Agricultural Products

ICI Americas Inc  
Wilmington, Delaware 19837

Contents: 110 gal (416 L)

089212

BEST AVAILABLE COPY

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

**CONDITIONS OF SALE AND LIMITED WARRANTY:**  
The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. **SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.**

When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set forth below. The **EXCLUSIVE REMEDY OF BUYER OR USER** and the **LIMIT OF LIABILITY** of Seller will be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. **SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.**

This product is sold only for uses stated on the label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America.

*DEVIRINOL*, *DYFONATE*, and *TILLAM* are trademarks of ICI Group Companies.  
*Compen* is a trademark of Kato, Inc.  
*Dasanit* is a trademark of Farbwerke Bayer, GmbH, Leverkusen, Germany.  
*Micro Cell* is a trademark of Manville Sales Corporation.  
*Mocap* is a trademark of Rhone-Poulenc Ag Company.  
*Paarlan* is a trademark of Dow Chemical Company.  
*Sponto* is a trademark of Witco Corporation.  
*Uni-Mix* is a trademark of Thompson Hayward Chemical Company.  
©1992 ICI Americas Inc.

<b>ACTIVE INGREDIENT:</b>	
S-propyl butylethylthiocarbamate . . . . .	77.32%
<b>INERT INGREDIENTS:</b> . . . . .	22.68%
<b>TOTAL</b> . . . . .	<b>100.00%</b>

Contains 6 pounds active ingredient per gallon.  
EPA Reg. No. 10182-158

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**  
**FIRST AID:** If a known exposure occurs or is suspected, immediately start the recommended procedures below. Simultaneously contact a Poison Control Center, a physician, or the nearest hospital. Describe the situation and follow the advice given.

**IF SWALLOWED:** Immediately give several glasses of water but **DO NOT** induce vomiting. This product contains hydrocarbon solvent. If vomiting occurs, give fluids again. Have a physician determine if condition of patient will permit induction of vomiting or evacuation of stomach. Do not give anything by mouth to an unconscious or convulsing person.

**IF ON SKIN:** Flush affected areas with large amounts of running water for at least 15 minutes and get medical attention. Remove and clean contaminated clothing and shoes.

**IF IN EYES:** Hold eyelids apart and flush eyes with large amounts of running water for 15 minutes. Get medical attention if irritation occurs.

**IF INHALED:** Remove to fresh air. Get medical attention if respiratory irritation occurs or if breathing becomes difficult.


**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E- (327-8633)**  
**FOR CHEMICAL EMERGENCY:** Spill, leak, fire, exposure, or accident Call CHEMTREC 1-800-424-9300

ICI Americas Inc  
Wilmington, Delaware 19897


**Agricultural Products**



Made in U.S.A.



**Tillam 6-E**  
**SELECTIVE HERBICIDE**



liquid herbicide  
for weed control  
in tobacco,  
tomatoes, and  
sugar beets

Complete Directions For Use

**CONTENTS**

Statement of Practical Treatment . . . . .	2
Precautionary Statements . . . . .	3
Weeds Controlled by TILLAM 6-E . . . . .	3
General Use Precautions . . . . .	4
Directions for Use of TILLAM 6-E . . . . .	5
TILLAM 6-E with Fluid Fertilizer . . . . .	7
Impregnation on Dry Fertilizer . . . . .	7
Cultural Practices . . . . .	10
Recommendations . . . . .	10
TILLAM 6-E/Dasanit Spray Concentrate Tankmix Combination . . . . .	14
TILLAM 6-E/Mocap EC Tankmix Combination . . . . .	15
TILLAM 6-E/Paarlan EC Tankmix Combination . . . . .	16
TILLAM 6-E/DEVIRINOL 50-WP, 50-DF, or 2-E Tankmix Combination . . . . .	17
TILLAM 6-E/DYFONATE 4-EC/ DEVIRINOL 50-WP, 50-DF, or 2-E Tankmix Combination . . . . .	21
Appendix I . . . . .	23
Appendix II . . . . .	25
Storage and Disposal . . . . .	29

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

**HARMFUL IF SWALLOWED.** Avoid contact with skin, eyes, and clothing. Avoid inhalation of spray mist. Wear rubber gloves and fresh clothing. Wear goggles to avoid contact with eyes. Wash with soap and water immediately after use or contact. Keep away from heat and flame. Do not contaminate food or feed.

Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. When using the solid set sprinkler method of application, the area being treated must be vacated by unprotected persons.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**WEEDS CONTROLLED**

<b>Annual Grasses:</b>	
Barnyardgrass (Watergrass)	<i>Echinochloa</i> spp
Bermudagrass, seedling	<i>Cynodon dactylon</i>
Crabgrass	<i>Digitaria</i> spp
Foxtails—Yellow, Green	<i>Setaria</i> spp
Goosegrass	<i>Eleusine indica</i>
Millet	<i>Setaria</i> spp
Narcissus	<i>Bracharia piperata</i>
Synthetic grass	
Wild Oats	<i>Avena fatua</i>
<b>Annual Broadleaf Weeds:</b>	
Black-eyed susan (Sweet Suzy)	<i>Rudbeckia serotina</i>

Deadnettle (Henbit)  
Florida pusley  
Hairy Nightshade\*  
Lambsquarters  
Nettleleaf goosefoot  
Prostrate pigweed  
Purslane, Common  
Redroot pigweed  
Shepherd's-purse

*Lamium amplexicaule*  
*Richardia scabra*  
*Solanum villosum*  
*Chenopodium album*  
*Chenopodium murale*  
*Amaranthus graecizans*  
*Portulaca oleracea*  
*Amaranthus retroflexus*  
*Capsella bursa-pastoris*

The above broadleaf weeds will be controlled only if TILLAM 6-E selective herbicide is applied when conditions are favorable for germination.

\*Suppression (partial control)

#### Perennial Weeds:

Yellow nutsedge *Cyperus esculentus*  
Purple nutsedge *Cyperus rotundus*

Existing stands of nutsedge must be turned under and chopped up thoroughly before treatment.

## GENERAL USE PRECAUTIONS

Read all label directions before using.

TILLAM 6-E should be used only for recommended purposes and at recommended rates (do not overdose).

Applied according to directions and under normal growing conditions, TILLAM 6-E will not harm the treated crop. During germination and growth, extended periods of unusually cold and wet or hot and dry weather, insect, nema-

tode, or plant disease attack, the use of certain soil applied systemic insecticides, highly saline or alkaline soil conditions, improperly placed fertilizers or soil insecticides may create abnormal conditions that weaken crop seedlings. TILLAM 6-E used under these abnormal conditions could result in crop injury.

## DIRECTIONS FOR USE OF TILLAM 6-E

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TILLAM 6-E is a selective herbicide which is incorporated into the soil, or applied below the surface of the soil, or applied through solid set sprinklers for control of weeds listed on this label. TILLAM 6-E controls weeds by interfering with normal seed germination and seedling development. It does not control established weeds. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

### SOLID SET SPRINKLERS

**USE PRECAUTIONS:** Apply this product only through solid set sprinklers irrigation system. Do not apply this product through any other type of irrigation system.

- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### USE REQUIREMENTS

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

- Do not apply when wind speed favors drift beyond the area intended for treatment.

## TILLAM 6-E WITH FLUID FERTILIZER

TILLAM 6-E may be combined with fluid (solution slurry, or suspension) fertilizers. However, physical compatibility with these fertilizers must be determined before combining in the spray tank. See Appendix I giving directions for these combinations. Even though found to be compatible, constant agitation is necessary to keep the TILLAM 6-E evenly mixed with the liquid fertilizer.

## IMPREGNATION ON DRY FERTILIZER

TILLAM 6-E may be impregnated on many dry bulk fertilizers for incorporation prior to transplanting of tobacco to control weeds listed on this label. To ensure optimum results, TILLAM 6-E must be uniformly distributed on the fertilizer particles. See Appendix II for special instructions regarding directions for impregnation and use.

## APPLICATION DIRECTIONS

The soil should be well-worked and dry enough to permit good soil mixing (incorporation) or subsurface application. Pour the recommended amount of TILLAM 6-E into the spray tank during the filling operations. Apply 10 to 50 gallons of water per acre.

For soil incorporated applications, use a properly calibrated, low-pressure, boom-type sprayer having good agitation. For subsurface applications, soil should be free of trash. Special equip-

ment designed for subsurface application must be used. Sweeps must be mounted immediately ahead of the planter unit.

In irrigated areas, do not apply TILLAM 6-E before preirrigation.

For all band applications, use proportionately less material per acre depending upon row spacing and band width to be treated. Do not use band application on rocky ground because thorough incorporation is not possible.

## INCORPORATION DIRECTIONS

TILLAM 6-E must be mixed (incorporated) into the soil IMMEDIATELY after application to prevent loss by evaporation of the herbicide. Whenever possible, application and incorporation should be done in the same operation.

**INCORPORATION BEFORE PLANTING:** Use power-driven cultivation equipment set to cut to a depth of 2 to 3 inches in light soils and 3 to 4 inches in heavy soils OR a tandem disc set to cut to a depth of 6 inches, followed by a spike-tooth harrow or some other leveling device which extends beyond ends of the disc. For thorough mixing, disc in two different directions (cross disc).

**INCORPORATION AT OR AFTER PLANTING:** Use power-driven cultivation equipment set to cut to a depth of 2 to 3 inches. OR on light soils only, ground-driven tillers (rolling cultivators, rotary hoe, row wheels, etc.) set to cut to a depth of 1½ inches and operated at 6 to 8 mph. When using ground-driven tillers preincubate if soil is crusted. When incorporating after planting care must be taken not to disturb the crop seedling.

## SUBSURFACE DIRECTIONS

(Tobacco only): Special equipment designed for subsurface sweep application MUST be used. Sweeps must be mounted immediately ahead of the planter unit.

**Covered Sweeps:** Set the sweeps to run below the soil surface deep enough to cover the TILLAM 6-E with 5 to 6 inches of soil. Calibrate by measuring the spray band width at the back of the sweep, not the sweep width. NOTE: When applying with sweeps, TILLAM 6-E must be applied deep enough to allow 3 to 4 inches of soil to remain over the treatment after the transplanting operation.

## BED-OVER METHOD OF APPLICATION

Apply TILLAM 6-E in a band immediately in front of bedding discs. Set discs so that TILLAM 6-E is covered with 6 inches of soil. Set bedding discs so that treated soil is not folded in. If bed leveling are used at transplanting, beds should be made higher to compensate for leveling, i.e., so that 3 to 4 inches of soil are left above the TILLAM 6-E treatment.

**SOLID SET SPRINKLER APPLICATION (Tomatoes only, Western Region):** Meter the TILLAM 6-E into the irrigation water during the entire period. Apply in 1.5 to 3 acre inch of water or sufficient water to penetrate the soil to a depth of 2 to 4 inches.

Time the TILLAM 6-E application to ensure that proper penetration of the herbicide corresponds with the end of the irrigation period. Flush the lines, then turn the water off promptly. Tap water (run-off water) from irrigation should be recirculated.

**BEST AVAILABLE COPY**

Application of more than recommended quantities of irrigation water may result in decreased product performance by removing the chemical from the zone of effectiveness.

## PLANTING

For maximum weed control benefits, crops should be planted or seeded immediately after application. Do not use a drag behind the planter as it may concentrate TILLAM 6-E over the seed row and cause crop injury.

## CULTURAL PRACTICES FOLLOWING APPLICATION

TILLAM 6-E is not persistent in the soil and susceptible weeds germinating later in the season may not be controlled. Shallow cultivation or approved postemergence herbicides may be necessary to control susceptible weeds that escape control by TILLAM 6-E as well as those weeds not susceptible to TILLAM 6-E. Do not cultivate deeper than the depth of herbicide incorporation.

When cultivating fields where TILLAM 6-E has been banded in the row, use shields to prevent the movement of untreated soil into the treated row.

## RECOMMENDATIONS

**SUR BEETS, PREPLANT:** Apply just prior to planting, see DIRECTIONS FOR USE for methods of application and incorporation. Recommended dosages are as follows:

For light textured soils and soils with less than 10% organic matter, apply 2½ qts. TILLAM 6-E per acre.

For heavy textured soils and soils with over 10% organic matter, apply 4 qts. TILLAM 6-E per acre.

### TOBACCO, FLUE-CURED

**Before bedding and transplanting:** Apply TILLAM 6-E at a broadcast rate of 2½ qts. per acre. Incorporate immediately with a tandem disc and bed up immediately prior to transplanting.

OR

**After bedding but before transplanting:** Apply TILLAM 6-E at a broadcast rate of 2½ qts. per acre prior to transplanting, reducing amount per acre proportionately depending upon band width and row spacing. Incorporate immediately using a power-driven rotary hoe with variable length blades (i.e., one with short blades on the inside and longer blades on the outside). Bed levelers, if used, should be kept shallow as excessive removal of treated soil will destroy weed control.

OR

**Subsurface sweeps after bedding but before transplanting:** Run at a depth so that when bed is in final shape, 3 to 4 inches of soil will cover the TILLAM 6-E. Apply TILLAM 6-E at 2 qts. per acre, just prior to transplanting. For band treatment, reduce amount per acre proportionately depending upon band width and row spacing.

### TOBACCO, LIGHT AIR CURED

Includes Type 31 (Bulley) & Type 32 Tobaccos, Type 41 Cigar Filler, Types 54 and 55 Cigar Binder.

**Before transplanting:** Apply TILLAM 6-E broadcast at 2½ qts. per acre. Incorporate immediately, only with a tandem disc on flat ground followed by transplanting. For band treatment reduce amount per acre proportionately depending upon band width and row spacing. Incorporate immediately, using a power driven rotary hoe.

## TOMATOES

SEE DIRECTIONS FOR USE FOR METHODS OF APPLICATION AND INCORPORATION

### DIRECT SEEDED (Western Region Only):

Apply TILLAM 6-E just prior to planting.

For light textured soils and soils with less than 10% organic matter, apply 2½ qts. TILLAM 6-E per acre.

For heavy textured soils and soils with more than 10% organic matter, apply 4 qts. TILLAM 6-E per acre.

OR

### DIPECT SEEDED, LAYBY (Not preplant):

Apply TILLAM 6-E as a directed spray to clean cultivated soil. Do not apply within eight (8) days of harvest.

For light textured soils and soils with less than 10% organic matter, apply 2½ qts. TILLAM 6-E per acre.

For heavy textured soils and soils with more than 10% organic matter, apply 4 qts. TILLAM 6-E per acre.

OR

### DIRECT SEEDED, SPRINKLER APPLICATION

(Western Region only): Apply TILLAM 6-E through solid set sprinklers prior to or immediately after planting, before weeds emerge.

For light textured soils and soils with less than 10% organic matter, apply 2½ qts. TILLAM 6-E per acre.

For heavy textured soils and soils with more than 10% organic matter, apply 4 qts. TILLAM 6-E per acre.

OR

### PLUG PLANTED, PREPLANT (Western Region

only): Prior to plug planting tomatoes, apply and incorporate up to 6½ quarts of TILLAM 6-E per acre. Use higher rate for hairy nightshade.

### TOMATO TRANSPLANTS (DO NOT USE HOTCAPS)

**Before transplanting:** Apply TILLAM 6-E prior to transplanting.

OR

**After transplanting:** Apply TILLAM 6-E as a directed spray to clean, cultivated soil. Do not apply within eight (8) days of harvest.

For light textured soils and soils with less than 10% organic matter, apply 2½ qts. TILLAM 6-E per acre.

For heavy textured soils and soils with more than 10% organic matter, apply 4 qts. TILLAM 6-E per acre.

### TOMATO TRANSPLANTS (South Carolina only):

**Before bedding and transplanting:** Apply TILLAM 6-E at a broadcast rate of 2 qts. per acre just prior to bedding and transplanting, reducing the amount per acre proportionately depending upon band width and row spacing. Use bed-over method of application.

OR

**After bedding but before transplanting:** Apply TILLAM 6-E at a broadcast rate of 2 qts. per acre just prior to transplanting, reducing amount per acre proportionately depending upon band width and row spacing. Apply to the top of the bed and incorporate 4 to 6 inches into the bed top.

OR

**Postemergence or after transplanting:** Apply TILLAM 6-E at a broadcast rate of 2 qts. per acre, reducing amount per acre proportionately depending upon band width and row spacing. Apply in a directed spray to soil around base of tomato plants using bands on either side of the row. Immediately after treatment, sweep out row centers to effectively cover TILLAM 6-E treat-

ment with a layer of soil 2 inches deep. Do not apply within eight (8) days of harvest.

## TILLAM 6-E/DASANIT® SPRAY CONCENTRATE TANKMIX COMBINATION

TILLAM 6-E IN COMBINATION WITH DASANIT SPRAY CONCENTRATE controls nematodes, wireworms and weeds controlled by TILLAM 6-E in tobacco (Flue Cured and Light Air Cured).

**NOTE:** Dasanit is a Restricted Use Pesticide. Refer to the label for Dasanit Spray Concentrate for pertinent directions and precautions for use, and restrictions.

## APPLICATION DIRECTIONS AND RECOMMENDATIONS

The soil should be well-worked and dry enough to permit good soil incorporation (mixing into the soil). Apply 2½ qts. TILLAM 6-E and 4 qts. Dasanit Spray Concentrate in 20 to 50 gallons water per acre. Use a properly calibrated low-pressure boom-type sprayer having good agitation.

## SOIL INCORPORATION

The tankmix must be incorporated (mixed) into the soil immediately after application to prevent loss of the herbicide by evaporation. Whenever possible, application and incorporation should be done in the same operation.

Use power-driven cultivation equipment set to cut to a depth of 3 inches in light soil and 3 to 4 inches in heavy soil or a tandem disc set to cut to a depth of 4 to 6 inches.

Cross disc at right angles for thorough mixing. The bedding operation should follow the incorporation as soon as is practical. Planting may be done immediately after bedding or at a later date, however, to ensure long term weed control, the crop should be planted as soon as possible after application and bedding.

## TILLAM 6-E/MOCAP® EC NEMATICIDE-INSECTICIDE TANKMIX COMBINATION

A tankmix of TILLAM 6-E and Mocap EC can be applied and mixed into the soil before transplanting for control of weeds, wireworms, and nematodes in tobacco.

**NOTE:** Mocap is a Restricted Use Pesticide. Refer to the label of Mocap EC for pertinent directions and precautions for use, and restrictions.

## TANK MIXING DIRECTIONS

The TILLAM 6-E/Mocap EC tankmix combination can be mixed with water and applied as a spray. Pour the recommended amount of TILLAM 6-E directly into the nearly filled spray tank while agitating. Next, add the recommended amount of Mocap EC. Fill the remaining portion of the tank with water, mix thoroughly. Maintain a moderate agitation to provide a uniform mixture upon application.

## APPLICATION DIRECTIONS AND RECOMMENDATIONS

The soil should be well-worked and dry enough to permit good soil mixing (incorporation). Apply 2½ qts. TILLAM 6-E plus 1 gallon Mocap EC in

BEST AVAILABLE COPY

10 to 50 gallons water per acre. Use a properly calibrated low-pressure, boom-type sprayer having good agitation

### INCORPORATION DIRECTIONS

The TILLAM 6-E/Mocap EC tankmix combination must be mixed (incorporated) into the soil immediately after application to prevent loss by evaporation. Whenever possible, application and incorporation should be done in the same operation. Use a tandem disc set to cut to a depth of 6 inches, followed by a spike-tooth harrow or some other leveling device which extends beyond the ends of the disc. For thorough mixing, disc in two different directions (cross disc).

### TILLAM 6-E/PAARLAN® EC HERBICIDE TANKMIX COMBINATION

A tankmix of TILLAM 6-E and Paartan EC can be applied and mixed into the soil before transplanting for additional weed control and longer season grass control in tobacco

NOTE: Refer to the label of Paartan EC for pertinent directions and precautions for use, and restrictions

### TANK MIXING DIRECTIONS

The TILLAM 6-E/Paarlan EC tankmix combination can be mixed with water and applied as a spray. Pour the recommended amount of TILLAM 6-E directly into the nearly filled spray tank while agitating. Next, add the recommended amount of Paartan EC. Fill the remaining portion of the tank with water, mix thoroughly. Maintain a moderate agitation to provide a uniform mix during application

### APPLICATION DIRECTIONS AND RECOMMENDATIONS

The soil should be well-worked and dry enough to permit good soil mixing (incorporation). Apply 2½ qts. TILLAM 6-E plus 2 pints Paartan EC in 10 to 50 gallons water per acre. Use a properly calibrated low-pressure, boom-type sprayer having good agitation

### INCORPORATION DIRECTIONS

The TILLAM 6-E/Paarlan EC tankmix combination must be mixed (incorporated) into the soil immediately after application to prevent loss by evaporation. Whenever possible, application and incorporation should be done in the same operation. Use a tandem disc set to cut to a depth of 6 inches, followed by a spike-tooth harrow or some other leveling device which extends beyond the ends of the disc. For thorough mixing, disc in two different directions (cross disc)

### TILLAM 6-E/DEVIRINOL® 50-WP, 50-DF, or 2-E SELECTIVE HERBICIDE TANKMIX COMBINATION APPLIED PREPLANT INCORPORATED

A tankmix combination of TILLAM 6-E and DEVIRINOL 50-WP, 50-DF, or 2-E can be applied and mixed (incorporated) into the soil before planting for weed control in tomatoes and tobacco

### WEEDS CONTROLLED

#### Annual Grasses:

Barnyardgrass *Echinochloa crus-galli*  
(Watergrass)

Bermudagrass, seedling  
Crabgrass  
Foxtails (Giant, Green, Yellow)  
Goosegrass  
Millet  
Narrow Leaf Signalgrass  
Wild Oats

#### Annual Broadleaf Weeds:

Blackeyed-susan *Rudbeckia serotina*  
(Sweet Suzy)  
Deadnettle (Henbit) *Lamium amplexicaule*  
Florida pusley *Richardia scabra*  
Hairy Nightshade *Solanum villosum*  
Lambsquarters *Chenopodium album*  
Nettleleaf Goosefoot *Chenopodium murale*  
Prostrate Pigweed *Amaranthus graecizans*  
Purslane, Common *Portulaca oleracea*  
Redroot Pigweed *Amaranthus retroflexus*  
Shepherd's-purse *Capsella bursa-pastoris*

\*Suppression (partial control)

The above broadleaf weeds will be controlled only when the TILLAM 6-E/DEVIRINOL 50-WP, 50-DF, or 2-E tankmix is applied when conditions are favorable for germination.

#### Perennial Weeds:

Purple Nutsedge *Cyperus rotundus*  
Yellow Nutsedge *Cyperus esculentus*

Existing stand of nutsedge must be turned under and chopped up thoroughly before treatment

Note: Observe all cautions and limitations on labeling of all products used in combination

Make only one application per season

In each case where a range of rates is given, the lower rates should be used on light (coarse) textured soils and soils low in organic matter. The higher rates should be used on heavy (fine) textured soils

After harvest or prior to planting of succeeding crops, a deep moldboard or disc plowing operation must be carried out

### TANK MIXING DIRECTIONS

The TILLAM 6-E/DEVIRINOL 50-WP, 50-DF, or 2-E tankmix combination can be mixed with water and applied as a spray. Pour the recommended amounts of TILLAM 6-E and DEVIRINOL 50-WP, 50-DF, or 2-E into a spray tank during the filling operations. When using DEVIRINOL 50-WP or 50-DF, add first to the spray tank, then add the TILLAM 6-E. Maintain adequate agitation of the tankmix. Apply in 25 to 50 gallons of water per acre

### APPLICATION DIRECTIONS AND RECOMMENDATIONS

The soil should be well-worked and dry enough to permit good soil mixing (incorporation)

For soil incorporated applications, use a properly calibrated low-pressure, boom-type sprayer having good agitation

In irrigated areas do not apply before preirrigation. For all band applications, reduce dosage proportionately depending upon row spacing and band to be treated. Do not use band application on rocky ground because thorough incorporation is not possible

*Cynodon dactylon*

*Digitaria* spp.  
*Setaria* spp.

*Eleusine indica*  
*Setaria* spp.  
*Brachiana pilgera*

*Avena fatua*

### INCORPORATION DIRECTIONS

The herbicides must be mixed (incorporated) into the soil immediately after application to prevent loss by evaporation. Whenever possible, application and incorporation should be done in the same operation

### INCORPORATION BEFORE PLANTING

Use power-driven cultivation equipment set to cut to a depth of 2 to 3 inches in coarse soils and 3 to 4 inches in heavy soils OR a tandem disc set to cut to a depth of 6 inches, followed by a spike-tooth harrow or some other leveling device which extends beyond ends of the disc. For thorough mixing, disc in two different directions (cross disc)

### INCORPORATION AT PLANTING

Use power-driven cultivation equipment set to cut to a depth of 2 to 3 inches OR, on coarse soils only, ground-driven tillers (rolling cultivators, rotary hoe, row wheels, etc.) set to cut to a depth of 1½ inches and operated at 6 to 8 mph. When using ground-driven tillers, pre-cultivate if soil is crusted.

### PLANTING

Crops should be planted or seeded immediately after application and incorporation. Do not use a drag behind the planter as it may concentrate the herbicides over the seed row and cause crop injury

### CULTURAL PRACTICES FOLLOWING APPLICATION

Shallow cultivation may be necessary to control

those susceptible weeds that escape control as well as those weeds not susceptible to either herbicide. Do not cultivate deeper than the depth of herbicide incorporation

When cultivating fields where the tankmix has been banded in the row, use shields to prevent the movement of untreated soil into the treated row

### RECOMMENDATIONS

**TOMATOES (Direct Seeded [California Only] and Transplanted):** Apply 2 to 4 pounds DEVIRINOL 50-WP or 50-DF, or 2 to 4 qts DEVIRINOL 2-E plus 2½ to 4 qts TILLAM 6-E per broadcast acre. Use the lower rates on light soils (coarse textured-sandy) and the higher rate on heavy soils (fine textured-clay). Do not use hotcaps on transplanted tomatoes

**TOBACCO (Transplanted):** Before transplanting apply 2 pounds DEVIRINOL 50-WP or 50-DF or 2 qts DEVIRINOL 2-E plus 2½ qts TILLAM 6-E per broadcast acre

To avoid injury to crops not specified on the label, do not plant until 12 months after the last DEVIRINOL application

### TILLAM 6-E/DYSCONATE® 4-EC EMULSIFIABLE LIQUID INSECTICIDE/DEVIRINOL 50-WP, 50-DF, or 2-E TANKMIX COMBINATION

A tankmix combination of TILLAM 6-E, DYSCONATE 4-EC and DEVIRINOL 50-WP, 50-DF or 2-E can be applied and mixed into the soil before transplanting tomatoes and tobacco for control of insects and soil insects. Refer to TILLAM 6-E DEVIRINOL Tankmix Combination for list of weeds controlled

NOT A COPY

**NOTE: DYFONATE 4-EC is a Restricted Use Residue.** Observe all cautions and limitations on labeling of all products used in combinations. Make only one application per season. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

The recommended tankmix controls weeds by interfering with normal seed germination and seedling development. It does not control established weeds. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

For GENERAL USE PRECAUTIONS and DIRECTIONS FOR USE refer to TILLAM 6-E/DEVIRINOL Tankmix Recommendations for methods of tank mixing, application and incorporation.

**RECOMMENDATIONS**

**TOMATOES (Direct Seeded [California Only] and Transplanted):** Apply 2 1/2 to 4 qts. TILLAM 6-E plus 2 to 4 pounds DEVIRINOL 50-WP or 50-DF, or 2 to 4 qts. DEVIRINOL 2-E plus 2 qts. DYFONATE 4-EC for control of symphylans or 4 qts. for control of symphylans and wireworms in 25 to 50 gallons of water per broadcast acre. Use the lower DEVIRINOL and TILLAM herbicide rates on light soils (coarse texture-sandy) and the higher rates on heavy soils (fine textured-clay). Do not use hotcaps on transplanted tomatoes.

**ACCORNS (Transplanted):** Before transplanting, apply 2 1/2 qts. TILLAM 6-E plus 2 pounds DEVIRINOL 50-WP or 50-DF, or 2 qts. DEVIRINOL 2-E plus 2 qts. DYFONATE 4-EC for control of wireworms in 25 to 50 gallons of water per broadcast acre.

To avoid injury to crops not specified on the label, do not plant until 12 months after the last DEVIRINOL application.

**APPENDIX I**

**TILLAM 6-E WITH FLUID FERTILIZERS**

The following procedure is suggested for determining whether TILLAM 6-E may be combined with a specific fluid fertilizer for spray tank application.

**MATERIALS REQUIRED:**

- 1. TILLAM 6-E
- 2. Fluid fertilizer to be used
- 3. Adjuvant for fertilizer tankmix: Complex™, Sponto™ 168-D, Uni-Mix™, or equivalent. The adjuvant which provides the best emulsification depends on the specific fertilizer under consideration.
- 4. Two one quart, wide mouth glass jars with lid or stopper.
- 5. Measuring spoons (a 25 mL pipette or graduated cylinder provides more accurate measurement).
- 6. Measuring cup, 8 oz (237 mL).

**PROCEDURE:**

1. Pour a pint (about 473 mL) of the fluid fertilizer into each of the quart jars.
2. Add adjuvant to one of the jars and mix (see rate table).

3. Add the TILLAM 6-E to both jars (see rate table).
4. Close both jars with lid or stopper and mix the contents by turning the jars upside down ten times.
5. Inspect the surface and body of the mixtures--
  - (A) Immediately after completing the jar inversions.
  - (B) After allowing the jars to stand quietly for 30 minutes.
  - (C) And then again after turning the jars upside down 10 times.

If a uniform mix cannot be made, the mixture should not be used. If either mixture remains uniform for 30 minutes, the combination may be used. Should either mixture separate after 30 minutes, but readily remix uniformly with the jar inversions, the mixture can be used if adequate agitation is maintained in the tank. If the mixture with adjuvant is satisfactory, but the one without adjuvant is not, be sure to use the adjuvant in the spray tank. Add the adjuvant first at a rate of 3 pints per 100 gallons of fluid fertilizer; foaming can be minimized by using moderate agitation.

If nondispersible oil, sludge, or clumps of solids form in the mixtures, the combinations should not be used.

**RATE TABLE FOR TILLAM 6-E AND ADJUVANT\*\* WITH THE FLUID FERTILIZER**

Gallons of fluid fertilizer to be applied per acre	mL or Tsp. of TILLAM 6-E* to be added to 1 pint of fertilizer	
	6-E	
	mL	Tsp.
10	8	1 2/3
15	6	1
20	4	3/4
25	3	2/3
30	2 1/2	2
40	2	3/4

Based on field rate of 1 pound active ingredient per acre in the fertilizer volumes indicated. Increase volume proportionately to correspond with intended field rate in terms of pounds active ingredient per acre (e.g. for field rate of 4 pounds actual TILLAM 6-E in 40 gallons fertilizer per acre, add 8 mL or 2 tsp. TILLAM 6-E to each jar for compatibility testing purposes).

Two (2) mL or one half (1/2) teaspoon of adjuvant to be added to 1 pint of fluid fertilizer in order to equal the rate of 3 pints of adjuvant per 100 gallons of fluid fertilizer.

**APPENDIX II**

**TILLAM 6-E IMPREGNATION ON DRY BULK FERTILIZERS FOR USE ON TOBACCO**

**CAUTION:** TILLAM 6-E must not be impregnated on ammonium nitrate, sodium nitrate, or potassium nitrate. Such mixtures may cause explosion and fire.

TILLAM 6-E may be impregnated on many dry bulk fertilizers\* and applied and incorporated in the soil before planting for the control of grasses and broadleaf weeds in tobacco.

Field results have shown that TILLAM 6-E on bulk dry fertilizers gives weed control equal to TILLAM 6-E applied as a spray in water. However, uniform impregnation of TILLAM 6-E on the dry fertilizer particles and uniform application in the field are necessary to assure good results.

For impregnating TILLAM 6-E on dry fertilizer, use a closed rotary-drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle (or nozzles) should be positioned inside of the mixer to provide uniform spray coverage of the tumbling fertilizer. The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with TILLAM 6-E provides a satisfactory, dry mixture. If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry, flowable mixture. Micro-Cel™ E is the recommended absorbent powder. It should be added separately and uniformly to the prepared TILLAM 6-E fertilizer mixture, in a quantity that is sufficient to provide a suitably flowable mixture. Generally less than 2% by weight of Micro-Cel E is required. The amount of TILLAM 6-E actually required in the manufacture of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of TILLAM 6-E actually contained in the mixture applied to the soil represents the correct rate of use.

ALL TILLAM 6-E supplementary literature instructions and label recommendations regarding rates per acre, soil incorporation, application, cautions, general use precautions, and other directions must be followed.

All individual state regulations relating to bulk dry fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the fertilizer and TILLAM 6-E mixture.

A minimum of 200 pounds and a maximum of 2,000 pounds of approved ingredients\* impregnated with TILLAM 6-E at the recommended rate must be applied per acre.

Bulk fertilizer impregnated with TILLAM 6-E should be applied immediately, not stored. It is recommended that all bulk containers be tightly covered while the product is being transported and applied to reduce chances of TILLAM 6-E loss via volatilization.

**\*APPROVED DRY FERTILIZER INGREDIENTS**

	N	P	K
Ammonium sulfate	21	0	0
Diammonium phosphate	18	45	0
Potassium chloride	0	0	60
Potassium sulfate	0	0	52
Super-phosphate (single)	0	20	0
Super-phosphate (triple)	0	45	0
Urea	45	0	0
Ammonium phosphate sulfate (1:48-0)	16	20	0
1:48-0	11	48	0

**NOTE:** K Mag and two MAP fertilizers (12-5-50 0 11-54-0) have been shown to be compatible with TILLAM 6-E and are approved for use.



8048

**RATE CHART FOR IMPREGNATION OF DRY  
BULK FERTILIZERS WITH TILLAM 6-E**

Fertilizer Rate Per Acre	Tillam 6-E Rate Per Acre (2½ qts./acre)
200	26½ qts./ton
300	17½ qts./ton
400	13½ qts./ton
500	10½ qts./ton
600	8½ qts./ton
800	6½ qts./ton
1000	5½ qts./ton
1500	3½ qts./ton
2000	2½ qts./ton

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, feed, or fertilizer by storage or disposal. Open dumping is prohibited.

**STORAGE:** Keep containers closed when not in use. Do not store near seeds, fertilizer, or food-stuffs. Protect from temperatures below 20°F but do not store near sources of heat or open flame.

**COMBUSTIBLE LIQUID**

**PESTICIDE DISPOSAL:** Rinse spray equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:**

**METAL**—Triple rinse (or equivalent), then offer for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

**PLASTIC**—Triple rinse (or equivalent), then offer for recycling, or reconditioning or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**FOR BULK AND MINI-BULK:**

**Container Disposal:** Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

**Container Precautions:** Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

**REFILL ONLY WITH TILLAM 6-E.** The contents of this container cannot be completely removed by cleaning. Refilling with materials other than TILLAM 6-E will result in contamination and may weaken container.

After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**