10182-178

United States Environmental Protection Agency Washington, D.C. 20460

> Office of Prevention, Pesticides and Toxic Substances

007 2 1993

Wayne R. Hillebrecht ZENECA INC ZENECA AG PRODUCTS NEW MURPHY ROAD & CONCORD PIKE BOX 751 WILMINGTON DE 19897

pm 25

Subject: Label Amendment Submission of 06/10/93 in Compliance with WPS Labeling Requirements EPA Reg No. 10182-178 RO-NEET 6-E

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted subject to the comments listed below.

Based on your certification, the Agency has accepted only those changes to your labeling which are necessary to comply with PR Notices 93-7 and 93-11, which reflect the WPS labeling requirements of 40 CFR part 156, subpart K. Any other labeling changes submitted in connection with this amendment application and not directly related to compliance with PR Notice 93-7 or 93-11 have neither been reviewed nor accepted by the Agency. If you wish to make any such changes, you must submit a separate amendment application proposing them. If your product registration is currently suspended, acceptance of this labeling amendment does not affect the suspension in any way.

A copy of your proposed labeling stamped "Accepted with comments" is attached. Make any required changes described in the attached and send three copies of final labeling as soon as it is available to: Document Processing Desk (FIN-LABEL) Office of Pesticide Programs (H-7504C) U.S. Environmental Protection Agency 401 M Street SW Washington, DC 20460-0001

Hand or courier deliveries of final labels may be made to:

Document Processing Desk (FIN-LABEL) Room 266A Crystal Mall 2 1921 Jefferson Davis Highway Arlington, VA 22202

Please correct the typographical errors circled on the draft before printing final labeling.

In your final labeling the "Agricultural Use Requirements" text" must be contained in a clearly separate box. This box may be set apart by a line, by another graphical device, by a different color background, or in any other way that clearly distinguishes it from surrounding text.

hcerely,

James Tompkins, Deputy Chief Registration Support Branch (Registration Division (H7505W)

Page 1 of 7 7 24 RO-NEET 6-E Container RON178C.RSB - RS-052693B

, RO-NEET® 6-E Selective Herbicide

Liquid Herbicide for Weed Control in Sugar Beets, Table Beets, and Spinach

ACTIVE INGREDIENT:

S-ethyl cyclohexylethylthiocarbamate	 73.9%
INERT INGREDIENTS:	 <u>26.1%</u>
Total	 100.0%

Contains 6 pounds active ingredient per gallon.

EPA Reg. No. 10182-178 EPA Est. No. 10182-

•

Net Contents:

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACCEPTED vidi COMMENTS in EPA Letter Duted OCT | 2 1993

Made in U.S.A. ZENECA Ag Products ZENECA Inc. Wilmington, DE 19897

Page 2 RO-NEET 6-E Container RON178C.RSB - RS-052693B

STATEMENT OF PRACTICAL TREATMENT

FIRST AID

If a known exposure occurs or is suspected, immediately start the recommended procedures below. Simultaneously contact a Poison Control Center, a physician, or the nearest hospital. Describe the situation and follow the advice given.

IF SWALLOWED: Immediately give large quantities of water but DO NOT induce vomiting. *This product contains hydrocarbon solvent.* If vomiting does occur, give fluids again. Have a physician determine if condition of patient will permit evacuation of stomach. DO NOT give anything by mouth to an unconscious person. Call a physician immediately.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for 15 minutes. Get medical attention if irritation occurs.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

Page 3 RO-NEET 6-E Container RON178C.RSB - RS-052693B

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

HARMFUL IF SWALLOWED. Avoid contact with skin, eyes, and clothing. Wear-rubber gloves and protective clothing. Wash hands, arms, and face thoroughly with scap and water after handling and before cating or smoking. Routinely shower or bathe after work and wash all clothing with scap and hot water before reusing. In case of contact, immediately remove contaminated clothing. Avoid contamination of feed or food.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear.

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or nitrile rubber or nedprene rubber or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems rendosed cabs, or elicital in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR (170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing?
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

READ ENTIRE LABEL BEFORE USING

Protect from temperatures below 20°F.

Product crystallizes at lower temperatures.

Page 4 RO-NEET 6-E Container RON178C.RSB - RS-052693B

29

Warm or store at higher temperatures and mix to redissolve crystals and assure uniformity before use.

14

Page 5 RO-NEET 6-E Container RON178C.RSB - RS-052693B

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product marway that will contact workers or other persons wither directly or through Mint Monly protected handlers in vibe in the area during application affor any requirements specific to your State of Under consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

<u>....</u>

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling (i.e., attached bookdet) under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

Refer to attached booklet for use directions.

BEST AVAILABLE COPY

RO-NEET 6-E Container RON178C.RSB - RS-052693B

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Keep containers closed when not in use. Do not store near seeds, fertilizers or foodstuffs. Protect from temperatures below 20°F, but do not store near sources of heat or open flame.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

FOR BULK AND MINI-BULK CONTAINERS

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges dents, abrasions, and damaged or worn threads on closure devices.

REFILL ONLY WITH RO-NEET 6-E. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than RO-NEET 6-E will result in contamination and may weaken container.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Page 7 RO-NEET 6-E Container RON178C.RSB - RS-052693B

INPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIABILITY of Seller will be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States of America.

RO-NEET® is a trademark of a ZENECA Group Company.

Page 1 of 20 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

RO-NEET® 6-E Selective Herbicide

Liquid Herbicide for Weed Control in Sugar Beets, Table Beets, and Spinach

COMPLETE DIRECTIONS FOR USE

EPA Reg. No. 10182-178

. .

Made in U.S.A. ZENECA Ag Products ZENECA Inc. Wilmington, DE 19897

Page 2 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACTIVE INGREDIENT:

S-ethyl cyclohexylethylthiocarbamate	· • • •	 	 73.9%
INERT INGREDIENTS:		 	 <u>26.1%</u>
Total .		 · · · · · · · · · · ·	 100.0%

Contains 6 pounds active ingredient per gallon.

14

STATEMENT OF PRACTICAL TREATMENT

FIRST AID

If a known exposure occurs or is suspected, immediately start the recommended procedures below. Simultaneously contact a Poison Control Center, a physician, or the nearest hospital. Describe the situation and follow the advice given.

IF SWALLOWED: Immediately give large quantities of water but DO NOT induce vomiting. *This product contains hydrocarbon solvent.* If vomiting does occur, give fluids again. Have a physician determine if condition of patient will permit evacuation of stomach. DO NOT give anything by mouth to an unconscious person. Call a physician immediately.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for 15 minutes. Get medical attention if irritation occurs.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

Page 3 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

HARMFUL IF SWALLOWED. Avoid contact with skin, eyes, and clothing. Wear rubber gloves and protective clothing. Wash hands, arms, and face thoroughly with scap and water after handling-and before-cating or smoking. Routinely shower or bathe after work and wash all clothing with scap and hot water before reusing. In case of contact, immediately remove contaminated clothing. Avoid contamination of feed or food.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear.

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS:

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove dothing immediately if pesticide gets inside. Then wash thoroughly and put on clean dothing?
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

READ ENTIRE LABEL BEFORE USING

Protect from temperatures below 20°F.



Product crystallizes at lower temperatures.

Page 4 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D 29

137

Warm or store at higher temperatures and mix to redissolve crystals and assure uniformity before use.

Page 5 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, dect ntamination, notification, and emergency assistance. It also contains specific Instructions and exceptions pertaining to the statements on this label about personal protection equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soll-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralis
- Chemical-resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or viton
- Shoes plus socks

GENERAL INFORMATION

RO-NEET® 6-E is a selective herbicide which is mixed (incorporated) into the soil or applied below the surface of the soil for control of weeds listed on this label. RO-NEET 6-E controls weeds by interfering with normal seed germination and seedling development. It does not control established weeds. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

GENERAL USE PRECAUTIONS

BEST AVAILABLE COPY

Not for use or storage in or around the home.

RO-NEET 6-E should be used only for recommended purposes and at recommended rates (DO NOT OVERDOSE). Applied according to directions and under normal growing

Page 6 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

15 9 29

conditions, RO-NEET 6-E will not harm the treated crop. During germination and early growth, extended periods of unusually cold and wet or hot and dry weather, insect, nematode, or plant disease attack, carryover pesticide residues, the use of certain soil applied systemic insecticides, highly saline or alkaline soil conditions, improperly placed fertilizers or soil insecticides may create abnormal conditions that weaken crop seedlings. RO-NEET 6-E used under these abnormal conditions could result in crop injury.

RO-NEET 6-E may cause crop injury on very light sandy soil.

RO-NEET 6-E may be combined with fluid (solution, slurry, or suspension) fertilizers. However, physical compatibility with these fluid fertilizers must be determined before combining in the spray tank. See special literature giving directions for these combinations. Even though found to be compatible, constant agitation is necessary to keep the RO-NEET 6-E evenly mixed with the fluid fertilizer.

Tank mix this product with fungicides, insecticides, or herbicides only as recommended.

Do not contaminate irrigation water or water used for domestic purposes.

Page 7 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D 6929

WEEDS CONTROLLED

ANNUAL GRASSES:

Annual Bluegrass Annual Ryegrass Barley, Volunteer Barnyardgrass (Watergrass) Crabgrass Foxtails Wild Oats

ANNUAL BROADLEAF WEEDS:

Black Nightshade Hairy Nightshade Henbit, Deadnettle Lambsquarters Nettleleaf Goosefoot* Common Purslane Redroot Pigweed Shepherd's-purse Small Stinging Nettle, Burning Nettle Velvetleaf** Poa annua Lolium multiflorum Hordeum spp. Echinochloa spp. Digitaria spp. Setaria spp. Avena fatua

Solanum nigrum Solanum villosum Lamium spp. Chenopodium album Chenopodium murale Portulaca oleracea Amaranthus retroflexus Capsella bursa-pastoris Urtica urens Abutilon theophrasti

*California excluded. **Partial control or suppression only.

The above broadleaf weeds will be controlled only if the application is made when conditions are favorable for germination. RO-NEET 6-E SELECTIVE HERBICIDE WILL NOT CONTROL ESTABLISHED WEEDS.

PERENNIAL WEEDS:

Yellow Nutsedge Purple Nutsedge Cyperus esculentus Cyperus rotundus

Existing stands of nutsedge must be turned under and chopped thoroughly before treatment. Prior to planting, RO-NEET 6-E must be thoroughly mixed into the scal or applied subsurface to the soil with injectors for nutsedge control.

Page 8 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

SOIL PREPARATION: Prepare the soil for seeding according to good agricultural practice. All weed growth should be thoroughly worked into the soil before treatment. DO NOT APPLY BEFORE PREIRRIGATION.

APPLICATION: Apply RO-NEET 6-E to well-worked soil which is dry enough to permit thorough mixing with incorporation equipment or proper injection. Use standard low-pressure (20-50 psi) boom spray equipment. DO NOT USE SINGLE NOZZLE BOOM-JET TYPE SPRAYERS. Application equipment should be carefully calibrated before use and checked frequently during application to be sure nozzles are free from clogging and delivering a uniform pattern. Apply UNIFORMLY the recommended rate of RO-NEET 6-E in 20 to 50 gallons of water per acre. Avoid overlaps that will increase RO-NEET 6-E dosage above recommended limits because plant injury will occur.

RO-NEET 6-E WITH FLUID FERTILIZER: RO-NEET 6-E may be combined with fluid (solution, slurry, or suspension) fertilizers. However, physical compatibility with these fluid fertilizers must be determined before combining in the spray tank. See Appendix I giving directions for these combinations. Even though found to be compatible, constant agitation is necessary to keep the RO-NEET 6-E evenly mixed with the fluid fertilizer.

IMPREGNATION ON DRY BULK FERTILIZERS: RO-NEET 6-E may be impregnated on many dry bulk fertilizers for use on sugar beets. RO-NEET impregnated on dry bulk fertilizers may be applied and incorporated into the soil either in the fall (in states where fall application for RO-NEET 6-E is recommended) before the ground freezes or before planting. However, uniform distribution of RO-NEET 6-E on fertilizer particles and uniform application are necessary to assure good results. See Appendix II for special instructions regarding directions for impregnation and use.

INCORPORATION DIRECTIONS

Immediately (within minutes) after application thoroughly mix RO-NEET 6-E into the soil to a depth of 2 to 3 inches. (California only: Incorporate to a depth of 3 inches.) Thorough soil mixing is necessary for good weed control.

RO-NEET 6-E must be incorporated into the soil immediately to prevent loss of the herbicide. Whenever possible, application and incorporation should be done in the same operation. Use equipment which has been proven to incorporate thoroughly to the recommended depth.

SOIL MIXING (INCORPORATION) BEFORE PLANTING: The following equipment commonly is used for soil mixing (incorporation) before planting:

For Broadcast (Overall) Application:

- Power Driven Cultivation Equipment (recommended on all soil types). Set to cut to a depth of 2 to 3 inches.
- **Tandem Discs** (recommended on all soil types). Set to cut to a depth of 4 to 6 inches, operated at 4 to 6 mph followed by a spiked-tooth harrow or sume other leveling device which extends beyond the ends of the discs. For more thorough mixing (for nutsedge, heavier soils) disc in two different directions (cross disc).

- Field Cultivators (recommended on lighter soils in good tilth only). Not recommended in California. Use 3 to 4 rows of sweeps, spaced at 7 inch or less intervals and staggered so that no soil is left unturned, followed by a spiked-tooth harrow pulled behind the cultivator. Do not use chisels or points. Set the cultivator to cut 4 inches deep, operated at 5 mph or more. Run the equipment over the field twice, the second run at an angle to the first.
- Rotary Ground Driven or Spring-Tooth Cultivators (recommended on lighter soils in good tilth only). Not recommended in California. Set to penetrate to a depth of 4 to 6 inches and operated at 5 to 8 mph in two different directions.

For Band (Row) Application:

Uniformly mix to a depth of 2 to 3 inches.

- Hooded power-driven rotary tillers.
- Hooded ground-driven rotary tillers.

Rolling:

Under most conditions, the treated area should be firmly rolled between the incorporation unit and the planters.

PLANTING

For maximum weed control benefits, crops should be planted or seeded immediately after application. Do not use a drag behind the planter as it may concentrate RO-NEET 6-E over the seed row and cause crop injury.

CULTUR' ACTICES FOLLOWING APPLICATION

RO-NEET 6-E is not persistent in the soil and susceptible weeds germinating later during the growing season may not be controlled. Shallow cultivation or approved postemergence herbicides may be necessary to control these susceptible weeds that escape control by RO-NEET 6-E as well as those weeds not susceptible to RO-NEET 6-E. Do not cultivate deeper than the depth of the herbicide incorporation.

When cultivating fields where RO-NEET 6-E has been banded in the row, use shields to prevent the movement of untreated soil into the treated row.

Page 10 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

DOSAGE RECOMMENDATIONS Solid Broadcast Treatment			
CROP	WHEN TO APPLY	GALLONS RO-NEET 6-E PER ACRE	REMARKS
Sugar Beets*	Preplant (soil incorporation) or at planting (soil injection)	1/2 to 2/3	Use on mineral soils ONLY. Use lower rate on sandy soils. Use higher dosage rate on heavier soil. Injury may result in highly saline or alkaline soils.
	Fall application	2/3	Recommended only in the states of Idaho, Minnesota, Montana, North Dakota, Oregon, Wyoming, and Washington. Apply and incorporate in late fall before the ground freezes.
Table Beets	Preplant (soil incorporation)	1/2 to 2/3	Use on mineral soils CNLY. Use lower rate on sandy soils. Use higher dosage rate on heavier soils. When applying RO-NEET 6-E in combination with fluid fertilizer, do not apply over 150 lbs. actual nitrogen per acre.
Spinach**	Preplant (soil incorporation	1/2	Use on sandy mineral soils only.

 Make only one (1) application per growing season. If RO-NEET 6-E is applied in the fall, do not reapply RO-NEET 6-E the following spring.

** Recommended on spinach in Arkansas, Connecticut, Delaware, Illinois, Maine, Maryland, Massachusetts, Mississippi, New Hampshire, New Jersey, New York, Ohio, Oklahoma, Pennsylvania, Texas, Vermont, Virginia and western Tennessee only.

Band Treatment:

The rates given above are for solid coverage basis. For band treatment, reduce the amount of RO-NEET 6-E proportionately depending on the row spacing and band width to be treated. DO NOT USE BAND APPLICATION ON ROCKY GROUND BECAUSE THOROUGH INCORPORATION IS NOT POSSIBLE.

Soil injection application on sugar beets - Special equipment designed for soil injection must be used. The injector units must be rigidly mounted immediately ahead or immediately behind the planter unit.

Soil Injection - Injector shanks must be spaced 2-1/2 to 3 inches apart and mounted in staggered positions to avoid trash building up on them. Injector shanks must be set to inject

RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

RO-NEET 6-E 1-1/2 to 2 inches below the soil surface. The width of the band in which weed control is desired will determine the number of injector shanks required per row (that is, four injector shanks for a 10 to 12-inch band, six injector shanks for a 15 to 18-inch band). The two injector shanks adjacent to the drill row must be 1-1/4 to 1-1/2 inches on either side of it.

RO-NEET 6-E AND EPTAM® 7-E SELECTIVE HERBICIDE TANKMIX FOR PREPLANT USE IN SUGAR BEETS FOR MICHIGAN, MINNESOTA, AND THE RED RIVER VALLEY AREA OF NORTH DAKOTA

The RO-NEET 6-E and EPTAM 7-E herbicide combination is a selective tankmix which must be incorporated into the soil. This combination controls weeds by interfering with normal germination and seedling development. This tank mixture does not control established weeds. The RO-NEET/EPTAM combination will give equal to or greater control of the weeds listed on this label than either product used separately. This tank mixture can be applied only once per growing season. The combination may be used in the fall or in the spring.

APPLICATION DIRECTIONS

During filling, pour the recommended rates of both RO-NEET 6-E and EPTAM 7-E into a properly calibrated, low-pressure boom sprayer having good agitation, and mix thoroughly. Apply the material uniformly in 10 to 50 gallons of water per acre. Check calibration frequently during application and observe the nozzles to ensure a uniform spray pattern. Do not use single nozzle boom-jet type sprayers. The soil should be well-worked prior to application and dry enough to permit thorough mixing with incorporation equipment.

SPRING PREPLANT INCORPORATED APPLICATION RATES* (Pints/Acre)			
SOIL TEXTURE**	ORGANIC MATTER%	EPTAM 7-E	RO-NEET 6-E
Loam or Coarser	3 to 4.5	1.25	3.3
	4.5>	1.25	4.0
Loam to Clay Loam	3 to 4.5	1.25	3.3
	4.5>	1.75	3.3
Clay Loam or Finer	3 to 4.5	1.75	3.3
	4.5>	2.25	3.3

* Make only one application per growing season. Do not apply the RO-NEET 6-E/EPTAM 7-E combination in the spring if either this combination or RO-NEET 6-E alone was applied in the fall.

** EPTAM 7-E should not be used on Loam or Coarser soils with less than 3% organic matter. RO-NEET 6-E at 4 to 5-1/3 pints/Acre should be used on these soils.

FALL PREPLANT INCORPORATED APPLICATION RATES* (Pints/Acre)				
SOIL TEXTURE	ORGANIC MATTER %	EPTAM 7-E	RO-NEET 6-E	
Loam or Coarser**	3 to 4.5 4.5>	1.25 1.25	4.0 4.0	
Loam to Clay Loam	3 to 4.5 4.5>	1.25 1.75	4.0 4.0	
Clay Loam or Finer 3 to 4.5 2.25 2.67 4.5> 2.25 4.0				

 Make only one application per growing season. Do not apply the RO-NEET 6-E/EPTAM 7-E combination in the spring if either this combination or RO-NEET 6-E alone was applied in the fall.

** EPTAM 7-E should not be used on Loam or Coarser soils with less than 3% organic matter. RO-NEET 6-E at 4 to 5-1/3 pints/Acre should be used on these soils.

SOIL INCORPORATION

The RO-NEET 6-E and EPTAM 7-E tank mixture must be immediately incorporated (mixed) into the top 2 to 3 inches of soil after spraying to prevent loss of the herbicides. Whenever possible, application and incorporation should be done in the same operation. Refer to the RO-NEET 6-E and EPTAM 7-E labels for complete incorporation procedures and the use of equipment which has been proven to incorporate thoroughly to the recommended depth.

GENERAL USE PRECAUTIONS

The RO-NEET 6-E and EPTAM 7-E combination may cause crop injury on very light sandy soil and when used under adverse environmental conditions that weaken crop seedlings. Refer also to RO-NEET 6-E and EPTAM 7-E labels for recommendations regarding cultural practices after application.

A tank mixture of RO-NEET 6-E and EPTAM 7-E will give equal to or greater control of the following listed weeds than either product used separately. This combination does not control established weeds.

Green foxtail Yellow foxtail Wild Oats Yellow nutsedge Purple nutsedge

Lambsquarters, common* Pigweed, redroot* Setaria viridis Setaria glauca Avena fatua Cyperus esculentus Cyperus rotundus

Chenopodium album Amaranthus retroflexus

*Partial control only (suppression).

Read both the RO-NEET 6-E and EPTAM 7-E labels carefully before using this tankmix. Observe all cautions and limitations on each label. The most restrictive labeling applies in tankmixes.

Do not contaminate water when disposing of equipment washwaters.

25

Page 14 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

٢

ĩ

APPENDIX I RO-NEET 6-E WITH FLUID FERTILIZERS

The following procedure is suggested for determining whether RO-NEET 6-E may be combined with a specific fluid fertilizer for spray tank application.

MATERIAL REQUIRED:

- 1. RO-NEET 6-E.
- 2. Fluid fertilizer to be used.
- 3. Adjuvant for fertilizer tankmix: Compex[™], Sponto 168-D[™], Unite[™], or equivalent. The adjuvant which provides the best emulsification depends on the specific fertilizer under consideration.
- 4. Two one-quart, wide-mouth glass jars with lid or stopper.
- 5. Measuring spoons (a 25 mL pipette or graduated cylinder provides more accurate measurement).
- 6. Measuring cup, 8 oz. (237 mL).

PROCEDURE:

- 1. Pour a pint (about 473 mL) of the fluid fertilizer into each of the quart jars.
- 2. Add adjuvant to one of the jars and mix (see rate table).
- 3. Add the RO-NEET 6-E to both jars (see rate table).
- 4. Close both jars with lid or stopper and mix the contents by turning the jars upside down ten times.
- 5. Inspect the surface and body of the mixtures
 - (A) Immediately after completing the jar inversions.
 - (B) After allowing the jars to stand quietly for 30 minutes.
 - (C) And then again after turning the jars upside down 10 times.

If the uniform mix cannot be made, the mixture should not be used. If either mixture remains uniform for 30 minutes, the combination may be used. Should either mixture separate after 30 minutes, but readily remix uniformly with the jar inversions, the mixture can be used if adequate agitation is maintained in the tank. If the mixture with adjuvant is satisfactory, but the one without adjuvant is not, be sure to use the adjuvant in the spray tank. Add the adjuvant first at a rate of 3 pints per 100 gallons of fluid fertilizer; foaming can be minimized by using moderate agitation.

If nondispersible oil, sludge or clumps of solids form in the mixtures, the combinations should not be used.

Page 15 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

<u>ş</u>.

r

i.

RATE TABLE FOR RO-NEET 6-E AND ADJUVANT** WITH THE FLUID FERTILIZER			
Gallons of fluid fertilizer to be applied per acre	mL or Tsp. of RO-NEET 6-E [*] to be added to 1 pint of fertilizer 6-E		
	mL	Tsp.	
10	8	1-2/3	
15	5	1	
20	4	3/4	
25	3	2/3	
30	2-1/2	1/2	
40	2	2/5	

* Based on field rate of 1 pound active ingredient per acre in the fertilizer volumes incicated. Increase volume proportionately to correspond with intended field rate in terms of pounds active ingredient per acre (e.g., for field rate of 4 pounds actual RO-NEET 6-E in 40 gallons fertilizer per acre, add 8 mL or 2 tsp. RO-NEET 6-E to each jar for compatibility testing purposes).

** Two (2) milliliters or one-half (1/2) teaspoon of adjuvant to be added to 1 pint of fluid fertilizer in order to equal the rate of 3 pints of adjuvant per 100 gallons of fluid fertilizer.

<u>...</u>

Page 16 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

APPENDIX II

RO-NEET 6-E IMPREGNATION ON DRY BULK FERTILIZERS

CAUTION: Nitrate fertilizers represent a potential explosive hazard, particularly in contact with organic substances such as RO-NEET 6-E. Do not impregnate RO-NEET 6-E on ammonium nitrate. Do not use fertilizers containing ammonium, potassium or sor⁷. Jm nitrate. Such mixtures may cause explosion.

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the RO-NEET fertilizer mixtures.

RO-NEET 6-E may be impregnated on many dry bulk fertilizers (Table 1) and applied and incorporated into the soil before planting for the control of labeled grasses and broadleaf weeds in sugar beets.

All RO-NEET 6-E label and supplementary literature instructions and precautions regarding rates per acre, soil type and soil incorporation, application, and other directions must be followed.

Test results have shown RO-NEET 6-E on many dry bulk fertilizers gives weed control equal to RO-NEET 6-E applied as a spray in water or liquid fertilizer. However, uniform impregnation of the pesticide on the dry fertilizer particles and uniform application in the field are necessary to assure good results.

A minimum of 200 pounds and a maximum of 700 pounds of approved fertilizer ingredients (Table 2) impregnated with the appropriate amount of RO-NEET 6-E must be applied per acre.

For impregnating the pesticides on dry fertilizens, use a closed rotary-drum type mixer equipped with suitable spraying equipment. The spray nozzles should be positioned inside of the mixer to provide uniform spray coverage of the tumbling fertilizer. The RO-NEET 6-E should be sprayed uniformly onto the fertilizer using a fine spray pattern.

The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with RO-NEET 6-E provides a satisfactory dry mixture.

If the absorptivity is not adequate, use of a highly absorptive powder is required to provide a dry, free-flowing mixture. Micro-Cel[™] E is the recommended absorbent powder. It should be added separately and uniformly to the prepared pesticide-fertilizer mixture, in a quantity that is sufficient to provide a suitable free-flowing mixture. Generally, less than 2% by weight of Micro-Cel E is required.

The amount of RO-NEET 6-E actually required in the manufacture of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amounts of RO-NEET actually contained in the mixture applied to the soil represent the correct rates of use.

Page f7 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

Bulk fertilizers impregnated with RO-NEET 6-E should be *applied immediately*, NOT STORED. All bulk containers should be tightly covered while the products are being transported and applied to reduce chances of RO-NEET loss via volatilization.

APPROVED DRY FERTILIZER INGREDIENTS FOR USE WITH RO-NEET 6-E			
	<u>N</u>	<u>P</u>	K
Ammonium sulfate	21	0	0
Ammonium Phosphate-sulfate	16	20	0
Diammonium ohosphate	18	46	0
Monoammonium Phosphate	11	56	0
Potassium chloride	0	0	60
Potassium sulfate	0	0	[′] 52
Single superphosphate	0	20	0
Treble superphosphate	0	46	0
Urea	45	0	0

Table 1.

NOTE: SUL-PO-MAG, K-MAG and 11-48-0 have been shown to be compatible with RO-NEET 6-E and are approved for use.

RO-NEET 6-E Physical Data

Specific Gravity 20/20°C: Pounds/gallon, 20/20°C: Flashpoint: Viscosity Freeze Point:

0.976 8.12 168°F (Tagliabue Closed Cup) Sprayable down to 25°F 20°F (seeded)

Page 18 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D 29

7

Table 2.

RATE CHART FOR THE IMPREGNATION OF DRY BULK FERTILIZERS WITH RO-NEET 6-E:			
	RO-NEET 6-E Rate Per Acre		
Fertilizer Rate Per Acre	2 quarts per acre	2-2/3 quarts per acre	
200 lbs.	20 quarts/ton	26-2/3 quarts/ton	
250 lbs.	16 quarts/ton	21-1/3 quarts/ton	
300 lbs.	13-1/3 quarts/ton	17-3/4 quarts/ton	
350 lbs.	11-2/5 quarts/ton	15-1/4 quarts/ton	
400 lbs.	10 quarts/ton	13-1/3 quarts/ton	
450 lbs.	8-7/8 quarts/ton	11-7/8 quarts/ton	
500 lbs.	8 quarts/ton	10-2/3 quarts/ton	
550 lbs.	7-1/3 quarts/ton	9-3/4 quarts/ton	
600 lbs.	6-2/3 quarts/ton	8-7/8 quarts/ton	
650 lbs.	6-1/5 quarts/ton	8-1/4 quarts/ton	
700 lbs.	5-3/4 quarts/ton	7-5/8 quarts/ton	

4

.4

Page 19 RO-NEET 6-E Booklet RON178B.RSD - RS-052693D

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Keep containers closed when not in use. Do not store near seeds, fertilizers or foodstuffs. Protect from temperatures below 20°F, but do not store near sources of heat or open flame.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

FOR BULK AND MINI-BULK CONTAINERS

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage, such as cracks, punctures, burges dents, abrasions, and damaged or worn threads on closure devices.

REFILL ONLY WITH RO-NEET 6-E. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than RO-NEET 6-E will result in contamination and may weaken container.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIABILITY of Seller will be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLEP'3 SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States of America.

EPTAM® and RO-NEET® are trademarks of ZENECA Group Companies. Compex[™] is a trademark of KALO, Inc. Micro-Cel[™] is a trademark of Manville Sales Corporation. Sponto 168-D[™] is a trademark of Witco Corporation. Unite[™] is a trademark of HACO. Inc.