## **ZENECA** Ag Products

# RO-NEET® 6-E

**Selective Herbicide** 

Contains 6 pounds active ingredient per gallon. EPA Reg. No. 10182-178

EPA Heg. No. 10182-178 EPA Est. No. 10182-NB-1

Liquid Herbicide for Weed Control in Sugar Beets, Table Beets, and Spinach

KEEP OUT OF REACH OF CHILDREN CAUTION



See side and back panels for Statement of Practical Treatment and Precautionary Statements. ACCEPTED

WAY 7 ....
Under the Federal Insecting to Acre

Under the Federal Inserting, Fungicide, and Follow in Act, as amended, for the posteride registered under EPA Reg. No. 10182-178

Net Contents: 2.5 gái (9.46 L)

BEST AVAILABLE COPY

## **ZENECA** Ag Products

## RO-NEET® 6-E

Selective Herbicide

Liquid Herbicide for Weed **Control in Sugar Beets,** Table Beets, and Spinach

### COMPLETE DIRECTIONS FOR USE



Under the Federal Insecticide,

Fungicide, and Rodenticide Act. as amended, for the posticide registered under EPA Reg. No. 70/2277

079308Z

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:
The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks reterred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories). Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIABILITY of Seller with the claim of Celler with the claim of t be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States of America.

EPTAM® and RO-NEET® are trademarks of ZENECA Group Companies.
Compex™ is a trademark of KALO, Inc.
Micro-Cel™ is a trademark of Manville Sales Corporation.
Sponto 168-D™ is a trademark of Witco Corporation.
Unite™ is a trademark of HACO, Inc.
@1993 ZENECA Inc.

## **CONTEN**

Statement of Pr

Precautionary S

Directions for U

Soil Prepara

Application.

Dry Fertilize

General Use Pr

**Weeds Controll** 

Incorporation Di

Planting.....

**Cultural Practice** 

Dosage Recom

Appendix I......

Appendix II......

Storage and Dis

BEST AVAIL TO LE C

ላ)

KEEP OUT OF REACH OF CHILDREN CAUTION

**ACTIVE INGREDIENT:** 

Contains 6 pounds active ingredient per gallon.

EPA Reg. No. 10182-178

## STATEMENT OF PRACTICAL TREATMENT FIRST AID

If a known exposure occurs or is suspected, immediately start the recommended procedures below. Simultaneously contact a Poison Control Center, a physician, or the nearest hospital. Describe the situation and follow the advice given.

IF SWALLOWED: Immediately give large quantities of water but DO NOT induce vomiting. This product contains hydrocarbon solvent. If vomiting does occur, give fluids again. Have a physician determine if condition of patient will permit evacuation of stomach. DO NOT give anything by mouth to an unconscious person. Call a phys. an immediately.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for 15 minutes. Get medical attention if irritation occurs.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

2

HARMFUL IF SWALLOWED. Avoid contact with skin, eyes, and clothing. Avoid contamination of feed or food.

#### **Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or viton
- · Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## User Safety Recommendations Users should:

- Wash hands before eating, drinking, ing gum, using tobacco or using the to
- Remove clothing immediately if pes gets inside. Then wash thoroughly and clean clothing.
- Remove PPE immediately after had this product. Wash the outside of glow fore removing. As soon as possible, thoroughly and change into clean clot

#### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to area surface water is present or to intertidal area the mean high water mark. Do not cont water when disposing of equipment washy

**READ ENTIRE LABEL BEFORE US** 

Protect from temperatures below 20° Product crystallizes at lower temperatures Warm or store at higher temperatures mix to redissolve crystals and assur uniformity before use.

## **DIRECTIONS FOR US**

It is a violation of Federal law to use this in a manner inconsistent with its labeling.

Do not apply this product in a way that will workers or other persons, either direct through drift. Only protected handlers may the area during application. For any requispecific to your State or Tribe, consult the responsible for pesticide regulation.

Do not apply this product through any typication stratem.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protection equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or viton
- Shoes plus socks

6

### **GENERAL INFORMATION**

RO-NEET\* 6-E is a selective herbicide which is mixed (incorporated) into the soil or applied below the surface of the soil for control of weeds listed on this label. RO-NEET 6-E controls weeds by interfering with normal seed germination and seedling development. It does not control established or germinated weeds present at application. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

## GENERAL USE PRECAUTIONS

Not for use or storage in or around the home.

RO-NEET 6-E should be used only for recommended purposes and at recommended rates (DO NOT OVERDOSE). Applied according to directions and under normal growing conditions, RO-NEET 6-E will not harm the treated crop. During germination and early growth, extended periods of unusually cold and wet or hot and dry weather, insect, nematode, or plant disease attack, carryover pesticide residues, the use of certain soil applied systemic insecticides, highly saline or alkaline soil conditions, improperly placed fertilizers or soil insecticides may create abnormal conditions that weaken crop seedlings. RO-NEET 6-E used under these abnormal conditions could result in crop injury.

RO-NEET 6-E may cause crop injury on very light sandy soil.

Tank mix this product with fungicides, insecticides, or herbicides only as recommended.

Do not contaminate irrigation water or water used for domestic purposes.

## **WEEDS CONTROLLED**

#### **ANNUAL GRASSES:**

Annual Bluegrass
Annual Ryegrass
Barley, Volunteer
Barnyardgrass
Poa annua
Lolium multiflorum
Hordeum spp.
Echinochloa spp.

(Watergrass)

Crabgrass Digitaria spp.
Foxtails Setaria spp.
Wild Oats Avena fatua

#### **ANNUAL BROADLEAF WEEDS:**

Black Nightshade
Hairy Nightshade
Henbit, Deadnettle
Lamium spp.
Chegopodium albi

Lambsquarters
Nettleleaf Goosefoot\*
Common Purslane
Redroot Pigweed
Shepherd's-purse
Small Stinging Nettle,

Chenopodium album
Chenopodium murale
Portulaca oleracea
Amaranthus retroflexus
Capsella bursa-pastoris
Urtica urens

**Burning Nettle** 

Velvetleaf\*\* Abutilon theophrasti

\*California excluded.

The above broadleaf weeds will be controlled only if the application is made when conditions are favorable for germination. RO-NEET 6-E SELECTIVE HERBICIDE WILL NOT CONTROL ESTABLISHED OR GERMINATED WEEDS PRESENT AT APPLICATION.

#### **PERENNIAL WEEDS:**

Yellow Nutsedge Cyperus esculentus
Purple Nutsedge Cyperus rotundus

Existing stands of nutsedge must be turned under and chopped thoroughly before treatment. Prior to planting, RO-NEET 6-E must be thoroughly mixed into the soil or applied subsurface to the soil with injectors for nutsedge control.

TOT AVAILABLE COPY

<sup>\*\*</sup>Partial control or suppression only.

7

8

SOIL PREPARATION: Prepare the soil for seeding according to good agricultural practice. All weed growth should be thoroughly worked into the soil before treatment. DO NOT APPLY BEFORE PREIRRIGATION.

APPLICATION: Apply RO-NEET 6-E to well-worked soil which is dry enough to permit thorough mixing with incorporation equipment or proper injection. Use standard low-pressure (20-50 psi) boom spray equipment. DO NOT USE SINGLE NOZZLE BOOM-JET TYPE SPRAYERS. Application equipment should be carefully calibrated before use and checked frequently during application to be sure nozzles are free from clogging and delivering a uniform pattern. Apply UNIFORMLY the recommended rate of RO-NEET 6-E in 20 to 50 gallons of water per acre. Avoid overlaps that will increase RO-NEET 6-E dosage above recommended limits because plant injury will occur.

RO-NEET 6-E WITH FLUID FERTILIZER: RO-NEET 6-E may be combined with fluid (solution, slurry, or suspension) fertilizers. However, physical compatibility with these fluid fertilizers must be determined before combining in the spray tank. See Appendix I giving directions for these combinations. Even though found to be compatible, constant agitation is necessary to keep the RO-NEET 6-E evenly mixed with the fluid fertilizer.

IMPREGNATION ON DRY BULK FERTILIZERS: RO-NEET 6-E may be impregnated on many dry bulk fertilizers for use on sugar beets. RO-NEET impregnated on dry bulk fertilizers may be applied and incorporated into the soil either in the fall (in states where fall application for RO-NEET 6-E is recommended) before the ground freezes or before planting. However, uniform distribution of RO-NEET 6-E on fertilizer particles and uniform application are necessary to assure good results. See Appendix II for special instructions regarding directions for impregnation and use.

## INCORPORATION DIRECTIONS

Immediately (within minutes) after application thoroughly mix RO-NEET 6-E into the soil to a depth of 2 to 3 inches. (California only: Incorporate to a depth of 3 inches.) Thorough soil mixing is necessary for good weed control.

RO-NEET 6-E must be incorporated into the soil immediately to prevent loss of the herbicide. Whenever possible, application and incorporation should be done in the same operation. Use equipment which has been proven to incorporate thoroughly to the recommended depth.

SOIL MIXING (INCORPORATION) BEFORE PLANTING: The following equipment commonly is used for soil mixing (incorporation) before planting:

For Broadcast (Overall) Application:

- Power Driven Cultivation Equipment (recommended on all soil types). Set to cut to a depth of 2 to 3 inches.
- Tandem Discs (recommended on all soil types). Set to cut to a depth of 4 to 6 inches, operated at 4 to 6 mph followed by a spikedtooth harrow or some other teveling device which extends beyond the ends of the discs. For more thorough mixing (for nutsedge, heavier soils) disc in two different directions (cross disc).
- Field Cultivators (recommended on lighter soils in good tilth only). Not recommended in California. Use 3 to 4 rows of sweeps, spaced at 7 inch or less intervals and staggered so that no soil is left unturned, followed by a spiked-tooth harrow pulled behind the cultivator. Do not use chisels or points. Set the cultivator to cut 4 inches deep, operated at 5 mph or more. Run the equipment over the field twice, the second run at an angle to the first.

 Rotary Ground Driven or Spring vators (recommended on lighte tilth only). Not recommended in to penetrate to a depth of 4 to operated at 5 to 8 mph in two diffe

For Band (Row) Application:

Uniformly mix to a depth of 2 to 3 in

- Hooded power-driven rotary tillers
- Hooded ground-driven rotary tiller

Rolling:

Under most conditions, the treated a firmly rolled between the incorpor the planters.

### **PLANTING**

For maximum weed control benefits be planted or seeded immediately a except for fall applications. Do not use the planter as it may concentrate over the seed row and cause crop in

## CULTURAL PRACTIC FOLLOWING APPLICA

RO-NEET 6-E is not persistent in the ceptible weeds germinating later duri season may not be controlled. Shall and approved postemergence herb necessary to control those susce that escape control by RO-NEET & those weeds not susceptible to RO-not cultivate deeper than the depth cide incorporation.

When cultivating fields where RO-Nobeen banded in the row, use shields movement of untreated soil into the t

9

| DOSAGE RECOMMENDATIONS Solid Broadcast Treatment |                               |                                    |   |
|--|-------------------------------|------------------------------------|---|
| CROP   | WHEN TO APPLY                 | GALLONS<br>RO-NEET 6-E<br>PER ACRE | REMARKS   |
| Table Beets                                      | Preplant (soil incorporation) | ½ to ¾                             | Use on mineral soils ONLY. Use lower rate on sandy soils. Use higher dosage rate on heavier soils. When applying RO-NEET 6-E in combination with fluid fertilizer, do not apply over 150 lbs. actual nitrogen per acre. |
| Spinach**  | Preplant (soil incorporation) | 1/2                                | Use on sandy mineral soils only.  |

<sup>\*\*</sup>Recommended on spinach in Arkansas, Colorado, Connecticut, Delaware, Illinois, Maine, Maryland, Massachusetts, Mississippi, New Hampshire, New Jersey, New York, Ohio, Oklahoma, Pennsylvania, Texas, Vermont, Virginia and western Tennessee only.

| DOSAGE RECOMMENDATIONS Solid Broadcast Treatment |   |                                    |   |
|--|---|------------------------------------|---|
| CROP   | WHEN TO APPLY   | GALLONS<br>RO-NEET 6-E<br>PER ACRE | REMARKS   |
| Sugar Beets*                                     | Preplant (soil incorporation) or at planting (soil injection) | % to %                             | Use on mineral soils ONLY. Use lower rate on sandy soils. Use higher dosage rate on heavier soil. Injury may result in highly saline or alkaline soils.                   |
|  | Fall application  | **                                 | Recommended only in the states of Idaho, Minnesota, Montana, North Dakota, Oregon, Wyoming, and Washington. Apply and incorporate in late fall before the ground freezes. |

\*Make only one (1) application per growning season. If RO-NEET 6-E is applied in the fall, do not reapply RO-NEET 6-E the following spring.

applied only once per growing season. The combination may be used in the fall or in the spring. EPTAM 7-E should not be used on soils with less than 3% organic matter. RO-NEET 6-E at 4 to 5 ½ pints/acre should be used on these soils.

#### **APPLICATION DIRECTIONS**

During filling, pour the recommended rates of both RO-NEET 6-E and EPTAM 7-E into a properly calibrated, low-pressure boom sprayer having good agitation, and mix thoroughly. Apply the material uniformly in 10 to 50 gallons of water per acre. Check calibration frequently during application and observe the nozzles to ensure a uniform spray pattern. Do not use single nozzle boom-jet type sprayers. The soil should be well-worked prior to application and dry enough to permit thorough mixing with incorporation equipment.

| SPRING* PREPLANT INCORPORATED APPLICATION RATES** (Pints/Acre***) |                  |             |            |  |
|---|------------------|-------------|------------|--|
| SOIL  | ORGANIC MATTER % | EPTAM       | RO-NEET    |  |
| TEXTURE   |                  | 7-E         | 6-E        |  |
| Coarse  | 3.0 to 4.5       | 1.0         | 3.0        |  |
|   | >4.5             | 1.0 to 1.25 | 3.0        |  |
| Medium  | 3.0 to 4.5       | 1.0 to 1.25 | 3.0 to 4.0 |  |
|   | >4.5             | 1.0 to 1.75 | 3.0 to 4.0 |  |
| Fine  | 3.0 to 4.5       | 1.0 to 1.5  | 3.0 to 4.0 |  |
|   | >4.5             | 1.0 to 2.0  | 3.0 to 4.0 |  |

#### \*DO NOT USE THE TANKMIX IN THE SPRING IN MICHIGAN.

| FALL PREPLANT INCORPORATED APPLICATION RATES* (Pints/Acre**) |            |             |             |
|--|------------|-------------|-------------|
| SOIL   | ORGANIC    |             | RO-NEET     |
| TEXTURE  | MATTER %   |             | 6-E         |
| Coarse   | 3.0 to 4.5 | 1.0         | 4.0         |
|  | >4.5       | 1.0 to 1.25 | 4.0 to 4.33 |
| Medium   | 3.0 to 4.5 | 1.0 to 1.25 | 4.0 to 4.5  |

>4.5

3.0 to 4.5

>4.5

\*Make only one application per growing season. Do not apply the RO-NEET 6-E/EPTAM 7-E combination in the spring if either this combination or RO-NEET 6-E alone was applied in the fall.

1.0 to 1.5

1.0 to 1.5

1.0 to 2.0

4.0 to 4.5

3.0 to 4.5

#### SOIL INCORPORATION

Fine

The RO-NEET 6-E and EPTAM 7-E tank mixture must be immediately incorporated (mixed) into the top 2 to 3 inches of soil after spraying to prevent loss of the herbicides. Whenever possible, application and incorporation should be done in the same operation. Refer to the RO-NEET 6-E and EPTAM 7-E labels for complete incorporation procedures and the use of equipment which has been proven to incorporate thoroughly to the recommended depth.

#### **GENERAL USE PRECAUTIONS**

The RO-NEET 6-E and EPTAM 7-E combination may cause crop injury on very light sandy soil and when used under adverse environmental conditions that weaken crop seedlings. Refer also to RO-NEET 6-E and EPTAM 7-E labels for recommendations regarding cultural practices after application.

A tank mixture of RO-NEET 6-E and EPTAM 7-E will give equal to or greater control of the following listed weeds than either product used separately. This combination does not control established weeds.

Green foxtail
Yellow foxtail
Wild Oats
Yellow nutsedge
Purple nutsedge
Lambsquarters,
common\*
Pigweed, redroot\*

\*Partial control only (sup Read both the ROlabels carefully be Observe all cautions The most restrictive la Do not contaminate equipment washwate

## APF

#### **RO-NEET 6-E WIT**

The following proceduring whether RO-NI a specific fluid fertilize MATERIAL REQUIR

#### 1. RO-NEET 6-E.

- 1. NO-NEET O-E.
- 2. Fluid fertilizer to b
- Adjuvant for fert Sponto 168-D™, adjuvant which sification dependent under consideration
- Two one-quart, w or stopper.
- Measuring spoons cylinder provides r
- Measuring cup, 8

<sup>\*\*</sup>Make only one application per growing season. Do not apply the RO-NEET 6-E/EPTAM 7-E combination in the spring if either this combination or RO-NEET 6-E alone was applied in the fall.

<sup>\*\*\*</sup>Do not apply more than 5 pints of RO-NEET 6-E and EPTAM 7-E combined regardless of the ratio.

<sup>\*\*</sup>Do not apply more than 6 pints of RO-NEET 6-E and EPTAM 7-E combined regardless of the ratio.

#### PROCEDURE:

- 1. Pour a pint (about 473 mL) of the fluid fertilizer into each of the quart jars.
- 2. Add adjuvant to one of the jars and mix (see rate table).
- 3. Add the RO-NEET 6-E to both jars (see rate table).
- Close both jars with lid or stopper and mix the contents by turning the jars upside down ten times.
- 5. Inspect the surface and body of the mixtures
  - (A) Immediately after completing the jar inversions.
  - (B) After allowing the jars to stand quietly for 30 minutes.
  - (C) And then again after turning the jars upside down 10 times.

If the uniform mix cannot be made, the mixture should not be used. If either mixture remains uniform for 30 minutes, the combination may be used. Should either mixture separate after 30 minutes, but readily remix uniformly with the jar inversions, the mixture can be used if adequate agitation is maintained in the tank. If the mixture with adjuvant is satisfactory, but the one without adjuvant is not, be sure to use the adjuvant in the spray tank. Add the adjuvant first at a rate of 3 pints per 100 gallons of fluid fertilizer; foaming can be minimized by using moderate agitation.

If nondispersible oii, sludge or clumps of solids form in the mixtures, the combinations should not be used.

18

%

| RATE TABLE FOR RO-NEET 6-E<br>AND ADJUVANT** WITH THE<br>FLUID FERTILIZER   |     |      |  |
|---|-----|------|--|
| Gallons of fluid mL or Tsp. of RO-NEET 6-E* fertilizer to be to be added to applied per acre 1 pint of fertilizer |     |      |  |
|   | 6-E |      |  |
|   | mL  | Tsp. |  |
| 10  | 8   | 1%   |  |
| 15  | 5   | 1    |  |
| 20  | 4   | %    |  |
| 25  | 3   | 3/4  |  |
| 30  | 2½  | 1/2  |  |

\*Based on field rate of 1 pound active ingredient per acre in the fertilizer volumes indicated. Increase volume proportionately to correspond with intended field rate in terms of pounds active ingredient per acre (e.g., for field rate of 4 pounds actual RO-NEET 6-E in 40 gallons fertilizer per acre, add 8 mL or 2 tsp. RO-NEET 6-E to each jar for compatibility testing purposes).

2

40

"Two (2) milliliters or one-half (%) teaspoon of adjuvant to be added to 1 pint of fluid fertilizer in order to equal the rate of 3 pints of adjuvant per 100 gallons of fluid fertilizer.

### **APPENDIX II**

## RO-NEET 6-E IMPREGNATION ON DRY BULK FERTILIZERS

CAUTION: Nitrate fertilizers represent a potential explosive hazard, particularly in contact with organic substar.ces such as RO-NEET 6-E. Do not impregnate RO-NEET 6-E on ammonium nitrate. Do not use fertilizers containing ammonium, potassium or sodium nitrate. Such mixtures may cause explosion.

All individual state regulations relating fertilizer blending, registration, labeling, tion are the responsibility of the indivicompany selling the RO-NEET fertilize RO-NEET 8-E may be impregnated or

bulk fertilizers (Table 1) and applied a rated into the soil before planting for the labeled grasses and broadleaf weeds in

All RO-NEET 6-E label and supplementure instructions and precautions regarder acre, soil type and soil incorporation, and other directions must be followed the results have shown RO-NEET 6-

dry bulk fertilizers gives weed contraction. RO-NEET 6-E applied as a spray in watertilizer. However, uniform impregnate pesticide on the dry fertilizer particles application in the field are necessar good results.

A minimum of 200 pounds of approvingredients (Table 2) impregnated with priate amount of RO-NEET 6-E must per acre.

For impregnating the pesticides on druse a closed rotary-drum type mixer education suitable spraying equipment. The sp should be positioned inside of the mixed uniform spray coverage of the tumbling The RO-NEET 6-E should be sprayed onto the fertilizer using a fine spray pat

The physical properties of fertilizers valuarly in liquid absorptive capacity. Writivity is sufficient, simple spray impregriertilizer with RO-NEET 6-E provides a dry mixture.

If the absorptivity is not adequate, use absorptive powder is required to profree-flowing mixture. Micro-Cel<sup>TM</sup> E is mended absorbent powder. It should

17

separately and uniformly to the prepared pesticide-fertilizer mixture, in a quantity that is sufficient to provide a suitable free-flowing mixture. Generally, less than 2% by weight of Micro-Cel E is required.

The amount of RO-NEET 6-E actually required in the manufacture of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amounts of RO-NEET actually contained in the mixture applied to the soil represent the correct rates of use.

Bulk fertilizers impregnated with RO-NEET 6-E should be *applied immediately*, NOT STORED. All bulk containers should be tightly covered while the products are being transported and applied to reduce chances of RO-NEET loss via volatilization.

Table 1.

| APPROVED DRY FERTILIZER<br>INGREDIENTS FOR USE WITH<br>RO-NEET 6-E |          |    |    |
|--|----------|----|----|
| -  | <b>N</b> | ٩  | K  |
| Ammonium sulfate   | 21       | 0  | 0  |
| Ammonium Phosphate-sulfate   | 16       | 20 | 0  |
| Diammonium phosphate   | 18       | 46 | 0  |
| Monoammonium Phosphate   | 11       | 56 | 0  |
| Potassium chloride   | 0        | 0  | 8  |
| Potassium sulfate  |          | 0  | 52 |
| Single superphosphate  | 0        | 20 | 0  |
| Treble supernhasphate  | 0        | 46 | 0  |
| Urea   | 45       | 0  | 0  |

NOTE: SUL-PO-MAG, K-MAG and 11-48-0 have been shown to be compatible with RO-NEET 6-E and are approved for use.

#### **RO-NEET 6-E Physical Data**

Specific Gravity 20/20°C: 0.97 Pounds/gallon, 20/20°C: 8.12

Flashpoint:

168°F (Tag!iabue

Closed Cup)

Viscosity:

Sprayable down

to 25°F

Freeze Point:

20°F (seeded)

Table 2.

| RATE CHART FOR THE IMPREGNATION OF DRY BULK FERTILIZERS WITH RO-NEET 6-E |                           |                |  |
|--|---------------------------|----------------|--|
| Fertilizer   | RO-NEET 6-E Rate Per Acre |                |  |
| Rate   | 2 quarts                  | 2% quarts      |  |
| Per Acre   | per acre                  | per acre       |  |
| 200 lbs.   | 20 quarts/ton             | 26% quarts/ton |  |
| 250 lbs.   | 16 quarts/ton             | 21% quarts/ton |  |
| 300 lbs.   | 13% quarts/ton            | 17% quarts/ton |  |
| 350 lbs.   | 11% quarts/ton            | 15% quarts/ton |  |
| 400 lbs.   | 10 quarts/ton             | 13% quarts/ton |  |
| 450 lbs.   | 8% quarts/ton             | 11% quarts/ton |  |
| 500 lbs.   | 8 quarts/ton              | 10% quarts/ton |  |
| 550 lbs.   | 7% quarts/ton             | 9% quarts/ton  |  |
| 600 lbs.   | 6% quarts/ton             | 8% quarts/ton  |  |
| 650 lbs.   | 6% quarts/ton             | 8¼ quarts/ton  |  |
| 700 lbs.   | 5% quarts/ton             | 7% quarts/ton  |  |

## STORAGE AND DISPO

PROHIBITIONS: Do not contaminate food, or feed by storage or disposa dumping is prohibited.

STORAGE: Keep containers closed w in use. Do not store near seeds, fertifoodstuffs. Protect from temperature: 20°F, but do not store near sources of open flame. COMBUSTIBLE LIQUID.

PESTICIDE DISPOSAL: Waste resulti the use of this product may be dispose site or at an approved waste disposal

**CONTAINER DISPOSAL:** Triple ri equivalent); then offer for recycling or tioning, or puncture and dispose of in a landfill, or by other procedures approstate and local authorities.

FOR BULK AND MINI-BULK CONTA CONTAINER DISPOSAL: Reseal of and offer for reconditioning, or triple requivalent) and offer for recycling or toning, or clean in accordance with returner's instructions.

CONTAINER PRECAUTIONS: Before inspect thoroughly for damage, such as punctures, bulges, dents, abrasions, at aged or worn threads on closure device REFILL ONLY WITH RO-NEET 6-E. Thents of this container cannot be contained.

tents of this container cannot be cor removed by cleaning. Refilling with m other than RO-NEET 6-E will result in mination and may weaken container.

After filling and before transporting, check f

Do not refill or transport damaged ing container.

CONTAINER IS NOT SAFE FOR FEED, OR DRINKING WATER!

#### KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

#### FIRST AND-

If a known exposure occurs or is suspected, immediately start the recom-mended procedures below. Simultaneously contact a Poison Control Center, a physician, or the nearest hospital. Describe the situation and

IF SWALLOWED: Immediately give large quantities of water but DO NOT. induce varieing. This product contains hydrocarbon solvent, if varning does occur, one fluids again. Have a physician determine if condition of patient will permit evacuation of stomach. DO NOT give anything by outh to an unconscious person. Call a physician imme-

IF IN EYES: If id eyelds apart and flush eyes with large amounts of runon water fry 15 minutes. Get medical attention of imitation occurs. F ON SION: Wash with planty of soap and water. Get medical attention if

IF WHALED: Remove to fresh air. Seek medical attention if respiratory intelion occurs

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE. CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

#### **PRECAUTIONARY STATEMENTS**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

MARRIFUL IF SWALLOWED. Avoid contact with skin, eyes and clothing Avoid contamination of feed or food Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed. below. If you went more options, follow the instructions for category E on an EPA chemical resistance category selection chart Applicators and other handlers must weer

Long-sleeved shirt and long pants

 Chemical-registant gloves, such as barner laminate or nitrie rubber. or neoprene rubber or viton

Follow manufacturer's instructions for cleaning/maintaining PPE til no auch instructions for washables, use detergent and hot water. Keep and wash PPE separately from other leundry

then handlers use closed systems, enclosed cabs, or aircraft in a man ner that meets the requirements listed in the Worker Protection Standard WPS) for agricultural pesticides [40 CFR 170 240(d)(4-8)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations Users should

- . Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if posticide gets inside. Then wash
- thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washingters.

READ ENTINE LABEL SEFORE USING Protect from temperatures below 20°F. Product crystallizes at lower temperatures Warm or store at higher temperatures and mix to redissolve crystals and assure uniformity before use

with its labeling

> Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of impation system

AGRICU'.TURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling a attached bookiet) under 'AGRICULTURAL USE REQUIRE MENTS" in the DIRECTIONS FOR USE section for information about this standard

Refer to attached booklet for use directions

#### STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or dis posal. Open dumping it prohibited ...

STORAGE: Keep containers closed when not in use. Do not store near seeds, fertilizers or foodsaults. Protect from temperatures below 20°F, but do not store near sources of heat or open flame

COMBUSTIBLE LIQUIO

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be decored of on site or at an approved waste disposal facility CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or

trate rines (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions FOR BULK AND MINI-BULK CONTAINERS

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices

REFILL ONLY WITH RO-NEET® 6-E. The contents of this container canno be completely removed by cleaning. Refilling with materials other than RO-NEET 6-E will result in contamination and may weaken container After filling and before transporting, check for leaks

Do not refill or transport damaged or leaking container CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

IMPORTANT: Read the entire Directions for Use and Conditions of Sale and Warranty before using this product constitutes or SALE AND LIMITED WARRANT

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all notes inherently associated with the use of this product. Once mury, ineffectiveness or other unintended consequences may result because of such factors as immig and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influenc my factors in the use of the product, all of which are beyond the control of the safler. All such rate and to personal by the Book and Char and Book and Union and Safler harmines for any claims relating to such factors.

Seller warrents that this product conforms to the chemical description on the label and is probly fit for the purposes stated on the label subject to the inherent raks n to above, when used in accordance with directions under normal conditions of use. This wavenity does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of the Seller £ of Buyer and User assume the ret of any such use. SELLER DISCLAMIS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITHESS OR MERCHANTABILITY

When Buver or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theoneal), Buyer or User must premptly notify in writing Select of any claims to be elepted to receive either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIMITLITY of Sefer will be, at the election of Sefer, refund of the purchase price paid for product bought, or replacement of amount of product used SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL SPECIAL OR INDIRECT MAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLEN'S SOLE LIMINATY AND HAVEN'S AND UNEN'S EXCLUSIVE HENELY SHALL SE UNITED TO THE REPUND OF THE PURCHASE PRICE

The product is sold ordy for uses stated on its label. He express or maked bronce is granted to use or sell this product under any patent in any country incept as specified Country United States of America

RO-NEETS is a trademark of a ZENECA Group Company

**C1993 Zenece Inc.** 

#### IMPORTANT

See attached booklet for Use Precautions, **Directions for Use and Recommendations** Read all label directions before using.



PLACE Complete Directions for Use **Booklet** Here

\_\_\_\_\_

CONTENTS:

ZENECA A Products

érésesz

⊕ ACCEPTED Under the Description of 1 ... 7. ~1. 70; ·

BEST AVAILABLE COPY

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling (i.e., attached booklet) under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

Refer to attached booklet for use directions.

#### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**STORAGE:** Keep containers closed when not in use. Do not store near seeds, fertilizers or foodstuffs. Protect from temperatures below 20°F, but do not store near sources of heat or open flame. COMBUSTIBLE LIQUID.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, an of which are beyond the control of the Selfer. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Selfer harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warrantly does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably to eseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims loss is or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set torth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIABILITY of Seller will be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPEC, AL OH, INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States of America.

RO-NEET is a trademark of a ZENECA Group Company. ©1993 ZENECA Inc.

**ZENECA** Ag Products

ZENECA Inc. Wilmington, Delaware 19897

1.073308Z

#### STATEMENT OF PRACTICAL TREATMENT

#### **FIRST AID**

)

If a known exposure occurs or is suspected, immediately start the recommended procedures below. Simultaneously contact a Poison Control Center, a physician, or the nearest hospital. Describe the situation and follow the advice given.

IF SWALLOWED: Immediately give large quantities of water but DO NOT induce vomiting. This product contains hydrocarbon solvent. If vomiting does occur, give fluids again. Have a physician determine if condition of patient will permit evacuation of stomach. DO NOT give anything by mouth to an unconscious person. Call a physician immediately.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for 15 minutes. Get medical attention if irritation occurs.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633). FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

#### **PRECAUTIONARY STATEMENTS**

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

HARMFUL IF SWALLOWED. Avoid contact with skin, eyes, and clothing. Avoid contamination of feed or food.

#### **Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart. Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot we'er. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### **User Safety Recommendations**

Users should

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves perore removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

#### **READ ENTIRE LABEL BEFORE USING**

Protect from temperatures below 20°F.

Product crystallizes at lower temperatures.

Warm or store at higher temperatures and mix to redissolve crystals and assure uniformity before use.

Ro-Neet Booklet 8/17/93 1:51 PM Page 25 (PANTONE 470 CV plat

Mady in U.S.A

## ZENECA Ag Pro

ZENSCA Inc. Wilmington, Delaware 1969

BEST A