

ACCEPTED

JUN 14 1989

Under _____, and
Pesticide Act, as amended, for the
pesticide registered under
EPA Reg. No. **10182-180**

10-G

granular insecticide

...soil insecticide for pest control in a variety of crops

Active Ingredient:	
O-Ethyl S-phenyl ethylphosphonodithioate*	10 0%
Inert Ingredients	90 0%
Contains 10 pounds active ingredient per 100 pounds	100 0%

*DYFONATE

Keep Out of Reach of Children

WARNING AVISO

PRECAUCION AL USUARIO.
Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada completamente.

Statement of Practical Treatment FOR PESTICIDE EMERGENCY

CALL A PHYSICIAN IMMEDIATELY if a known exposure occurs, or if poisoning is suspected; do not wait for symptoms to develop. Immediately start the procedures given below and simultaneously contact a Poison Center, physician or the nearest hospital.

Note: Be sure to advise the person contacted that the compound is a cholinesterase inhibitor. Describe the type of exposure, the victim's symptoms, and follow the advice given.

POISON SYMPTOMS - DYFONATE is a highly toxic cholinesterase inhibitor. Symptoms of poisoning include headache, blurred vision, weakness, nausea, discomfort in the chest, abdominal cramps, vomiting, diarrhea, salivation, sweating and profuse pupils.

ALWAYS CALL A PHYSICIAN IN CASE OF SUSPECTED POISONING. ATROPINE BY INJECTION IS ANTIDOTAL.

If swallowed, immediately give several glasses of water and induce vomiting by placing a finger on the back of the victim's tongue. Give fluids until vomiting is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If on skin contact, flush with large amounts of running water for 15 minutes. Remove contaminated clothing and shoes. Get medical attention. Wash clothing before re-use.

For eye contact, immediately flush eyes with large amounts of running water for at least 15 minutes. Hold eyelids apart. Continue rinsing of the entire surface of the eye and lids with water. Get medical attention immediately.

If inhaled, remove to fresh air. If not breathing, clear victim's airway and start mouth-to-mouth artificial respiration.

If breathing is difficult, give oxygen, preferably with a physician's advice. Get medical attention immediately.

INFORMATION FOR MEDICAL PERSONNEL

Initial measures include removal of secretions, maintenance of a patent airway, and if necessary, artificial respiration. When cyanosis is relieved, atropine by injection is antidotal and may be administered in large therapeutic doses, repeated as necessary to the point of tolerance. If symptoms warrant further treatment, 2-PAM (Protopam Chloride) has shown utility as adjunctive therapy. Never use morphine. Continued absorption of the poison may occur, resulting in later relapse after initial improvement; very close supervision of the patient is indicated for at least 48 to 72 hours.

Do not transport or store Dyfonate 10-G with food or feed intended for human or animal consumption.

IN CASE OF SPILL - ISOLATE the spill. HOLD the package, other cargo and vehicles involved. IMMEDIATELY TELEPHONE (24 hour service for emergency assistance) 800-424-8300.

FOR 24 HOUR EMERGENCY MEDICAL ASSISTANCE, CALL ICI AMERICAS MEDICAL EMERGENCY INFORMATION CENTER (800) 321-8633 OR CHEMTREC (800) 424-8300.


For other product information, contact your local ICI Americas, Inc. distributor or see Product Safety Information Sheet.

READ ALL LABEL DIRECTIONS BEFORE USING

Keep Out of Reach of Children

WARNING

May Be Fatal If Swallowed



See Back Panel For Additional Precautionary Statements

EPA Reg. No. 10182-180
EPA Est. 476-180-1

50 LB. NET WT. 22.7 kg

A-688

AVOID CONTACT WITH SKIN. DO NOT GET IN THE EYES, ON SKIN, OR ON CLOTHING. AVOID BREATHING OF DUST.

Wear rubber gloves, shoes, protective clothing and goggles when handling or applying.

Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing. In case of contact, immediately remove contaminated clothing and wash thoroughly with soap and water.

Do not transport or store DYFONATE 10 G with any food or feed intended for human or animal consumption.

Not for use or storage in or around the home.

Environmental Hazards

This pesticide is toxic to fish and wildlife. Birds feeding in treated areas may be killed. Remove or cover granules if spills occur. Runoff from treated areas may be hazardous to fish in neighboring areas. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste. For guidance contact your regional office of the EPA. Apply this product only as specified on the label.

General Precautions

- Read all label directions before using.
- Do not apply on any crop or in any manner not specified on this label.
- Do not rotate with carrots.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Do not enter treated areas for 24 hours after application unless appropriate protective clothing is worn.
- Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.
- Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.
- Do not apply or allow to drift to areas occupied by beneficial insects or onto adjoining food, fiber or pasture crops. Do not contaminate food or feed.
- Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

When oral warnings are given, warnings shall be given in a language customarily understood by workers and must indicate the specific period of time that treated areas may not be entered without protective clothing. Oral warnings must also include appropriate first aid instructions in case of accidental exposure. These instructions are given in the STATEMENT OF PRACTICAL TREATMENT at the beginning of this label.

Written warnings must include the following information: "WARNING: Area treated with DYFONATE® Agricultural Insecticide on (date of application). Do not enter treated areas for 24 hours after application unless appropriate protective clothing is worn (or other reentry interval if your state has a more restrictive interval for this product). If accidental exposure occurs, follow the instructions below." (Written warnings must include the STATEMENT OF PRACTICAL TREATMENT given at the beginning of this label.)

is a violation of Federal laws.

ALABAMA

Luckdale, Limestone and Madison.

ARIZONA

Graham, Maricopa, Mohave, Pinal and Santa Cruz.

ARKANSAS

Benton and Polk.

CALIFORNIA

Butte, Colusa, Glenn, Imperial, Inyo, Kern, Los Angeles, Merced, Modoc, Orange, Riverside, Sacramento, San Bernardino, San Diego, Santa Barbara, Solano, Stanislaus, Sutter, Tehama, Ventura and Yuba.

FLORIDA

Alachua, Baker, Bradford, Brevard, Broward, Charlotte, Citrus, Clay, Collier, Columbia, Dade, DeSoto, Dixie, Duval, Flagler, Gadsden, Gilchrist, Glades, Hardee, Hendry, Hernando, Highlands, Hillsborough, Indian River, Jefferson, Lafayette, Lake, Lee, Leon, Levy, Madison, Manatee, Marion, Martin, Monroe, Nassau, Orange, Osceola, Ocala, Palm Beach, Pasco, Pinellas, Polk, Putnam, St. Johns, St. Lucie, Sarasota, Seminole, Sumter, Suwannee, Taylor, Union, Volusia and Wakulla.

GEORGIA

Braniff, Bryan, Bulloch, Burke, Camden, Candler, Chatham, Chatham, Effingham, Emanuel, Evans, Glascock, Glynn, Jefferson, Jenkins, Johnson, Liberty, Long, McIntosh, Pierce, Richmond, Screven, Ware, Washington and Wayne.

KANSAS

Clark, Comanche, Meade and Stafford.

MISSISSIPPI

Clayborne, Copah and Hinds.

MISSOURI

Barry, Benton, Camden, Christian, Dallas, Greene, Hickory, Jasper, Lawrence, Miller, Newton, Osage, Polk, St. Clair, Stone and Webster.

MONTANA

Garfield, McCone, Sheridan and Valley.

NEBRASKA

Boyd, Brown, Buffalo, Butler, Cass, Cedar, Colfax, Dawson, Dodge, Douglas, Hall, Hamilton, Holt, Howard, Kearney, Keya Paha, Knox, Nemack, Nance, Phelps, Platte, Polk, Rock, Sarpy and Saunders.

NEVADA

Clark.

NEW MEXICO

Chaves, DeBaca and Eddy.

NORTH DAKOTA

Benson, Bottineau, Burke, Burleigh, Divide, Dunn, Eddy, Emmons, Foster, Kidder, Logan, McHenry, McIntosh, McKenzie, McLean, Mercer, Morton, Mountrail, Nelson, Oliver, Pierce, Ramsey, Renville, Rolette, Sheridan, Sioux, Stutsman, Towner, Ward, Wells and Williams.

OHIO

Pickaway.

OKLAHOMA

Delaware, McCurtain and Pushmataha.

OREGON

Lake.

SOUTH CAROLINA

Aiken, Barnwell, Beaufort, Berkeley, Charleston, Clarendon, Dorchester, Georgetown, Hampton, Horry, Jasper and Marion.

SOUTH DAKOTA

Clay, Haakon, Hughes, Potter, Stanley, Sully, Union, Watworth, Yankton and Ziebach.

TENNESSEE

Lawrence and Wayne.

TEXAS

Arkansas, Austin, Bastrop, Burleson, Cameron, Colorado, Comal, Fort Bend, Goliad, Harris, Hays, Jeff Davis, Pecos, Reeves, Realguio and Victoria.

UTAH

Utah and Washington.

Place
Complete Directions For Use
Pamphlet
Here

Storage and Disposal

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Do not transport or store with any food or feed intended for human or animal consumption. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer and User, assuming the best of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Made in U.S.A. by
ICI Americas Inc.
Agricultural Products
Wilmington, DE 19887

WT-188

© ICI Americas Inc. 1988

Dyfonate[®] 10-G

granular insecticide

...soil insecticide for pest control
in a variety of crops

READ ALL LABEL DIRECTIONS BEFORE USING



ACCEPTED

JUN 14 1989

Under the provisions of the Federal
Pesticide Act, as amended, the
product is registered under
EPA Reg No. 10182-180

General precautions

- Read all directions before using
- Do not apply on any crop, or in any manner not specified on this label.
- Do not rotate with carrots

Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Reentry Statements

- Do not enter treated areas for 24 hours after application unless appropriate protective clothing is worn
- Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information
- Do not apply this product in such a manner, as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons

Do not apply or allow to drift to areas occupied by beneficial animals or onto adjoining food, fiber or pasture crops. Do not contaminate food or feed

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers

When oral warnings are given, warnings shall be given in a language customarily understood by the workers and must indicate the specific period of time that treated areas may not be entered without protective clothing. Oral warnings must also include appropriate first aid instructions in case of accidental exposure. These instructions are given in the STATEMENT OF PRACTICAL TREATMENT at the beginning of this label

Written warnings must include the following information, "WARNING, Area treated with DYFONATE[®] 10-G Granular Insecticide on (date of application). Do not enter treated areas for 24 hours after application unless appropriate clothing is worn (or other reentry interval if your state has a more restrictive interval for this product). If accidental exposure occurs, follow the instructions below" (Written warnings must include the STATEMENT OF PRACTICAL TREATMENT given at the beginning of this label)

Gauge settings and calibration instructions

NOTE: These settings should be used as starting points only. Continually check the amount of DYFONATE 10-G used against a known length of row or acreage and make further adjustments accordingly. Also, check calibration occasionally to make sure equipment wear, changing conditions, etc., have not caused a change in flow rate

GAUGE SETTINGS

DYFONATE 10-G 12 oz 1000 ft. of row and 22 oz 1000 ft. of row

Equipment	Speed in mph					
	4	6	8	10	12	14
Application Rate: Oz. per 1000 ft. of row	12	22	12	22	12	22
Gandy ball on bottom	26	34	31	41	35	48
Gandy snap-on bottom	26	34	31	41	35	47
Noble (round opening)	22	31	29	36	33	45
Noble (rectangular opening)	15	24	21	32	26	41
John Deere 1986 Maxi-Emerge II (adjustable knob-0-79)	33	50	45	58	37	66
John Deere 7000 Maxi-Emerge (Even) (0-10-20-30-40)	24	33	31	38	35	45
John Deere 7000 Maxi-Emerge (Odd) (5-15-25-35-45)	18	25	23	31	26	37
John Deere (metal box with 3 range settings and dial) Range Notch	2.13	3.6	2.32	3.28	3.9	-
International Harvester (one gauge per hopper) Gate Dial	2.7	4.5	3.8	6.4	4.8	8.3
International Harvester (two gauges per hopper) Gate Dial	2.0	4.8	4.1	6.7	5.1	8.6
Buffalo (with gauge)	2.1	3.1	2.7	4.0	3.3	-
Buffalo (fixed-roll 10-30—revolutions on hexagonal adjusting nut)	3 1/2	5 1/2	Uniform Flow at All Speeds			
Alfa Chalmers (old)	5	7	Uniform Flow at All Speeds			
Alfa Chalmers (new with Noble metering unit)	16	25	22	23	27	41
Microband - Case Industries	16**	27**	Uniform Flow at All Speeds			

**Unable to deliver quantity of DYFONATE required at speed indicated
**Number of teeth - Silver Rotor

EQUIVALENT LBS DYFONATE 10-G PER ACRE

Ounces of Dyfonate 10-G per 1000 ft. of row	Row Spacing in Inches							
	80	40	30	24	24	30	22	18
6.8	3.6	5.6	5.8	5.5	6.4	6.8	7.4	10.0
10.2	5.6	8.2	8.6	8.0	9.6	10.4	11.0	14.4
12	6.8	10.0	10.4	9.6	11.8	12.2	13.0	17.0
22	12.0	18.0	19.0	20.0	21.2	22.6	24.0	30.0
24	13.0	20.0	20.6	21.8	23.0	24.4	26.0	33.0
55.1	30.0	-	-	-	-	-	-	-
73.5	40.0	-	-	-	-	-	-	-

See reverse for Recommendations -



Made in U.S.A. by

Agricultural Products

ICI Americas Inc
Wilmington Delaware 19897


A-10MR

EPA Reg No 10182-180

DE 12-88

Recommendations

CROPS	APPLICATION METHOD	PESTS CONTROLLED	RATES OF DIFONATE 10-G	REMARKS	APPLICATION DIRECTIONS
ASPARAGUS	Up to 30 days prior to harvest	Symphytes	Broadcast: 30 to 100 pounds per acre	Recommended in California only. Reduce the rate proportionately for 1 and 2 treatments.	Broadcast. Apply DIFONATE 10-G and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment. Do not apply within 30 days of harvest.
BEANS, GARDEN (P. R.)	Prior to planting	Symphytes	Broadcast: 20 pounds per acre	For symphyte control, reduce the rate proportionately for band treatments.	Broadcast. Apply DIFONATE 10-G prior to planting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment.
		Wireworms	Broadcast: 40 pounds per acre		
		Seed Corn Maggot	Broadcast: 20 to 40 pounds per acre		
BEANS, LIMA	Prior to planting	Symphytes	Broadcast: 20 pounds per acre	Reduce the rate proportionately for band treatments.	Broadcast. Apply DIFONATE 10-G prior to planting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment.
COLE CROPS (BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND CAULIFLOWER)	Prior to planting or at planting	Symphytes	Broadcast: 20 pounds per acre	Reduce the rate proportionately for band treatments.	Broadcast. Apply DIFONATE 10-G prior to or at the time of seeding or transplanting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment.
CORN-FIELD (Dent seed grain silage; SWEET POPCORN)	Prior to planting or at planting	Northern, Southern and Western Corn Rootworm		May be used on conventional row or no-till conservation and no-till tillage systems.	Band. Apply DIFONATE 10-G in a 7 inch wide band using a row bander directly behind the seed furrow opener and in front of the press wheels or in a 7 inch wide band of rock chains or spring tines to insure incorporation. Under windy conditions the use of windshield is recommended to prevent drift. Some granules can fall into the open furrow without causing harm to the seed. However, to avoid stand reduction do not make "in furrow" applications of DIFONATE 10-G.
		Suppression of Cutworms			
		Suppression of White Grubs	Band: 12 oz per 1000 linear feet		
		Seedcorn Maggot Seedcorn Beetle	or Broadcast: 40 pounds per acre	DIFONATE 10-G is recommended for control of seedcorn beetle in the Central and Midwestern corn growing areas only.	
		Wireworms			
		Lesser Corn Rootworm Symphytes Maize Weevil	Band: 24 oz per 1000 linear feet Broadcast: 20 pounds per acre Broadcast: 40 pounds per acre	Band applications of DIFONATE 10-G will provide suppression of wireworms. Use the broadcast application for control.	
At Cultivation	Northern, Southern and Western Corn Rootworm	Band: 12 oz per 1000 linear feet	Apply DIFONATE 10-G when activity of rootworm larvae is first observed.	Band. Apply DIFONATE 10-G in a 7 inch wide band over the corn rows. Cover the treated bands with 2 to 3 inches of soil by making the application ahead of disc, harrow or cultivation equipment. Do not apply within 45 days of harvest, nor feed or graze livestock within 30 days of treatment.	
Over the Whorl	European Corn Borer, Southwestern Corn Borer		Band: 12 oz per 1000 linear feet	Consult your State Agricultural Experiment Station or extension service specialist for local treatment threshold and application timing information. Scouting for insects is strongly recommended.	Band or Broadcast. Apply DIFONATE 10-G over the corn rows so granules fall into the whorl. May be applied broadcast using air or ground equipment, or in a band directed over corn rows. Do not apply within 30 days of harvest, nor feed or graze livestock within 30 days of treatment.
		Reduction of Fall Armyworm Populations	Broadcast: 10 pounds per acre		
SWEET CORN (Green or Black Belt)	At Planting	Wireworms	Band: 22 oz per 1000 linear feet	Recommended in Florida only.	Band. Apply DIFONATE 10-G in a 2 inch wide band over the corn rows.
ONIONS, BULB	At Planting	Onion Maggot	In-Furrow: 6 oz per 100 linear feet	Use only in soils containing 10% or more organic matter.	In-Furrow. Apply DIFONATE 10-G into the furrow at time of planting.
PEANUTS (Southeastern U.S.)	At Pegging or up to 30 Days After Pegging Begins	Southern Corn Rootworm, Granulate Cutworm	Band: 22 oz per 1000 linear feet	For control of southern corn rootworm and granulate cutworm, apply DIFONATE 10-G at pegging.	Band. Apply DIFONATE 10-G in a 18 inch wide band.
		Lesser Corn Rootworm	or	For control of lesser corn rootworm, apply DIFONATE 10-G at or after pegging.	Broadcast. Apply DIFONATE 10-G at or after pegging.
		Suppression of White Mold (Sclerotium rot)	Broadcast: 40 pounds per acre	For heavy white mold infestations, suitable fungicide treatment is necessary.	



Recommendations

CROPS	APPLICATION METHOD	PESTS CONTROLLED	RATES OF DIFLONATE 10-G	REMARKS	APPLICATION DIRECTIONS
PEANUTS (Southwestern U.S.)	At Popping or up to 30 Days After Popping Begins	Southern Corn Rootworm	Band 22 oz per 1000 linear feet		Band Apply DIFLONATE 10-G in a 18 inch wide band
		Lesser Cornstalk Borer	Band 9 8 to 13 7 oz per 1000 linear feet		Band Apply DIFLONATE 10-G in a 14 inch wide band
		Burrowing Bug	Band 11 to 22 oz per 1000 linear feet	Recommended for control of burrowing bug in Texas only	Band Apply DIFLONATE 10-G in a 18 inch wide band
			Broadcast: 20 pounds per acre		Broadcast Apply DIFLONATE 10-G and lightly incorporate into the soil 1 to 2 inches of soil.
POTATOES (WASH)	Prior to Planting or at Planting	Wireworms	Subsurface Bands 20 oz per 1000 linear feet	The use of subsurface bands is recommended in the Pacific Northwest only. This application will control light to moderate infestations of wireworms. Use the 40 pound per acre broadcast application for heavy infestations of wireworms.	Subsurface Bands Apply DIFLONATE 10-G in subsurface bands 3 to 4 inches on both sides of and 2 inches below the seed pieces
		Symphyten	Broadcast: 40 pounds per acre Broadcast: 20 pounds per acre		Broadcast Apply DIFLONATE 10-G prior to or at time of planting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment
RADISH	Prior to planting or at planting	Cabbage Maggot	Broadcast: 20 pounds per acre		Broadcast Apply DIFLONATE 10-G prior to or at time of planting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment
		Symphyten		For symphyten control reduce the rate proportionately for band treatments	
SORGHUM (Wash)	At Planting	Northern, Southern and Western Corn Rootworm	Band 12 oz per 1000 linear feet		Band Apply DIFLONATE 10-G in a 7 inch wide band using a row band applicator directly behind the seed furrow opener and in front of the press wheels or in a 7 inch wide band directly behind the press wheels using drag chains or spring tines to insure incorporation. Under windy conditions the use of windshields is recommended to prevent drift. Some granules can fall into the seed furrow without causing harm to the seed. However to avoid stand reduction do not make in furrow applications of DIFLONATE 10-G
		Lesser Cornstalk Borer			Broadcast Apply DIFLONATE 10-G over the top using same equipment. Use only on emerged sorghum. Do not harvest nor feed or graze live stock within 14 days of application
	Over the Top	Corn Leaf Apter Greenbug	Broadcast: 75 pounds per acre	Recommended in Texas only. Live post-emergence use to 2 applications per crop	
SUGAR BEETS	Prior to planting or at planting	Sugar Beet Root Maggot	Band 6 8 to 10 5 oz per 1000 linear feet Subsurface Band 10 5 oz per 1000 linear feet	Recommended in Colorado and Northwest Sugar Beet growing areas only. To avoid stand reduction do not place DIFLONATE 10-G in direct contact with the seed	Band Apply DIFLONATE 10-G in a 7 inch wide band over the sugar beet rows. Incorporate lightly into the soil using suitable tillage equipment. Subsurface Band Apply DIFLONATE 10-G 2 inches from the seed furrow and 2 inches below the soil surface on the emerged side of the sugar beet rows. Recommended in the Big Horn Basin of Wyoming only.
		Symphyten	Broadcast: 20 pounds per acre	For symphyten control reduce the rate proportionately for band treatments	Broadcast Apply DIFLONATE 10-G prior to planting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment
		Wireworms	Broadcast: 40 pounds per acre		
SUGARCANE	At Planting	Wireworms	Band 55 1 to 73 5 oz per 1000 linear feet	A use rate of less than 40 pounds is only recommended in Florida	Band Apply DIFLONATE 10-G in a 14 inch band over the tops of the sugar cane seed pieces immediately ahead of covering discs
TABLE BEETS	Prior to Planting	Cabbage Maggot Symphyten	Broadcast: 20 pounds per acre	For symphyten control reduce the rate proportionately for band treatments	Broadcast Apply DIFLONATE 10-G prior to planting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment
TOBACCO	Prior to bedding	Wireworms	Broadcast: 10 to 20 pounds per acre		Broadcast Apply DIFLONATE 10-G prior to bedding and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment
TOMATOES	Prior to planting	Symphyten	Broadcast: 20 pounds per acre	Reduce the rate proportionately for band treatments	Broadcast Apply DIFLONATE 10-G prior to planting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment

NOTE: For all crops where APPLICATION METHOD is "Prior to planting" results are best when application is made as close as possible to planting.