

PM 11 21 1991



DYFONATE® II 15-G Granular Insecticide

Soil insecticide for Pest Control In A Variety of Crops

ACTIVE INGREDIENT:	
O-Ethyl S-phenyl ethylphosphonodithioate*	15.0%
INERT INGREDIENTS:	85.0%
Total	100.0%

Contains 15 pounds active ingredient per 100 pounds
*DYFONATE

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO
PRECAUCION AL USUARIO: Si usted no le ingles, no use este producto hasta que la etiqueta le haya sido explicado ampliamente.

See back panel for additional Precautionary Statements

EPA Reg. No. 10182-187
EPA Est. No.

50 Lbs. Net Wt. 22.7 Kg.

Made in U.S.A. by
ICI Agricultural Products
A business unit of ICI Americas Inc.
Wilmington, Delaware 19897

ACCEPTED
APR 29 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10182-187

BEST AVAILABLE COPY

STATEMENT OF PRACTICAL TREATMENT

FOR PESTICIDE EMERGENCY

CALL PHYSICIAN IMMEDIATELY. If a known exposure occurs, or if poisoning is suspected, do not wait for symptoms to develop. Immediately contact a poison Control Center, a physician, or the nearest hospital.

NOTE: Be sure to advise the person contacted that the compound is a cholinesterase inhibitor. Describe the type of exposure, the victim's symptoms, and follow the advice given.

POISON SYMPTOMS: DYFONATE II 15-G is a highly toxic cholinesterase inhibitor.

Symptoms of poisoning include headache, blurred vision, weakness, nausea, discomfort in the chest, abdominal cramps, vomiting, diarrhea, salivation, sweating and pinpoint pupils.

ALWAYS CALL A PHYSICIAN IN CASE OF SUSPECTED POISONING.

ATROPINE BY INJECTION IS ANTIDOTAL.

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by placing a finger on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Flush with large amounts of running water for 15 minutes.

Remove contaminated clothing and shoes. Get medical attention. Wash clothing before reuse.

IF IN EYES: Immediately flush eyes with large amounts of running water for at least 15 minutes. Hold the eyelids apart to ensure rinsing of the entire surface of the eye and lids with water. Get medical attention immediately.

IF INHALED: Remove to fresh air. If not breathing, clear victim's airway, start mouth-to-mouth artificial respiration. **IF BREATHING IS DIFFICULT,** give oxygen, preferably with a physician's advice. Get medical attention immediately.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

INFORMATION FOR MEDICAL PERSONNEL: Initial measures include removal of secretions, maintenance of patient airway, and if necessary, artificial respiration. When cyanosis is relieved, atropine by injection is antidotal and may be administered in large therapeutic doses, repeated as necessary to the point of tolerance. If symptoms warrant further treatment, 2-PAM (Protopam Chloride) has shown utility as adjunctive therapy. Never use morphine. Continued absorption of the poison may occur, resulting in fatal relapse after initial improvement; very close supervision of the patient is indicated for at least 48 to 72 hours.

BEST AVAILABLE COPY

DO NOT TRANSPORT OR STORE DYFONATE II 15-G WITH FOOD OR FEED INTENDED FOR HUMAN OR ANIMAL CONSUMPTION.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Back Panel

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. RAPIDLY ABSORBED THROUGH SKIN. DO NOT GET IN THE EYES, ON SKIN, OR ON CLOTHING. AVOID BREATHING OF DUST.

Wear rubber gloves, shoes, protective clothing and goggles when handling or applying. Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing. In case of contact, immediately remove contaminated clothing and wash thoroughly with soap and water.

Do not transport or store DYFONATE II 15-G with any food or feed intended for human or animal consumption.

Not for use or storage in or around the home.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Birds feeding in treated areas may be killed. Remove or cover granules if spills occur. Runoff from treated areas may be hazardous to fish in neighboring areas. Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters.

For guidance contact your regional office of the EPA. Apply this product only as specified on the label.

In case of significant spill:

HOLD THIS PACKAGE, OTHER CARGO AND VEHICLES INVOLVED. IMMEDIATELY TELEPHONE (24-HOUR SERVICE) FOR EMERGENCY ASSISTANCE (800) 424-9300.

BEST AVAILABLE COPY

GENERAL PRECAUTIONS

- Read all directions before using.
- Do not apply on any crop, or in any manner not specified on this label.
- Do not rotate with carrots.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to Label Booklet for Complete Use Directions

REENTRY STATEMENTS

- Do not enter treated areas for 24 hours after application unless appropriate protective clothing is worn.
- Foliar Applications (over the top) for Corn and Sorghum: Do not allow worker reentry into treated fields within 96 hours of application unless appropriate protective clothing is worn. Protective clothing means, at least, a hat or other suitable head covering, gloves, a long-sleeved shirt and long-legged trousers or coverall type garment (all closely woven fabric covering the body, including the arms and legs), shoes and socks.
- Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.
- Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.
- Do not apply or allow drift to areas occupied by beneficial animals or onto adjoining food, fiber or pasture crops. Do not contaminate food or feed.
- Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

When oral warnings are given, warnings shall be given in a language customarily understood by workers and must indicate the specific period of time and treated areas may not be entered without protective clothing. Oral warnings must also include appropriate first aid instructions in case of accidental exposure. These instructions are given in the STATEMENT OF PRACTICAL TREATMENT at the beginning of this label.

BEST AVAILABLE COPY

Written warnings must include the following information, "WARNING Area treated with DYFONATE II 15-G Agricultural Insecticide on (date of application). Do not enter treated areas for 24 hours (96 hours for foliar applications in corn and sorghum) unless appropriate protective clothing is worn (or other reentry interval if your state has a more restrictive interval for this product). ~~If accidental exposure occurs, follow the instructions below.~~" (Written warnings must include the STATEMENT OF PRACTICAL TREATMENT given at the beginning of this label.)

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Do not transport or store with any food or feed intended for human or animal consumption. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

BEST AVAILABLE COPY

**DIRECTIONS FOR USE AND
CONDITIONS OF SALE AND WARRANTY**

IMPORTANT: Read the Entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIABILITY of Seller will be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

DYF187C.RSB

BEST AVAILABLE COPY



DYFONATE® II 15-G Granular Insecticide

Soil Insecticide For Pest Control In a Variety of Crops

READ ALL LABEL DIRECTIONS BEFORE USING

GENERAL PRECAUTIONS

- Read all directions before using.
- Do not apply on any crop, or in any manner not specified on this label.
- Do not rotate with carrots.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application of a sulfonylurea herbicide such as Beacon* and Accent following application of a soil insecticide such as DYFONATE II 15-G may result in crop injury. Follow the Directions of Use and Precautions contained on the sulfonylurea herbicide label.**

REENTRY STATEMENTS

- Do not enter treated areas for 24 hours after application unless appropriate protective clothing is worn.
- Foliar Applications (over the top) for Corn and Sorghum: Do not allow reentry into treated fields within 96 hours of application unless appropriate protective clothing is worn. Protective clothing means, at least, a hat or other suitable head covering, gloves, a long-sleeved shirt and long-legged trousers or coverall type garment (all closely woven fabric covering the body, including the arms and legs), shoes and socks.
- Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.
- Do not apply this product in such a manner as to directly or through drift expose workers or other person. The area being treated must be vacated by unprotected persons.

* Beacon is a registered trademark of Ciba-Geigy Corporation

** Accent is a registered trademark of DuPont Company

BEST AVAILABLE COPY

- Do not apply or allow to drift to areas occupied by beneficial animals or onto adjoining food, fiber or pasture crops. Do not contaminate food or feed.
- Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

When oral warnings are given, warnings shall be given in a language customarily understood by the workers and must indicate the specific period of time that treated areas may not be entered without protective clothing. Oral warnings must also include appropriate first aid instructions in case of accidental exposure. These instructions are given in the STATEMENT OF PRACTICAL TREATMENT at the beginning of this label.

Written warnings must include the following information, "WARNING, Area treated with DYFONATE II 15-G Agricultural Insecticide on (date of application). Do not enter treated areas for 24 hours (96 hours after foliar applications in corn and sorghum) unless appropriate clothing is worn (or other reentry interval if your state has a more restrictive interval for this product). ~~If accidental exposure occurs, follow the instructions below.~~" (Written warnings must include the STATEMENT OF PRACTICAL TREATMENT given at the beginning of this label.)

GAUGE SETTINGS AND CALIBRATION INSTRUCTIONS

NOTE: These settings should be used as starting points only. Continually check the amount of DYFONATE 11 15-G used against a known length of row or acreage and make further adjustments accordingly. Also, check calibration occasionally to make sure equipment wear, changing conditions, etc., have not caused a change in flow rate.

BEST AVAILABLE COPY

GAUGE SETTINGS DYFONATE II 15-G 8 OZ/1000 FT. OF ROW AND 16 OZ/1000 FT. OF ROW						
EQUIPMENT	SPEED IN MPH					
	4		6		8	
APPLICATION RATE: OUNCES PER 1000 FEET OF ROW	8	14.7	8	14.7	8	14.7
Gandy bolt on bottom	27	25	22	23	22	21
Gandy snap on bottom	27	25	23	24	27	22
Noble (round opening)	21	23	20	21	24	1
Noble (rectangular opening)	15	25	21	25	21	23
John Deere 1986 Maxi-Emerge II (adjustable knob-0-79)	29	29	25	25	20	20
John Deere 7000 Maxi-Emerge (Even) (0-10-20-30-40)	25	24	22	21	26	1
John Deere 7000 Maxi-Emerge (Odd) (5-15-25-35-45)	19	22	22	24	22	21
John Deere (metal box with 3 range settings and dial) Range/Notch	2/10	2/26	2/22	2/4	2/30	2/12
International Harvester (one gauge per hopper) Gate/Dial	2/6	2/6	2/7	2/0	2/9	2/3
International Harvester (two gauges per hopper) Gate/Dial	2/1	2/2	2/4	2/8	2/7	2/1
Buffalo (with gauge)	20	20	22	25	22	23
Buffalo (fluted-roll feed revolutions on hexagonal adjusting nut)	1	1	2	2	2	2
Allis Chalmers (old)	1	1	2	2	2	2
Allis Chalmers (new, with Noble metering unit)	18	25	22	23	25	22
Microband (Canaan Industries)		2/2	2/2	2/2	2/2	2/2

Unable to deliver quantity of DYFONATE II 15-G required at speed indicated.
 Number of teeth on sprockets. For additional rates/settings or for Double-Box units
 contact Canaan Industries at 1-800-633-7500.
 Uniform Flow at All Speeds

BEST AVAILABLE COPY

REQUIRED POUNDS OF DYFONATE II 15-G PER ACRE									
Ounces of DYFONATE II 15-G per 1000 ft. of row	ROW SPACING IN INCHES								
	60	40	38	36	34	32	30	22	18
8.5	25	8.7	8.9	9.1	9.3	9.5	9.7	9.9	10.2
8.8	27	9.6	9.8	10.1	10.3	10.5	10.7	10.9	11.2
9.0	28	9.8	10.0	10.3	10.5	10.7	10.9	11.1	11.4
14.7	8.0	12.0	12.6	13.3	14.1	15.0	16.0	17.9	26.7
36.0	19.6	29.4	31.0	32.7	34.6	36.6	39.2	43.5	65.3
48.0	26.1	39.2	41.3	43.6	46.1	49.0	52.3	58.3	87.1
72.0	39.2	58.8	61.9	65.3	69.2	73.5	78.4	106.9	130.7

BEST AVAILABLE COPY

RECOMMENDATIONS					
CROPS	APPLICATION METHOD	PESTS CONTROLLED	RATES OF DYFONATE II 15-G	REMARKS	APPL
COLE CROPS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower)	Immediately prior to planting or at planting	Cabbage Maggot	Broadcast: 27 pounds per acre.	Not recommended in the Northwestern U.S.	Broa II 1 time tran inco to a inch t11

RECOMMENDATIONS					
CROPS	APPLICATION METHOD	PESTS CONTROLLED	RATES OF DYFONATE II 15-G	REMARKS	APPLICATION DIRECTIONS
CORN-FIELD, (Dent, Seed, Grain, Silage) SWEET POPCORN	Immediately prior to planting or at planting	Northern, Southern, & Western Corn Rootworm Black Cutworm	Band: 8 oz. per 1000 linear feet. or Broadcast: 27 pounds per acre	May be used on conventional, minimum or conservation and no-till tillage systems. With heavy infestations of cutworms, a sufficient number of larvae may survive the DYFONATE treatment to cause damage to the corn. If this occurs, a rescue treatment with AMBUSH Insecticide or another insecticide registered for this use can be made to control these larvae. DYFONATE II 15-G is recommended for control of seed corn beetle in the central and midwestern corn growing areas only.	Band: Apply DYFONATE II 15-G in a 7 inch wide band using a row bander directly behind the seed furrow opener and in front of the press wheels, or in a 7 inch wide band directly behind the press wheels using drag chains or spring tines to insure incorporation. Under windy conditions the use of windshields is recommended to prevent band drift. Some granules can fall into the open furrow without causing harm to the seed. However, to avoid stand reduction, do not make "in-furrow" applications of DYFONATE II 15-G. Broadcast: Apply DYFONATE II 15-G prior to planting and incorporate into the soil to a depth of 2 to 3 inches using suitable tillage equipment.
		Suppression of White Grubs Seedcorn Maggot Seedcorn Beetle <u>Wireworms</u> Lesser Cornstalk Borer <u>Symphylans only</u>	Band: 14.7 oz per 1000 linear feet. Broadcast: 13.4 pounds per acre.	Band applications of DYFONATE II 15-G will provide suppression of wireworms. Use the broadcast application for control.	

RECOMMENDATIONS					
CROPS	APPLICATION METHOD	PESTS CONTROLLED	RATES OF DYFONATE II 15-G	REMARKS	APPLICATION DIRECTIONS
	At Cultivation	Maize Billbug	Broadcast: 27 pounds per acre.	DYFONATE II 15-G is recommended for control of maize billbug in the Southeastern U.S. only.	
	Over the Whorl	Northern, Southern, and Western Corn Rootworm	Band: 8 oz. per 1000 linear feet.	Apply DYFONATE II 15-G when activity of rootworm larvae is first observed.	Band: Apply DYFONATE II 15-G in a 7 inch wide band over the corn rows. Cover the treated bands with 2 to 3 inches of soil by making the application ahead of disc hillers or cultivation equipment. Do not apply within 45 days of harvest nor feed or graze livestock within 30 days of treatment.
		European Corn Borer, Southwestern Corn Borer	Band: 8 oz. per 1000 linear feet.	Consult your State Agricultural Experiment Station or extension service specialist for local treatment threshold and application timing information. Scouting for insect damage is strongly recommended.	
		Reduction of Fall Armyworm Populations	or Broadcast: 6.75 pounds per acre.		Band or Broadcast: Apply DYFONATE II 15-G over the corn rows so granules fall into the whorl. May be applied broadcast, using air or ground equipment, or, in a band directed over corn rows. Do not apply within 30 days of harvest nor feed or graze livestock within 30 days of treatment.
SWEET CORN (Grown on Muck Soils)	At Planting	Wireworms	Band: 14.7 oz. per 1000 linear feet.	Recommended in Florida only.	Band: Apply DYFONATE II 15-G in a 2 inch wide band over the corn rows.

BEST AVAILABLE COPY

RECOMMENDATIONS					
CROPS	APPLICATION METHOD	PESTS CONTROLLED	RATES OF DYFONATE II 15-G	REMARKS	APPLICATION DIRECTIONS
PEANUTS	At Early Flowering or up to 30 days After Pegging Begins	Southern Corn Rootworm, Granulate Cutworm Lesser Cornstalk Borer suppression of White Mold (Sclerotinia rolfsii)	Band: 14.7 oz. per 1000 linear feet. or Broadcast: 26.67 pounds per acre	Recommended for the Southeastern U.S. only. For control of southern corn rootworm and granulate cutworm apply DYFONATE II 15-G at early flowering or pegging. For control of lesser cornstalk borer, apply DYFONATE II 15-G at first signs of borer infestation after pegging begins. For heavy white mold infestations a suitable fungicide is necessary.	Band: Apply DYFONATE II 15-G in an 18 inch wide band. May be used up to 30 days after pegging begins. Broadcast: Apply DYFONATE II 15-G by air or ground equipment.
SORGHUM (milo)	Over the Top	Corn Leaf Aphid Greenbug	Broadcast: 5 pounds per acre.	Recommended in Texas only. Limit post-emergency use to 2 applications per crop.	Broadcast: Apply DYFONATE II 15-G over the top using aerial equipment. Use only on irrigated sorghum. Do not harvest nor feed or graze livestock within 14 days of application.
SUGAR BEETS	At Planting	Sugar Beet Root Maggot	Band: 4.5 to 6.8 oz. per 1000 linear feet.	To avoid reduction, do not place DYFONATE II 15-G in direct contact with the seed.	Band: Apply DYFONATE II 15-G in a 7 inch wide band over the sugar beet rows. Incorporate lightly into the soil using suitable tillage equipment.

RECOMMENDATIONS					
CROPS	APPLICATION METHOD	PESTS CONTROLLED	RATES OF DYFONATE II 15-G	REMARKS	APPLICATION DIRECTIONS
SUGARCANE	Post-Emergence	White Grubs	Band: 48 to 72 oz. per 1000 linear feet.	Recommended in the Southwestern U.S. only. Consult your State Agricultural Experiment Station or extension service specialist for local treatment threshold and application timing information.	Band: Apply DYFONATE II 15-G over the top of the cane row and lightly incorporate as close to the plants as possible. For best results, irrigate the application.
	At Planting	Wireworms	Band: 36 to 48 oz. per 1000 linear feet.	Use only 48 oz. rate outside of Florida.	Apply in 14 inch band on cane rows 60 inches apart. Apply over the tops of sugarcane seed pieces immediately ahead of covering discs.

EPA Reg. No. 10182-187

Made in U.S.A. by
 ICI Agricultural Products
 A business unit of ICI Americas Inc.
 Wilmington, Delaware 19897

DYF187B.RSD

BEST AVAILABLE COPY