

PM14 10182-195

10/16

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

5407672 $\frac{300}{18}$

DEC 4 1991

Mr. Andrew A. Davidson
ICI Americas Inc.
Agricultural Products Division
Delaware Corporate Center
Wilmington, DE 19897

Dear Mr. Davidson:

Subject: Dyfonate-Tillam 1-4E Insecticide-Herbicide
EPA Registration no. 10182-195
Your Application for Pesticide Amendment
Dated November 27, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the revised labeling:
 - a. As discussed on the phone, the product name has been changed and according to our files the name is now **DYFONATE-TILLAM 1-4E Insecticide-Herbicide**.
 - b. Also as discussed, under the section **DYFONATE-TILLAM 1-4E IMPREGNATION ON DRY BULK FERTILIZERS** clarify the use of the word "approved" by stating "ICI approved" or "approved by ICI".
2. Submit a copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Robert A. Forrest
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (H7504C)

~~MADISON~~ DISK 3 12/4/91

CONCURRENCES							
SYMBOL	enclosure						
SURNAME							
DATE							

208/16

DYFONATE-TILLAM 1-4E
Insecticide-Herbicide

Emulsifiable Liquid for the Control of Weeds and Insects in Tobacco

ACTIVE INGREDIENTS:

O-Ethyl S-phenyl ethylphosphonodithioate	12.6%
S-propyl butylethylthiocarbamate**	50.4%
INERT INGREDIENTS***:	37.0%
Total	100.0%

Contains 1 lb. fonofos* per gallon and 4 lbs. pebulate** per gallon.

***Contains xylene-range aromatic solvent.

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicado ampliamente.

EPA Reg. No. 10182-195
EPA Est. No. 10182-

Net Gals. _____

Made in U.S.A. by
ICI Agricultural Products
A business unit of ICI Americas Inc.
Wilmington, Delaware 19897

ACCEPTED
with COMMENTS
in EPA Letter Dated:

DEC 4 1991

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

10182-195

STATEMENT OF PRACTICAL TREATMENT

FOR PESTICIDE EMERGENCY

CALL PHYSICIAN IMMEDIATELY: If a known exposure occurs, or if poisoning is suspected, do not wait for symptoms to develop. Immediately contact a Poison Control Center, a physician, or the nearest hospital.

Note: Be sure to advise the person contacted that the compound is a cholinesterase inhibitor. Describe the type of exposure, the victim's symptoms, and follow the advice given.

POISON SYMPTOMS: DYFONATE® insecticide is a highly toxic cholinesterase inhibitor. Symptoms of poisoning include headache, blurred vision, weakness, nausea, discomfort in the chest, abdominal cramps, vomiting, diarrhea, salivation, sweating, and pinpoint pupils.

**ALWAYS CALL A PHYSICIAN IN CASE OF SUSPECTED POISONING.
ATROPINE BY INJECTION IS ANTIDOTAL.**

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by placing a finger on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Flush with large amounts of running water for 15 minutes. Remove contaminated clothing and shoes. Get medical attention. Wash clothing before reuse.

IF IN EYES: Immediately flush eyes with large amounts of running water for at least 15 minutes. Hold the eyelids apart to ensure rinsing of the entire surface of the eye and lids with water. Get medical attention immediately!

IF INHALED: Remove to fresh air. If not breathing, clear victim's airway and start mouth-to-mouth artificial respiration. Get medical attention.

IF BREATHING IS DIFFICULT, give oxygen, preferably with a physician's advice. Get medical attention immediately.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

INFORMATION FOR MEDICAL PERSONNEL: Initial measures include removal of secretions, maintenance of a patent airway, and if necessary, artificial respiration. When cyanosis is relieved, atropine by injection is antidotal and may be administered in large therapeutic doses, repeated as necessary to the point of tolerance. If symptoms warrant further treatment, 2-PAM (Protopam Chloride) has shown utility as adjunctive therapy. Never use morphine. Continued absorption of the poison may occur, resulting in fatal relapse after initial improvement; very close supervision of the patient is indicated for at least 48 to 72 hours.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. RAPIDLY ABSORBED THROUGH THE SKIN. DO NOT GET IN THE EYES, ON SKIN, OR ON CLOTHING. AVOID BREATHING OF SPRAY MIST.

Use MSHA/NIOSH approved respirator for pesticides.

Wear rubber gloves, shoes, protective clothing, and goggles when handling or applying. Protective clothing means, at least, a hat or other suitable head covering, gloves, a long-sleeved shirt and long-legged trousers or coverall type garment (all closely woven fabric covering the body, including the arms and legs), shoes and socks.

Wash hands, arms, and face thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing. In case of contact, immediately remove contaminated clothing and wash thoroughly with soap and water.

Do not transport or store DYFONATE®-TILLAM® 1-4E insecticide-herbicide with any food or feed intended for human or animal consumption.

Not for use or storage in or around the home.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Birds feeding in treated areas may be killed. Remove or cover if spills occur. Runoff from treated areas may be hazardous to fish in neighboring areas. ~~Keep out of lakes, ponds, or streams.~~ Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. For guidance contact your regional office of the EPA. Apply this product only as specified on the label.

In case of spill, isolate the spill.

Hold this package, other cargo and vehicles involved.

Immediately telephone (24-hour service) for emergency assistance CHEMTREC 1-800-424-9300.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

GENERAL PRECAUTIONS

- Read all directions before using.
- Do not apply to any crop or in any manner not specified on this label.

50416

- Do not rotate to carrots.
- Applied according to directions and under normal growing conditions, this product will not harm the treated crop. During germination and early growth, extended periods of unusually cold and wet or hot and dry weather, insect, nematode, or plant disease attack, the use of certain soil applied systemic insecticides, highly saline or alkaline soil conditions, improperly placed fertilizers or soil insecticides may create abnormal conditions that weaken crop seedlings. DYFONATE-TILLAM 1-4E used under these abnormal conditions could result in crop injury.
- Do not tank mix with insecticides, fungicides or fluid fertilizers unless their compatibility has been tested and proven to be satisfactory prior to use or except as specified on this label.
- Do not store near seeds or fertilizers.

50416

60416

SOIL INSECTS CONTROLLED¹

COMMON NAME	SCIENTIFIC NAME
Black cutworm	<i>Agrotis ipsilon</i>
Com wireworm	<i>Melanotus communis</i>
Southern potato wireworm	<i>Conoderus falli</i>
Tobacco wireworm	<i>Conoderus vespertinus</i>
Tobacco flea beetle larvae ²	<i>Epitrix hirtipennis</i>

- ¹ For control under severe insect infestations, tank mix 1 quart/acre of DYFONATE 4-EC.
- ² For suppression only; a transplant or setter water treatment of a suitable insecticide is necessary for control.

WEEDS CONTROLLED

ANNUAL GRASSES	
COMMON NAME	SCIENTIFIC NAME
Barnyardgrass, Watergrass	<i>Echinochloa crus-galli</i>
Bermudagrass, seedling	<i>Cynodon dactylon</i>
Crabgrass	<i>Digitaria spp.</i>
Foxtail (Giant, Green, Yellow)	<i>Setaria spp.</i>
Goosegrass	<i>Eleusine indica</i>
Millet	<i>Setaria spp.</i>
Narrowleaf signalgrass, Alexandergrass	<i>Brachiaria piligera</i>
Wild oats	<i>Avena fatua</i>

70116

The above following broadleaf weeds will be controlled only if DYFONATE-TILLAM 1-4E is applied when conditions are favorable for germination.

ANNUAL BROADLEAF WEEDS	
COMMON NAME	SCIENTIFIC NAME
Blackeyed susan, Sweet suzy	<i>Rudbeckia hirta</i>
Common lambsquarters	<i>Chenopodium album</i>
Common purslane	<i>Portulaca oleracea</i>
Deadnettle, Henbit	<i>Lamium amplexicaule</i>
Florida pusley	<i>Richardia scabra</i>
Hairy nightshade, Groundcherry ¹	<i>Solanum villosum</i>
Nettleleaf goosefoot	<i>Chenopodium murale</i>
Prostrate pigweed	<i>Amaranthus graecizans</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Shepherd's-purse	<i>Capsella bursa-pastoris</i>

¹Suppression or partial control only.

PERENNIAL WEEDS	
COMMON NAME	SCIENTIFIC NAME
Yellow nutsedge	<i>Cyperus esculentus</i>
Purple nutsedge	<i>Cyperus rotundus</i>

Existing stands of nutsedge must be turned under and chopped up thoroughly before treatment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application should be made by ground power equipment and by trained personnel only. Do not apply with hand equipment.

Do not apply this product with any type of irrigation equipment.

REENTRY STATEMENTS

- Do not enter treated areas for 24 hours after application unless appropriate protective clothing is worn.
- Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.
- Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.
- Do not apply or allow to drift to areas occupied by beneficial animals or onto adjoining food, fiber or pasture crops. Do not contaminate food or feed.
- Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

When oral warnings are given, warnings shall be given in a language customarily understood by workers and must indicate the specific period of time that treated areas may not be entered without protective clothing. Oral warnings must also include appropriate first aid instructions in case of accidental exposure. These instructions are given in the **STATEMENT OF PRACTICAL TREATMENT section**.

Written warnings must include the following information, ***WARNING. Area treated with DYFONATE-TILLAM 1-4E Agricultural Insecticide on (date of application). Do not enter treated areas for 24 hours after application unless appropriate protective clothing is worn (or other reentry interval if your state has a more restrictive interval for this product). If accidental exposure occurs, follow the instructions below.*** (Written warnings must include the **STATEMENT OF PRACTICAL TREATMENT section**.)

This product contains a selective herbicide which is mixed (incorporated) into the soil for control of weeds listed on this label. It controls weeds by interfering with normal seed germination and seedling development. It does not control established weeds. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

USE RATE

Apply DYFONATE-TILLAM 1-4E at the rate of 1 gallon per acre.

BEST AVAILABLE COPY

APPLICATION DIRECTIONS:

- The soil should be well-worked and dry enough to permit good soil mixing (incorporation).
- Pour the recommended amount of product into the spray tank during the filling operations.
- Apply in 20 to 50 gallons of water per acre. Use a properly calibrated, low-pressure, boom-type sprayer having good agitation.

APPLICATION TIMING:

- Apply DYFONATE-TILLAM 1-4E before bedding and transplanting. Incorporate and bed up immediately after application.

INCORPORATION DIRECTIONS:

- DYFONATE-TILLAM 1-4E must be mixed (incorporated) into the soil IMMEDIATELY after application to prevent loss by evaporation of the herbicide.
- Whenever possible, application and incorporation should be done in the same operation.

Tandem disc:

- Can be used on all soil types.
- Set disc to cut 4 to 6 inches deep.
- Operate disc at 4 to 6 MPH.
- Follow disc by a harrow or leveling device slightly wider than the disc.
- Two passes in different directions will improve control. The second pass should be at a slight angle to the first.
- On the second pass the disc should be operated no deeper than on the first pass and
- If deep germinating weeds are present, a second incorporation will improve control or suppression. Weeds which fall into this category are seedling johnsongrass and nutsedge.

Power Driven cultivation equipment:

- Soil should have previous primary tillage.
- Ground speed must be adjusted with PTO driven cultivation tools to ensure thorough incorporation.
- Set horizontal action line equipment to cut 4 to 6 inches deep.
- Set vertical action line equipment to cut 3 to 4 inches deep.

Field cultivator:

- Use on light soils in good till.
- Equip cultivator with 3 to 4 rows of sweeps spaced at 7 inches or less and staggered from row-to-row to leave no soil unturned or undisturbed.
- Set the field cultivator to cut 4 to 6 inches deep.
- Operate at 5 MPH or faster.
- Follow by a harrow or leveling device.
- Chisel plows or points should not be used.
- Two passes in different directions will improve control. The second pass should be at a slight angle to the first.
- If deep germinating weeds are present, use a tandem disc for the first pass.

BEST AVAILABLE COPY

~~Incorporate using tandem discs set to cut to a depth of 6 inches, followed by a spike-toothed harrow or some other leveling device which extends beyond the ends of the discs. For thorough mixing, disc in two different directions (cross disc).~~

~~Before bedding and transplanting, apply EDGE, broadcast, at 1 gallon per acre. Incorporate and bed up immediately.~~

TRANSPLANTING:

For maximum weed control, the tobacco may be transplanted immediately after application or application and bedding.

TOBACCO RECOMMENDATIONS

~~Flue-Cured Tobacco:~~

CULTIVATION:

- This product is not persistent in the soil and susceptible weeds germinating late in the season may not be controlled.
- Cultivation may be necessary to control these susceptible weeds that escape control as well as those weeds not susceptible.
- Do not cultivate deeper than the depth of incorporation of the product.

TANKMIX COMBINATIONS

**DYFONATE-TILLAM 1-4E/DEVIRINOL® 50-DF and
DYFONATE-TILLAM 1-4E/DEVIRINOL 50-WP**

Additional weeds controlled by DYFONATE-TILLAM 1-4E/DEVIRINOL tankmix

GRASSES	
COMMON NAME	SCIENTIFIC NAME
Johnsongrass, seedling	<i>Sorghum halepense</i>
Fall panicum	<i>Panicum dichotomiflorum</i>
Witchgrass	<i>Panicum capillare</i>
Stinkgrass	<i>Eragrostis cilianensis</i>

BROADLEAVES	
COMMON NAME	SCIENTIFIC NAME
Carpetweed	<i>Mollugo verticillata</i>
Purple cudweed	<i>Gnaphalium purpureum</i>
White clover	<i>Trifolium repens</i>

USE RATE:

- Tank mix DEVRINOL® 50-WP or 50-DF selective herbicide at the rate of 2 pounds per acre with the recommended rate of 1 gallon of DYFONATE-TILLAM 1-4E. The DEVRINOL 50-DF or 50-WP should be added to the tank first and properly agitated and dispersed before adding the DYFONATE-TILLAM 1-4E. Agitation should be maintained in the spray tank until empty.

APPLICATION DIRECTIONS:

- The tankmix directions are the same as for DYFONATE-TILLAM 1-4E alone.

APPLICATION TIMING:

- Apply the tankmix of DEVRINOL and DYFONATE-TILLAM 1-4E before bedding and transplanting.
- Incorporate and bed up immediately after application.

INCORPORATION DIRECTIONS:

- The tankmix directions are the same as for DYFONATE-TILLAM 1-4E alone.

TRANSPLANTING:

- For maximum weed control, the tobacco may be transplanted immediately after application or application and bedding.

DYFONATE-TILLAM 1-4E/DYFONATE 4-EC

Tank mix 1 quart of DYFONATE 4-EC for the control of severe insect infestations.

Follow the use directions for DYFONATE-TILLAM 1-4E for application, incorporation, and transplanting. Follow all use precautions on both the DYFONATE-TILLAM 1-4E and DYFONATE 4-EC labels.

DYFONATE-TILLAM 1-4E IMPREGNATION ON DRY BULK FERTILIZERS

CAUTION: DYFONATE-TILLAM 1-4E MUST NOT BE IMPREGNATED ON AMMONIUM NITRATE, SODIUM NITRATE, OR POTASSIUM NITRATE. SUCH MIXTURES MAY CAUSE EXPLOSION AND FIRE.

ALL DYFONATE-TILLAM 1-4E AND DYFONATE 4-EC HANDLING INSTRUCTIONS AND LABEL RECOMMENDATIONS REGARDING RATES PER ACRE, SOIL INCORPORATION, APPLICATION, CAUTIONS, GENERAL USE PRECAUTIONS, AND OTHER DIRECTIONS MUST BE FOLLOWED.

ALL INDIVIDUAL STATE REGULATIONS RELATING TO BULK DRY FERTILIZER BLENDING, REGISTRATION, LABELING, AND APPLICATION ARE THE RESPONSIBILITY OF THE INDIVIDUAL AND/OR COMPANY SELLING THE FERTILIZER AND DYFONATE-TILLAM 1-4E MIXTURE.

DYFONATE-TILLAM 1-4E may be impregnated on many dry bulk fertilizers (see following chart of approved dry fertilizer ingredients) and applied and incorporated in the soil before planting for the control of insects, grasses, and broadleaf weeds in tobacco.

DYFONATE-TILLAM 1-4E on bulk dry fertilizers provides weed and insect control equal to DYFONATE-TILLAM 1-4E applied as a spray in water. However, uniform impregnation of DYFONATE-TILLAM 1-4E on the dry fertilizer particles and uniform application in the field are necessary to assure good results.

For impregnating DYFONATE-TILLAM 1-4E on dry fertilizer, use a closed rotary-drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle (or nozzles) should be positioned inside of the mixer to provide uniform spray coverage of the tumbling fertilizer. The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with DYFONATE-TILLAM 1-4E provides a satisfactory, dry mixture. If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry, flowable mixture. Micro-Cel™ E is the recommended absorbent powder. It should be added separately and uniformly to the prepared DYFONATE-TILLAM 1-4E fertilizer mixture in a quantity that is sufficient to provide a suitably flowable mixture. Generally less than 2% by weight of Micro-Cel E is required. The amount of DYFONATE-TILLAM 1-4E actually required in the manufacture of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of DYFONATE-TILLAM 1-4E actually contained in the mixture applied to the soil represents the correct rate of use.

A minimum of 200 pounds and a maximum of 2,000 pounds of approved ingredients (see following chart of approved dry fertilizer ingredients) impregnated with DYFONATE-TILLAM 1-4E at the recommended rate must be applied per acre.

Bulk fertilizer impregnated with DYFONATE-TILLAM 1-4E should be applied immediately, not stored. It is recommended that all bulk containers be tightly covered while the product is being transported and applied to reduce chances of DYFONATE-TILLAM 1-4E loss via volatilization.

BEST AVAILABLE COPY

130416

APPROVED DRY FERTILIZER INGREDIENTS

	N	P	K
Ammonium sulfate	21	0	0
Diammonium phosphate	18	16	0
Potassium chloride	0	0	60
Potassium sulfate	0	0	52
Super-phosphate (single)	0	20	0
Super-phosphate (triple)	0	48	0
Urea	45	0	0
Ammonium phosphate sulfate	16	20	0
11-48-0	11	48	0

Note: K-Mag and two MAP fertilizers (12.5-50-0, 11-54-0) have been shown to be compatible with DYFONATE-TILLAM 1-4E and are approved for use.

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZERS WITH DYFONATE-TILLAM 1-4E

Fertilizer Rate Per Acre (Lbs.)	DYFONATE-TILLAM 1-4E Rate Per Acre (4 qts./acre)
200	40 qts./ton
300	26.7 qts./ton
400	20.0 qts./ton
500	16.0 qts./ton
600	13-1/3 qts./ton
800	10 qts./ton
1000	8 qts./ton
1500	5.4 qts./ton
2000	4 qts./ton

5 1/3 qts./ton
12/2/91
EAL

BEST AVAILABLE COPY

140416

For insect control under severe infestation, add DYFONATE 4-EC to the spray mix as follows:

Fertilizer Rate Per Acre (Lbs.)	DYFONATE 4-EC
200	30 qts./ton
300	6-2/3 qts./ton
400	5 qts./ton
500	4 qts./ton
600	3-1/3 qts./ton
800	2-1/2 qts./ton
1000	2 qts./ton
1500	1-1/3 qts./ton
2000	1 qt./ton

150416

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Do not transport or store with any food or feed intended for human or animal consumption. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

~~**Glass:** Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.~~

Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

)
)
Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

)
)
When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIABILITY of Seller will be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

)
)
This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States of America.

DEVRINOL®, DYFONATE®, and TILLAM® are trademarks of ICI Group Companies.
Micro-Cel™ is a trademark of Marville Sales Corporation.

BEST AVAILABLE COPY