



ACTIVE INGREDIENTS:

- S ethyl disobutylthio carbamate
Atrazine 2-chloro 4 (ethylamino) 6 (isopropylamino) s triazine
Related triazines

18.0%
5.7%
0.3%
76.0%
100.0%

INERT INGREDIENTS

Contains 18 pounds of S ethyl disobutylthio carbamate per 100 pounds of active ingredient.

Directions For Use

2012

Use in violation of Federal law to use this product for a purpose not intended with its labeling.
This product may not be mixed or loaded, or used within 10 feet of all wells, including abandoned wells, through wells, and sink holes.
Do not apply this product through any type of irrigation system.
SUTAZINE + 1B (6-G) is a selective herbicide that is active against non-crop plants and weeds for control of weeds listed on this label by interfering with normal germination and seedling development. It does not control established weeds. All weeds, grasses and crop stubble should be thoroughly worked into the soil before treatment.
Minimum application to corn must be made before corn reaches 1 1/2 feet in height.
The maximum application rate for corn is 3 pounds of atrazine active ingredient per acre per calendar year.

Keep Out of Reach of Children

CAUTION

Harmful If Swallowed

STATEMENT OF PRACTICAL TREATMENT

- If a known exposure occurs or is suspected, immediately start the first aid measures given below. If further treatment is required, you must contact a Poison Center, a physician, or the nearest hospital.
If Swallowed: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed in the back of the victim's tongue. Give fluids until vomitus is clear. If victim is unconscious, do not induce vomiting. Do not give anything by mouth.
If On Skin: Flush all affected areas with plenty of water for several minutes. Remove and clean contaminated clothing and shoes before reuse. Seek medical attention if irritation occurs.
If In Eyes: Hold eyelids apart and flush with plenty of water for 15 minutes. Seek medical attention if irritation occurs.
If Inhaled: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633)
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident
CALL CHEMTREC 1-800-424-9300

For other product information, contact your local ICI Americas Distributor or see Material Safety Data Sheet

PRECAUTION STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

HARMFUL IF SWALLOWED

Avoid contact with skin, eyes and clothing. Avoid inhalation of dust. Wash thoroughly with soap and water after use. Users are required to wear long sleeve shirts and long pants or equivalent, chemical resistant gloves, and boots (when needed). In addition, persons involved in mixing or loading operations are required to wear boots of resistant rubber or neoprene gloves and a face shield or goggles.

ENVIRONMENTAL HAZARDS

Atrazine leaches readily and accepted label rates may be leached to result in contamination of water supply if a way of groundwater. Therefore, users are advised to avoid use of atrazine in well drained soils, particularly those having high groundwater tables. Ground water contamination may be reduced by the design and timing of permanent liquid bulk storage sites with an impermeable material. This pesticide does not bioaccumulate. Do not apply directly to water or wetlands. Runoff from treated areas may be hazardous to aquatic organisms in nearby streams. Do not contaminate water when disposing of equipment and mixtures.

GENERAL USE PRECAUTIONS

- Read all label directions before using. Apply this product as specified on this label.
Make only one application per crop.
After treatment with SUTAZINE + 1B (6-G) do not plant any crop except corn the following year, or injury may occur.
Do not use on milk or sirghum. Do not use on corn seed stock (breeders, foundation or breeder) SUTAZINE + 1B (6-G) is not to be used on Production Seed corn.
This product is recommended for use on mineral soils only.
Applied according to directions and under normal growing conditions, SUTAZINE + 1B (6-G) will control the broadleaf weeds, grasses, and annual weeds listed in the table below and will not injure corn. However, in areas of drought or disease attack, the use of certain soil applied systems may be needed, especially plant herbicides or soil insecticides, to create a normal condition that will allow crop seedlings of SUTAZINE + 1B (6-G) to establish under these abnormal conditions. Consult your crop injury.
Do not take or allow children, pregnant women, or other persons to be near treated areas until the product has dried through the soil.
Do not contaminate food or feed.
Do not store near seeds, fertilizers, or foodstuffs.
Do not contaminate irrigation water or water used for domestic purposes.

WEEDS CONTROLLED

This product will not control established weeds.

ANNUAL GRASSES

- Barnyardgrass (Echinochloa crusgalli)
Cyngrass (Digitaria sp.)
Fall Panicum (Panicum de hybridum)
Giant Foxtail (Setaria faberii)
Goosegrass (Eleusine indica)
Green Foxtail (Setaria verticillata)
Seeding Bermudagrass (Cynodon dactylon)
Seeding Johnsongrass (Sorghum halepense)
Shattercane (Sorghum bicolor)
Wild Oats (Avena fatua)
Witchgrass (Panicum capillare)
Yellow Foxtail (Setaria glauca)
Texas Panicum (Buffalograss) (Lolopogon grass)
(Texas Panicum)

\*Cultivation is suggested in addition to the SUTAZINE + 1B (6-G) treatment in fields with heavy infestation of shattercane or buffalograss. (Texas panicum)

ANNUAL BROADLEAF WEEDS

- Annual Morningglory (Ipomoea purpurea)
Carpetweed (Mollis verticillata)
Cockspur (Common) (Ranunculus abortivus)
Jimsonweed (Datura stramonium)
Lambquarters (Rumex crispus)
Mustards (Brassica sp.)
Nightshade (Solanum sp.)
Common Purslane (Portulaca oleracea)
Ragweed (Ambrosia sp.)
Redroot Pigweed (Amaranthus retrofractus)
Prostrate Pigweed (Amaranthus granifolius)
Smooth Pigweed (Amaranthus hybridus)
Smartweed (Polygonum sp.)
Velvetleaf (Abutilon theophrasti)
Flower Purslane (Ricinus communis)

PERENNIAL GRASSES

- Purple Nutsedge (Cyperus rotundus)
Yellow Nutsedge (Cyperus esculentus)

Existing stands of nutgrasses must be treated under and/or topped up thoroughly before treatment. Cultivation is suggested in addition to the SUTAZINE + 1B (6-G) treatment in fields with moderate to heavy nutgrass infestations.

STORAGE AND DISPOSAL

- PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
STORAGE: Keep container closed when not in use. Do not store near seeds, fertilizers, or foodstuffs.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER: Completely empty bags into applicable equipment. Then dispose of empty bags in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Do not reuse empty containers.

Recommendations

Do not use on any type corn in Arizona or in the ten southernmost California counties (except Kern County where field and sugar corn may be treated).
NOTE: All doses expressed as broadcast rates per acre. For band applications, reduce dosage proportionately depending upon the row spacing and band width to be treated.

BEFORE PLANTING (PRE-PLANT) APPLICATION

In the North Central, Northeastern and Midwestern Corn Growing Areas
Apply and incorporate 16.5 to 22.5 pounds of SUTAZINE + 1B (6-G) per broadcast acre. Use the higher rates on extremely heavy soils (the heaviest clay soils with heavy subsoil matter). On fields with moderate to heavy infestations of shattercane or nutgrass use the higher rate regardless of soil texture except on low sandy and the light soils (course textured) sandy soil of the Eastern Corn Belt States. Cultivations are suggested in addition to the SUTAZINE + 1B (6-G) treatment in fields with moderate to heavy infestations of shattercane or nutgrass.

In All Other Corn Growing Areas

Apply and incorporate 16.5 pounds of SUTAZINE + 1B (6-G) per broadcast acre. Cultivation is suggested in addition to the SUTAZINE + 1B (6-G) treatment in fields with moderate to heavy infestations of shattercane or nutgrass or buffalograss (Texas pan corn).

AFTER PLANTING APPLICATION

NOTE: This corn crop is in the following states only: North and South Dakota, Nebraska, Kansas, Missouri and Iowa.
In all other conditions in the above named states, make pre-plant SUTAZINE + 1B (6-G) application unless the product may be applied and mixed into the soil at any time after planting before emergence of corn. Apply and incorporate 16.5 pounds of SUTAZINE + 1B (6-G) per broadcast acre. For band applications, use standard planter after bed preparation (disc, moldboard, or other) application of granular herbicides. Normally, application and incorporation are done in the same operation using a system that allows for the back of the applicator to be immediately adjustable, and similar equipment for separate application of herbicide. A rotary harrow can be used set for a depth of 1 1/2 inches, or operated at 2 to 3 inches.

NOTE: After planting application is recommended for control of nutgrass. Topsoil must be stirred after the BEFORE PLANTING (PRE-PLANT) APPLICATION for control of these weeds.

CULTURAL PRACTICES FOLLOWING APPLICATION

Shallow weeding, a shallow cultivation or rotary hoeing will generally result in better weed control. When cultivating for any reason, it should be shallow, not more than 1/2 the depth the herbicide was incorporated.

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, inefficient use or other unintended consequences may result from the use of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is a reliable ally in the purposes stated on the label subject to the inherent risks referred to above when used in accordance with the label under normal conditions of use. This warranty does not extend to the use of this product in violation of the label or in violation of any state, federal, or other conditions or order conditions not reasonably foreseeable to or beyond the control of Seller and Buyer, and they assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use of this product, the following limitations shall apply: (1) the amount of such losses or damages shall be limited to the actual loss sustained by Buyer or User, and (2) the amount of such losses or damages shall be limited to the amount of the purchase price paid for the product. Seller and Buyer agree to hold Seller harmless for any claims relating to such factors.

This product is not registered for use in California, Oregon, Washington, Idaho, Utah, Nevada, Arizona, New Mexico, Texas, Oklahoma, Missouri, Arkansas, Louisiana, Mississippi, Alabama, Georgia, Florida, South Carolina, North Carolina, Virginia, West Virginia, Maryland, Delaware, Pennsylvania, New Jersey, New York, Connecticut, Rhode Island, Massachusetts, Vermont, New Hampshire, Maine, New Brunswick, Nova Scotia, Prince Edward Island, New Brunswick, Ontario, Quebec, and the District of Columbia.


EPA Reg No 10182-201
EPA Est No 10182-NB-1

AVAILABLE COPY
NET CONTENTS 1500 LB. 680 kg



Agricultural Products

ICI Americas Inc.
Wilmington, Delaware 19897



090104