

FEB 24 1994

Ms. Cheryl Bohenick
Zeneca Inc.
Zeneca Ag Products
P.O. Box 751
Wilmington, DE 19897

Dear Ms. Bohenick:

Subject: Devrinol® 2-E Selective Herbicide
Worker Protection Standards
EPA Reg. No. 10182-219
Your submission dated Feb. 1, 1994

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable and a stamped copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

BEST AVAILABLE COPY

DEVINOL 2-E

Selective Herbicide—Soluble Liquid

For use on Tomatoes, Peppers, Strawberries
and Tobacco

COMPLETE DIRECTIONS FOR USE

AGRICULTURAL
with COMMENTS
in EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

ACTIVE INGREDIENT
N,N-dimethyl-2-(1-naphthyl)propionamide 21.8%
INERT INGREDIENTS 78.2%
TOTAL 100.0%

Contains 2 pounds active ingredient per gallon
EPA Reg. No. 10182-219

KEEP OUT OF REACH OF CHILDREN DANGER

PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

FIRST AID

CALL A POISON CENTER OR A PHYSICIAN IMMEDIATELY

If a known exposure occurs or is suspected, immediately start the procedures given below. Simultaneously contact a Poison Center, a physician or the nearest hospital. NOTE: Be sure to advise the person contacted that the product contains a hydrocarbon solvent. Describe the type of exposure, the victim's symptoms, and follow the advice given.

IF SWALLOWED: Immediately give several glasses of water but do not induce vomiting. If vomiting does occur, give fluids again. Have a physician determine if condition of patient will permit induction of vomiting or evacuation of stomach. Do not give anything by mouth to an unconscious or convulsing person.

IN CASE OF CONTACT: Immediately flush eyes and skin with large amounts of running water for at least 15 minutes while removing contaminated clothing and shoes. For eyes, hold eyelids apart during the flushing to ensure rinsing of the entire surface of the eye and lids with water. Get medical attention immediately. Wash clothing before reuse.

IF INHALED: Remove to fresh air. If victim is breathing, oxygen may be given from a demand-type or continuous-flow inhaler, preferably with a physician's advice. If not breathing, clear victim's airway and start mouth-to-mouth artificial respiration. Get medical attention immediately.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633)

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9308.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

CORROSIVE. CAUSES EYE DAMAGE AND SKIN IRRITATION. Do not get in eyes, on skin, or on clothing. Harmful if swallowed or absorbed through the skin. Avoid contamination of food or feed. Do not use, store, spill, or pour near heat or open flame.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category H on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves, such as Barrier Laminar or Viton 214 mils.
- Shoes plus socks.
- Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or airblast in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters. Apply this product as specified on this label. Do not contaminate water used for domestic purposes. Avoid contamination of Food and Feed.

GENERAL PRECAUTIONS

Read all label directions before using. Use only for recommended purposes at recommended rates. Do not overdo. Do not contaminate irrigation water or water used for domestic purposes. Do not store near seeds or fertilizer. Do not apply more than once per season. DEVINOL 2-E is recommended for use on mineral soils only. Not recommended for use on soils with over 10% organic matter. Spraying on ground heavily covered with leaves or trash may adversely affect weed control.

ZENECA

AGRICULTURAL
with COMMENTS
in EPA Letter Dated

FEB 24 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

10182-219

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. The Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protection equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical-resistant gloves, such as Barrier Laminar or Viton 214 mils.
- Shoes plus socks.
- Protective eyewear.

DEVINOL 2-E is a selective, surface-applied herbicide for use in strawberries. It is also recommended for weed control in certain vegetables and tobacco, applied and incorporated prior to planting.

DEVINOL 2-E controls weeds before they emerge. It does not control established weeds. All weed growth and crop stubble in fields intended for vegetables or tobacco must be thoroughly worked into the soil prior to planting.

Spray equipment should be carefully calibrated before use and checked frequently during application to be sure nozzles are free from clogging and are delivering a uniform spray pattern. Apply the recommended rate of DEVINOL 2-E in 20 to 100 gallons of water per acre.

STRAWBERRIES

NOTE: DEVINOL does not control established weeds.

WEEDS CONTROLLED	
ANNUAL GRASSES:	
Annual Bluegrass	<i>Poa annua</i>
Barnyardgrass—Watergrass	<i>Echinochloa spp.</i>
Bristly Foxtail	<i>Setaria verticillata</i>
Large (Heavy) Crabgrass	<i>Digitaria sanguinalis</i>
Ripgut Brome	<i>Bromus rigidus</i>
Southwestern Cupgrass	<i>Eriochloa gracilis</i>
Wild Barley	<i>Hordeum lapponicum</i>
Wild Oats	<i>Avena fatua</i>
ANNUAL BROADLEAF WEEDS:	
Chickweed	<i>Stellaria media</i>
Common Fiddleneck	<i>Amsinckia intermedia</i>
Common Knotweed	<i>Polygonum aviculare</i>
Common Purslane	<i>Portulaca oleracea</i>
Common Sowthistle	<i>Sonchus oleraceus</i>
Fleese	<i>Erodium spp.</i>
Groundsel	<i>Senecio spp.</i>
Little Mallow	<i>Malva parviflora</i>
Lambquarters	<i>Chenopodium album</i>
Pineappleweed	<i>Melicaria melicarioides</i>
Prickly Lettuce	<i>Lactuca scariola</i>
Redroot Pigweed	<i>Amaranthus retrofractus</i>

*From seed.

See reverse for Recommendations

BEST AVAILABLE COPY

Made in U.S.A.

ZENECA Ag Products

ZENECA Inc.
Wilmington, Delaware 19887

for band or strip treatment.

STRAWBERRIES (Post Transplant and Direct Seeded): For control of all weeds listed, apply 2 qts. DEVRINOL 2-E per broadcast acre to the soil surface. Apply in the fall through early spring, prior to weed emergence. Do not apply DEVRINOL 2-E to frozen ground. Do not apply DEVRINOL 2-E from bloom through harvest. For best results, the treatment must reach the zone of weed seed germination. Treatments made other than in November through February must be irrigated-in within 24 hours (sprinkler or flood) with sufficient water to wet the soil to a depth of at least 2 to 4 inches. Irrigation or shallow incorporation is recommended for treatments made from November through February if no rainfall occurs within one week of application.

VEGETABLES & TOBACCO

NOTE: DEVRINOL 2-E does not control established weeds.

WEEDS CONTROLLED	
ANNUAL GRASSES:	
Barnyardgrass	<i>Echinochloa</i> spp.
Crabgrass	<i>Digitaria</i> spp.
Fall Panicum	<i>Panicum dichotomiflorum</i>
Goosegrass	<i>Elymus indica</i>
Seeding Johnsongrass	<i>Sorghum halepense</i>
ANNUAL BROADLEAF WEEDS:	
Carpweed	<i>Malva verticillata</i>
Common Purslane	<i>Portulaca oleracea</i>
Lambquarters	<i>Chenopodium album</i>
Redroot Pigweed	<i>Amaranthus retrofractus</i>

DIRECTIONS FOR USE

On flat-landed or bedded crops apply, before planting, to well worked soil that is dry enough to permit thorough incorporation to a depth of 1 to 2 inches. Incorporate the same day as applied, using equipment which will result in uniform incorporation of the herbicide to the desired depth.

After harvest or prior to planting of succeeding crops, a deep moldboard or disc plowing operation must be carried out.

To avoid injury, do not seed to alfalfa, small grains, sorghum, corn, lettuce or sugar beets for 12 months after DEVRINOL 2-E application.

RECOMMENDATIONS

NOTE: All rates are given on a broadcast basis. Reduce rates proportionately for band or strip treatment.

TOMATOES (Direct Seeded and Transplanted):

Apply 1/2 to 1 gallon DEVRINOL 2-E per broadcast acre. Use the lower rate on light soil (coarse textured—sandy) and the higher rate on heavy soil (fine textured—clay).

PEPPERS (Direct Seeded and Transplanted):

Apply 1/2 to 1 gallon DEVRINOL 2-E per broadcast acre. Use the lower rate on light soil (coarse textured—sandy) and the higher rate on heavy soil (fine textured—clay).

TOBACCO (Transplanted):

Before transplanting, apply 1/2 to 1 gallon DEVRINOL 2-E per broadcast acre. Use the lower rate on light soils (coarse textured—sandy) and the higher rate on heavy soil (fine textured—clay).

DEVRINOL 2-E/TILLAM 6-E HERBICIDE TANKMIX COMBINATION

A tankmix combination of DEVRINOL 2-E and TILLAM 6-E herbicide can be applied and mixed (incorporated) into the soil before planting for weed control in tomatoes and tobacco.

IMPORTANT:

- Observe all cautions and limitations on labeling of all products used in combination.
- Make only one application per season.
- In each case where a range of rates is given, the lower rates should be used on light (coarse) textured soils and soils low in organic matter. The higher rates should be used on heavy (fine) textured soils.
- After harvest or prior to planting of succeeding crops, a deep moldboard or disc plowing operation must be carried out.
- To avoid injury, do not seed to alfalfa, small grains, sorghum, corn, lettuce or sugar beets for 12 months after application of DEVRINOL 2-E/TILLAM 6-E tankmix.

The tankmix is recommended for control of the following weeds.

WEEDS CONTROLLED	
ANNUAL GRASSES:	
Barnyardgrass—Watergrass	<i>Echinochloa</i> spp.
Bermudagrass	<i>Cynodon dactylon</i>
Crabgrass	<i>Digitaria</i> spp.
Foxtail—yellow, giant, green	<i>Setaria</i> spp.
Goosegrass	<i>Elymus indica</i>
Millet	<i>Setaria</i> spp.
Signalgrass	<i>Brachiaria</i> spp.
Wild Oats	<i>Avena fatua</i>
ANNUAL BROADLEAF WEEDS:	
Black-eyed Susan; Sweet Suzy	<i>Rudbeckia hirta</i>
Common Purslane	<i>Portulaca oleracea</i>
Deadnettle; Henbit	<i>Lamium amplexicaule</i>
Florida Pusley	<i>Richardia scabra</i>
Hairy Nightshade	<i>Solanum villosum</i>
Lambquarters	<i>Chenopodium album</i>
Nasturtium Goosefoot	<i>Chenopodium murale</i>
Prostrate Pigweed	<i>Amaranthus prostratus</i>
Redroot Pigweed	<i>Amaranthus retrofractus</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
PERENNIAL WEEDS:	
Yellow Nutsedge (Yellow Nutgrass)	<i>Cyperus esculentus</i>
Purple Nutsedge (Purple Nutgrass)	<i>Cyperus rotundus</i>

Existing stands of nutsedge must be turned under and chopped up thoroughly before treatment.

DIRECTIONS FOR USE

The recommended tankmix controls weeds by interfering with normal seed germination and seedling development. It does not control established weeds. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

TANK MIXING DIRECTIONS

Be sure the sprayer tank, boom and nozzle have been thoroughly flushed with clean water if another pesticide has been used previously.

Sufficient jet or mechanical agitation should be provided during the filling operation and during application. Fill the spray tank at least one-fourth full of clean water and while pump and agitator are running, add the recommended amounts of the herbicides to the tank. Fill the tank with water to desired level. Do not allow the mixture to stand in the spray tank without agitation.

Allow the pump to run several minutes before application to insure proper suspension and mixing. The tankmix should not be left in the spray tank for prolonged periods; batches should be mixed and applied the same day.

APPLICATION DIRECTIONS

The soil should be well worked and dry enough to permit good soil mixing (incorporation). Pour the recommended amounts of DEVRINOL 2-E and TILLAM 6-E into a spray tank during the filling operation. Apply in 25 to 50

in irrigated areas, do not apply before reduce dosage proportionately if the soil is to be treated. Do not use on rocky ground because thorough incorporation is not possible.

INCORPORATION DIRECTIONS

The herbicides must be mixed (incorporated) into the soil IMMEDIATELY after application to prevent loss by evaporation. Whenever possible, application and incorporation should be done in the same operation.

Incorporation Before Planting: Use power-driven cultivation equipment set to cut to a depth of two to three inches in coarse soils and three to four inches in heavy soils OR a tandem disc set to cut to a depth of six inches, followed by a spike-toothed harrow or some other leveling device which extends beyond ends of the disc. For thorough mixing, disc in two different directions (cross disc).

Incorporation At Planting: Use power-driven cultivation equipment set to cut to a depth of two to three inches OR, on coarse soils only, ground-driven tillers (rolling cultivators, rotary hoe, row wheels, etc.) set to cut to a depth of 1 1/2 inches and operated at 6 to 8 mph. When using ground-driven tillers, pre-cultivate if soil is crusty.

PLANTING

Crops should be planted or seeded immediately after application and incorporation. Do not use a drag behind the planter as it may concentrate the herbicides over the seed row and cause crop injury.

CULTURAL PRACTICES FOLLOWING APPLICATION

Shallow cultivation may be necessary to control those susceptible weeds that escape control as well as those weeds not susceptible to either herbicide. Do not cultivate deeper than the depth of herbicide incorporation.

When cultivating fields where the tankmix has been banded in the row, use shields to prevent the movement of untreated soil into the treated row.

RECOMMENDATIONS

TOMATOES (Direct Seeded (California Only) and Transplanted): Apply 1/2 to 1 gallon DEVRINOL 2-E plus 2 1/2 to 4 quarts TILLAM 6-E per broadcast acre. Use the lower rates on light soils (coarse textured—sandy) and the higher rate on heavy soils (fine textured—clay). Do not use hotcaps on transplanted tomatoes.

TOBACCO (Transplanted):

Before transplanting, apply 1/2 gallon DEVRINOL 2-E plus 2 1/2 quarts TILLAM 6-E per broadcast acre.

DEVRINOL 2-E/TILLAM 6-E/DYFONATE 4-EC

INSECTICIDE TANKMIX COMBINATION

A tankmix combination of DEVRINOL 2-E, TILLAM 6-E and DYFONATE 4-EC insecticide can be applied and mixed into the soil before planting tomatoes or tobacco, for control of weeds and soil insects. Refer to DEVRINOL/TILLAM Tankmix Combination for list of weeds controlled. Refer to the DYFONATE 4-EC label for list of insects controlled.

NOTE: DYFONATE 4-EC is a Restricted Use Pesticide. Observe all cautions and limitations on labeling of all products used in combinations. Make only one application per season. Do not contaminate water by cleaning equipment or disposal of wastes.

DIRECTIONS FOR USE

The recommended tankmix controls weeds by interfering with normal seed germination and seedling development. It does not control established weeds. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

For GENERAL USE PRECAUTIONS AND DIRECTIONS FOR USE refer to DEVRINOL 2-E/TILLAM 6-E Tankmix Recommendations for methods of tank mixing, application and incorporation.

RECOMMENDATIONS

TOMATOES (Direct Seeded (California Only) and Transplanted): Apply 1/2 to 1 gallon DEVRINOL 2-E plus 2 1/2 to 4 qts. TILLAM 6-E plus 2 qts. DYFONATE 4-EC for control of symphytines or 4 qts. for control of symphytines and wireworms in 25 to 50 gallons of water per broadcast acre. Use the lower DEVRINOL 2-E and TILLAM 6-E rates on light soils (coarse textured—sandy) and the higher rates on heavy soils (fine textured—clay). Do not use hotcaps on transplanted tomatoes.

TOBACCO (Transplanted):

Before transplanting, apply 1/2 gallon DEVRINOL 2-E plus 2 1/2 qts. TILLAM 6-E plus 2 qts. DYFONATE 4-EC for control of wireworms in 25 to 50 gallons of water per broadcast acre.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse container.

STORAGE: Store in a cool dry place. Do not store below 20°F. Keep container closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherent in the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIABILITY of Seller will be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

DEVRINOL®, DYFONATE®, and TILLAM® are trademarks of ZENECA Group Companies.

BEST AVAILABLE COPY


ACTIVE Ingredient(s)	
N,N-diethyl-2-(1-naphthylthio) propanesulfonate	21.0%
INERT Ingredient(s)	79.0%
TOTAL	100.0%

Contains 2 pounds of active ingredient per gallon

EPA Reg. No. 10102-10

EPA Est. No. 10102-CA-2

To obtain the order, call to us today. Because a deposit goes toward the purchase of your car, you can get the full value of the order when you trade in your car. (See the full details on the back of the order.)



Contents: 5 gal (18.9 L)

BEST AVAILABLE COPY

with COMMENTS
to EPA Letter Dated:

FEB 24 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the purpose of
registered under EPA Reg. 40.1.

~~10182-219~~

STATEMENT OF PRACTICAL TREATMENT

[illegible]

PRECAUTIONARY STATEMENTS

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-7-A-S-T-A-S-E (287-6886).
FOR CHEMICAL EMERGENCY 24 Hrs. Incl. Ev. response, or accident call CHEMTREC 1-800-424-9300.

DANGER

[illegible]

Personal Protective Equipment
Some materials that are chemical-resistant by this product are listed below. If you wear appropriate clothing and use the appropriate PPE for Category II or III chemical resistance category, select in Part 1 Applications and other hazards made up of:

- Large round and long pants
- Chemical-resistant gloves, such as Butyl Latex or Viton, 18 mils
- Shoes plus socks
- Protective eyewear

DEVRINO! 2-E

Selective Her. John—Lumber/Mobile & Land

**For use on Tomatoes, Peppers,
Strawberries and Tobacco**



Decided clothing and other apparel exports that have been designated as highly restricted with that country's apparel exports. From 1990 to 1992, the U.S. Department of Commerce's Bureau of Economic Analysis reported that PPE is not such indicators for countries, we developed and we have also used PPE regularly from other country.

When analyzing the data on exports, we used the data on exports of apparel in the U.S. Department of Commerce's Bureau of Economic Analysis. The data on exports of apparel in the U.S. Department of Commerce's Bureau of Economic Analysis is reported in the U.S. Department of Commerce's Bureau of Economic Analysis.

UN 1993



FLAMMABLE LIQUID

3

Flammable liquid, n.o.s.
(contains chlorobenzene).

ZENECA Ag Products

DYFONATE® II 20-G

**Granular Insecticide
Organophosphate**

**Soil Insecticide for Protection Against Corn
Rootworm, Black Cutworm, and Other Major Pests**

ACTIVE INGREDIENT:

O-Ethyl-S-phenylethylphosphonodithioate.....20.0%

INERT INGREDIENTS:.....80.0%

Total.....100.0%

Contains 20 pounds active ingredient per 100 pounds.

*Ionofos

EPA Reg. No. 10182-135

EPA Est. No. 10182-NB-1

**KEEP OUT OF
REACH OF CHILDREN
DANGER ☠ POISON**

Hazardous to Livestock,
Humans, and Domestic Animals

PELIGRO

Si usted no entiende la etiqueta, busque a
alguien para que se la explique o llame al
ataje. (If you do not understand the label
find someone to explain it to you in detail.)

UN 2783



BEST AVAILABLE COPY

See Back Panel For Additional
Precautionary Statements

Net contents: 50 lb (22.7 kg)

In EPA Letter Dated:

FEB 24 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

10182-219

