Ⓐ Do not apply more than 4.6 lbs. of DEVRINOL 2-G per 1,000 sq. ft. in California. PM 25

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

MAY BE HARMFUL IF SWALLOWED. Do not breathe dust. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after use.
- Avoid Contamination of Food and Feed

ENVIRONMENTAL HAZARDS
- Do not apply directly to lakes, streams, and/or ponds
- Do not contaminate irrigation water or water used for domestic purposes

GENERAL PRECAUTIONS
- Read all instructions before using
- Use only for recommended purposes and at recommended rates. Do not overdose

WHAT THIS PRODUCT DOES

DEVRINOL 2-G Ornamental controls certain annual grassy and broadleaf weeds as they germinate. It does not control established weeds.

The weeds controlled are listed below.

WEEDS CONTROLLED BY DEVRINOL 2-G

Annual Grasses:

Barnyardgrass, Junglerice (*Echinochloa spp.*)	Giant Foxtail (*S. faberi*)
Wild Barley (*Hordeum leporinum*)	Green Foxtail (*S. viridis*)
Annual Bluegrass (*Poa annua*)	Yellow Foxtail (*S. lutescens*)
Bromes (*Bromus*)	Goosegrass (*Eleusine indica*)
Cheatgrass (*B. secalinus*)	Seedling Johnsongrass (*Sorghum halepense*)
Downy Brome (*B. tectorum*)	Wild Oats (*Avena fatua*)
Ripgut Brome (*B. rigidus*)	Panicums (*Panicum*)
Soft Chess (*B. mollis*)	Guineagrass (*P. maximum*)
Large (Hairy) Crabgrass (*Digitaria sanguinalis*)	Fall Panicum (*P. dichotomiflorum*)
Smooth Crabgrass (*Digitaria ischaemum*)	Texas Panicum (*P. texanum*)
Southwestern Cupgrass (*Eriochloa gracilis*)	Witchgrass (*P. capillare*)
Foxtails (*Setaria*)	Italian Ryegrass (*Lolium multiflorum*)
Bristly Foxtail (*S. verticillata*)	Sandbur (*Cenchrus spp.*)
	Red Sprangletop (*Leptochloa filiformis*)
	Stinkgrass (*Eragrostis cilianensis*)

Annual Broadleaf Weeds

Carpetweed (*Mollugo verticillata*)	Pineappleweed (*Matricaria matricarioides*)
Common Chickweed (*Stellaria media*)	Common Purslane (*Portulaca oleracea*)
Coast Fiddleneck (*Amsinckia intermedia*)	Horse Purslane (*Trianthema portulacastrum*)
Filaree (*Erodium spp.*)	Common Ragweed** (*Ambrosia artemisiifolia*)
Groundsel (*Senecio spp.*)	Annual Sowthistle (*Sonchus oleraceus*)
Prostrate Knotweed (*Polygonum aviculare*)	
Lambsquarters (*Chenopodium album*)	
Prickly Lettuce (*Lactuca serriola*)	
Little Mallow* (*Malva parviflora*)	
Redroot Pigweed (*Amaranthus retroflexus*)	

*From seed
**Suppression

WHERE TO USE THIS PRODUCT

DEVRINOL 2-G Ornamental can be applied over small ornamental plants or around larger plants in rose gardens, container plantings, ornamental landscapes and ground covers including dichondra.
Use DEVRINOL 2-G Ornamental for weed control in the following plants:

Deciduous and Evergreen Trees

Acer	Dogwood	Japanese Larch	Pin Oak
Ash	Douglas Fir	Laburnum	Pinus
Birch	Eucalyptus	Malus	Pittosporum
Black Walnut	Ficus	Olive (Olea)	Podocarpus
Cedar	Fir	Palm (Washingtonia robusta)	Populus
Citrus	Gleditsia		Prunus
Crataegus	Hemlock	Picea	Pears
Cypress			Pistachios

Deciduous and Evergreen Shrubs

Abelia	Crape Myrtle	Hypericum	Pittosporum
Asparagus (Sprengeri)	Dogwood	Ilex	Podocarpus
Aucuba	Euonymus	Juniperus	Pyracantha
Azalea	Forsythia	Leucothe	Raphiolepis
Bottlebrush	Gardenia	Ligustrum	Rhododendron
Bougainvillea	Heather (Calluna)	Nandina	Rosa
Buxus	Hibiscus	Natal Plum	Star Jasmine
Camellia	Honeysuckle (Lonicera)	Oleander	Taxus
Cotoneaster		Osmanthus	Viburnum
		Photinia	Xylosma

Ground Covers

Ajuga	Erysimum	Iceplant	Osteospermum
Carpobrotus	Gazania	Ivy	Pachysandra
Delosperma	Hedera	Lantana	Sedum
Dichondra	Hypericum	Liriope	Vinca

CONTAINER STOCK

Abelia	Cypress	Ilex	Podocarpus
Ageratum	Dogwood	Juniperus	Pyracantha
Agapanthus	Douglas Fir	Leucothe	Raphiolepis
Asparagus	Eucalyptus	Ligustrum	Rhododendron
Azalea	Euonymus	Liriope	Rosa
Bottlebrush	Fir	Nandina	Sedum
Bougainvillea	Forsythia	Pachysandra	Star Jasmine
Buxus	Gardenia	Palm (Washingtonia robusta)	Thuja
Camellia	Geranium		Vinca
Cedar	Heather	Photinia	Xylosma
Cotoneaster	Hedera	Pinus Spp.	
Crape Myrtle	Hibiscus	Pittosporum	

Continued on right panel

Devrinol™
2G-Ornamental
Selective Herbicide

A Pre-emergence Herbicide for Weed Control in Home Flower Gardens and Landscape Plantings. Use in Deciduous and Evergreen Trees and Shrubs, Ground Covers, Perennial Flowers, Fruit and Nut Crops, Container Grown Plants, and Turfgrass.

ACTIVE INGREDIENT:	
N,N-diethyl-2-(1-naphthalenyloxy)-Propionamide	2.0%
INERT INGREDIENTS:	98.0%
	100.0%

Contains 2 Pounds Active Ingredient per 100 Pounds

Keep Out of Reach of Children
CAUTION
STATEMENT OF PRACTICAL TREATMENT

If an adverse reaction occurs after exposure, promptly initiate the recommended procedures below. If further treatment is required, seek medical attention.

In case of contact: Flush eyes and skin with plenty of water for several minutes. Get medical attention if eye or skin irritation occurs.

If swallowed: Immediately give several glasses of water and induce vomiting by placing a finger on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If inhaled: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL ICI AMERICAS MEDICAL EMERGENCY INFORMATION CENTER (800) 327-8633 OR CHEMTREC (800) 424-9300

For other product information, contact your local ICI Americas Distributor or see Material Safety Data Sheet.

EPA Reg. No. 10182-243 EPA Est. No. 476-NB-1

A-5R

40 LB. NET WT. 18kg

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country United States Patent No. 3,480,671, 3,718,455

Made in U.S.A. by
ICI AMERICAS INC.
AGRICULTURAL PRODUCTS • WILMINGTON, DE 19897

WT 3-88

CONTAINER STOCK (Cont.)

Flowers

African Daisy	Chrysanthemum	Geranium
Aster	Dahlia	Gladiolus
Bird of Paradise	Daisy	Narcissus

Fruit & Nuts

Almond	Filbert	Orange
Apple	Grape	Peach
Apricot	Lemon	Pear
Cherry	Nectarine	Pecan

Strawberry:
(Application should be delayed until the desired number of daughter plants are established. Do not apply from bloom through harvest.)

WHEN TO USE THIS PRODUCT

DEVRINOL 2-G Ornamental should be used before weeds germinate ... established weeds.

Application can be made anytime of the year to weed free soil. Fall ... will provide excellent weed control the following spring and summer.

In containers, apply after potting and watering the plants.

HOW TO USE THIS PRODUCT

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply this product as specified on this label.

DEVRINOL 2-G Ornamental should be spread evenly on the soil surface ... or granular herbicide applicator.

Do not apply when weather conditions favor drift from areas treated.

When weed infestation is heavy or when broadleaves are the primary w... on 145 square feet (6.9 lbs./1,000 sq. ft.). For annual grassy weeds or w... is light, use one pound on 218 square feet (4.6 lbs./1,000 sq. ft.) Ⓐ

DEVRINOL 2-G Ornamental should be incorporated evenly into the s... vator to a depth of 1-2 inches or irrigation or natural moisture (rainfall ... within 2-3 days) for optimum results.

For weed control in dichondra, apply at the time of seeding or to establis... dichondra sod. Irrigate with a minimum of one inch of water.

RECOMMENDATIONS: TURF GRAS...
(For Use on Warm Season Grasses O...

DEVRINOL 2-G is recommended for annual bluegrass, crabgrass and ... the following established turf grasses: bahia, bermudagrass, centiped... fescue.

TIMING OF APPLICATION:
For control of crabgrass and goosegrass, apply DEVRINOL 2-G at the ... Application should be made 2-3 weeks prior to weed emergence. DEVR... applied with a gravity flow drop spreader or a cyclone type spreader. A... cation can be made by spreading half the required amount over the are... applying the remainder at right angles.

DEVRINOL 2-G requires a minimum of one half (½) inch of rainfall or irr... application for adequate control. If irrigation methods are not available ... plication of DEVRINOL 2-G should be moved back several weeks to al... between application and weed germination.

RATES OF APPLICATION:

For Crabgrass Control: Apply 100 to 150 lbs. of DEVRINOL 2-G unifo... 3.38* lbs per 1,000 sq. ft.). Use the higher rate when applying before J... cable north of South Carolina or when weed infestation is expected to ...

For Goosegrass Control: Apply 100 to 150 lbs. of DEVRINOL 2-G per ... per 1,000 sq. ft.). Goosegrass usually starts to germinate 6 to 7 wee... crabgrass germination.

Split Application For Crabgrass and Goosegrass Control: When b... are to be controlled, apply 100 lbs. of DEVRINOL 2-G spread uniformly ... per 1,000 sq. ft.) for crabgrass and another 100 lbs. of DEVRINOL 2-G ... grass. Apply 8-10 weeks apart. Timing should be adjusted to the germi... your area. Split applications give longer and better weed control than o...

For Annual Bluegrass (Poa Annua) Control: Annual bluegrass is a w... of the warm season turf regions. Therefore, application of DEVRINOL 2-G ... the fall before the annual bluegrass germinates. Apply 100-150 lbs. of ... acre (2.25-3.38 lbs. per 1,000 sq. ft.).

Reseeding: Do not reseed or overseed within six months after applicat...

*Where a rate range is given, use the higher rate when weed infestation is expecte...

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE: Keep container closed when not in use. Store in a cool, d... store near seeds or fertilizer or food stuffs.

PESTICIDE DISPOSAL: Wastes resulting from the use of this produ... disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application eq... dispose of empty bag in a sanitary landfill or by incineration, or, if al... and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER AND USER
Seller warrants that this product conforms to the chemical descriptio... is reasonably fit for the purposes stated on the label when used in ac... rections under normal conditions of use. This warranty does not ext... this product contrary to label instructions, or under abnormal use co... conditions not reasonably forseeable to Seller, and Buyer and User a... any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXP... PLIED INCLUDING ANY WARRANTY OF FITNESS OR MERC... SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIA... DAMAGES RESULTING FROM THE USE OR HANDLING OF ... SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S ... SHALL BE LIMITED TO THE REFUND OF THE PURCHASE P...