

Reg # 10182-260

PM-23

UN 2900

UN 2900



PESTICIDES, LIQUID, TOXIC, FLAMMABLE
N.O.S. (molinate/propanil/aromatic solvents)



Agricultural Products

Arrosolo® 3-3E


**emulsifiable liquid herbicide for
use on rice grown in the southern
United States only**

ACTIVE INGREDIENTS:

S-methyl hexahydro-1H-azepine-1-carboxylate	33.1%
N-(3,4-dichlorophenyl) propanamide	33.1%
INERT INGREDIENTS:	33.8%

TOTAL	100.0%
--------------	---------------

Contains 3 Pounds of ORDRAM® and 3 pounds of PROPANIL (Active Ingredients) Per Gallon
EPA Reg. No. 10182-260
EPA Est. No. 10182-AR 1



Contents: 30 gal (114 L)

BEST AVAILABLE COPY

Keep Out
of Reach
of Children

WARNING

See side panel
for Statement of Practical
Treatment and Precaution
ary Statements



Arrosolo® 3-3E
Emulsifiable Liquid Herbicide

STATEMENT OF PRACTICAL TREATMENT

CALL A POISON CONTROL CENTER OR PHYSICIAN IMMEDIATELY if a known exposure occurs or is suspected immediately start the procedures given below and contact a Poison Control Center, physician or the nearest hospital

NOTE: Be sure to advise the person contacted that the product is a weak cholinesterase inhibitor in experimental animals. Describe the type and extent of exposure, the victim's symptoms, and follow the advice given.

NOTE TO MEDICAL PERSONNEL: If cholinesterase inhibition is suspected, atropine by injection is antidotal. It has not been determined if pralidoxime chloride (2-PAM, protropam) is antidotal for the product.

IF SWALLOWED: Immediately give several glasses of water but DO NOT induce vomiting. If vomiting occurs, give fluids again. Have a physician determine if condition of patient will permit induction of vomiting or evacuation of stomach. Do not give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Immediately flush eyes with large amounts of water for at least 15 minutes. Hold eyelids apart during flushing to ensure rinsing of the entire surface of the eye and lids with water. Get medical attention immediately.

IF ON SKIN: Flush affected areas with plenty of water for several minutes. Remove contaminated clothing and shoes. Get medical attention if irritation occurs.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs or breathing becomes difficult.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-A-I-E-D (327-4632)

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

CAUSES SUBSTANTIAL BUT TEMPORARY EYE IRRITATION. Wear goggles or face shield. Harmful if swallowed. Avoid contact with skin, eyes and clothing. Wash with soap and water after use. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water except as specified on this label. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water used for domestic use.

Water drained from treated rice fields must not be used to irrigate other crops or released within 1 mile upstream of a potable water intake (i.e., river, stream, etc.) or within 1 mile of a potable water intake in a standing body of water such as a lake, pond or reservoir.

PHYSICAL AND CHEMICAL HAZARDS

Combustible Liquid

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use the product in a manner inconsistent with the label. Do not apply this product through any type of irrigation system.

GENERAL USE PRECAUTIONS

- Read all label directions before using. Do not overdose.
- ARROSOLO 3-3E emulsifiable liquid herbicide should be used only for purposes and at recommended rates. ARROSOLO 3-3E may cause crop injury on extreme soil or climatic conditions or if directions are not followed.
- Except as specified on this label, the use of fertilizers, oils, surfactants, tank mixes with ARROSOLO 3-3E may result in crop injury and/or phytotoxicity of ARROSOLO 3-3E. A small scale physical compatibility test, as it should be, conducted on all herbicide tank mixes before combining in a tank.
- Avoid spray drift to all other crops and non-target areas.
- Do not apply carbamate or organophosphorus insecticides in combination with ARROSOLO 3-3E application. Severe injury or kill of rice plants may result from combination or separate sprays of ARROSOLO 3-3E and certain insecticides.
- Do not apply ARROSOLO 3-3E when rice is exhibiting stress from verticillium, drought conditions, or injury from previously applied pesticides (diseases).
- ARROSOLO 3-3E should only be applied to fields which have been or will be exposed to susceptible weeds.
- To avoid injury do not apply when the air temperature is above 100°F.
- Weed control may be improved if ARROSOLO 3-3E is applied when the temperature is below 70°F.
- For best results apply ARROSOLO 3-3E to actively growing weeds in rice.
- Do not apply if rainfall is anticipated within 6 hours after application.
- Apply only when wind velocity is low (0-5 mph).
- Do not apply more than 3 pounds of active ingredient of ORDRAM® per acre per season.

GENERAL APPLICATION DIRECTIONS

ARROSOLO 3-3E is a selective broad spectrum postemergence rice herbicide for annual grasses and broadleaf weeds.

All equipment used in the application of ARROSOLO 3-3E should be calibrated before use and checked frequently during application to be sure the application overlaps that will increase ARROSOLO 3-3E dosage above the recommended rate because crop injury may occur.

Apply ARROSOLO 3-3E in 20 to 50 gallons of water per acre by ground or 10 to 15 gallons of water per acre by airplane.

Shake well before using.

DIRECTIONS FOR USE AND RECOMMENDATIONS

SOUTHERN UNITED STATES (Arkansas, Louisiana, Mississippi, Missouri, North Carolina, and Texas)

POSTEMERGENCE, PREFLOOD: For use on dry-seeded and water

**NOTIFICATION
LABEL NOT REVIEWED
PER FR NOTICE 88-4**

DATE: 6/30/93



Arrosolo 3-3E

Emulsifiable Liquid Herbicide

For control of Barnyardgrass, Springbroom, Dayflower and Broadleaf Signalgrass, apply ARROSOLO 3-3E at 3 to 4 quarts per acre depending on weed size as shown in the table below. For best results, apply ARROSOLO 3-3E when weeds are in the 1-2 leaf stage. Flood the rice within 5 days after application to prevent weed reinfestation.

Weeds Controlled by ARROSOLO 3-3E and Rate Chart

Weeds	Leaf Stage	
	3 Quarts	4 Quarts
Barnyardgrass	Echinochloa spp. 1-2	3-4
Crabgrass, large	Digitaria sanguinalis	1-early 4
Crab, waxy	Crabro capitatus	-
Dayflower, common	Commelina communis	1-2
Jack-in-the-box	Rumex crispus	-
Portulaca	Portulaca spp.	1-early 4
Goosegrass	Echinochloa indica	1-early 4
Gulf Cockspur	Echinochloa crus-galli	1-early 4
Hemp, Sesbania or Coffee bean	Sesbania exaltata	-
Hoargrass or Fringedush	Erigeron spp.	1-early 4
Horned Beardless	Rhynchospora corniculata	-
Juncus	Echinochloa colozum	1-early 4
Parasitic	Panicum purpurascens	1-early 4
Pigweed, redroot	Amaranthus retrofractus	-
Rudbeckia	Melicope caroliniana	-
Signalgrass, broadleaf	Brachiaria distachya	1-2
Speargrass	Echinochloa spp.	1-early 4
Springbroom (maximum 1 in high)	Echinochloa spp.	1-2*
Texas Millet	Panicum texanum	1-early 4
Texasweed	Cyperus polystachyus	-

*For control of seeding broadleaf weeds.
**Suppression only in the states of Arkansas, Mississippi, Louisiana and Missouri.

APPENDIX I

Procedure for Testing the Compatibility of ARROSOLO 3-3E in Tankmix Combinations. The following procedure is suggested for determining whether or not ARROSOLO 3-3E may be combined with a specific herbicide for spray tank application.

- Materials Required**
- ARROSOLO 3-3E
 - Tankmix herbicides
 - Adjuvant for spray tankmix combinations, Compent[®], Scento 164-D[®], Unile[®] or equivalent. The adjuvant which provides the best emulsification depends on the specific herbicide under consideration.
 - Two one quart wide mouth glass jars with lids or stoppers
 - Measuring spoons. A 25 mL syringe or graduated cylinder provides more accurate measurement. 1 tablespoon equals 3 teaspoons, equals approximately 5 milliliters.
 - Measuring cup. 8 ounces = 237 mL.

Procedure:

1. Pour a pint or about 473 mL of water into each quart jar.
 2. Add adjuvant to one of the jars and mix. See Table 1 for rate for adjuvant rate.
 3. The order of addition of tankmix herbicides is emulsifiable powders first, followed by wettable dispersible granules, followed by herbicides, followed by ARROSOLO 3-3E. See Table 2 for rates of tankmix herbicides to add to both jars.
 4. If the tankmix herbicide being tested is a wettable powder, premix the wettable powder in a cup of water prior to addition to the pint of water.
 5. Add ARROSOLO 3-3E to both jars. See Table 1 for rate to use.
 6. Close both jars with lid or stopper and mix the contents by turning the jars upside down and back upright ten times.
 7. Inspect the surface and body of the mixtures.
 - Immediately after completing the jar inversions.
 - After allowing the jars to stand undisturbed for 30 minutes.
 - Again after turning the jars upside down and upright again ten times after the 30 minute wait.
- Observations:**
- If a uniform mixture cannot be made, the mixture should not be used.
 - If either mixture remains uniform for 30 minutes, the combination may be used.
 - Should either mixture separate after 30 minutes, but readily re-mixes uniformly with ten additional jar inversions, the mixture can be used if adequate agitation is maintained in the tank.
 - If the mixture with adjuvant is satisfactory, but the one without adjuvant is not, be sure to use the adjuvant in the spray tank.
 - If adjuvant is needed, add to the spray tank first at a rate of 3 pints per 100 gallons of water.
 - Foaming can be minimized by using moderate agitation.
 - If nondispersible or sludge or clumps of solids form in the mixture, the combination should not be used.

TABLE 1
RATE TABLE FOR ARROSOLO 3-3E AND ADJUVANT*

GALLONS OF SPRAY VOLUME TO BE APPLIED PER ACRE	ARROSOLO 3-3E TO ADD TO 1 PINT OF WATER FOR 3 QUART PER ACRE RATE		ARROSOLO 3-3E TO ADD TO 1 PINT OF WATER FOR 4 QUART PER ACRE RATE	
	ML	TEAS.	ML	TEAS.
10	34	7 $\frac{1}{2}$	45	9
15	22 $\frac{1}{2}$	4 $\frac{1}{2}$	31	6 $\frac{1}{2}$
20	17	3 $\frac{1}{2}$	22 $\frac{1}{2}$	4 $\frac{1}{2}$
25	13 $\frac{1}{2}$	3	18 $\frac{1}{2}$	4
30	11 $\frac{1}{2}$	2 $\frac{1}{2}$	15 $\frac{1}{2}$	3 $\frac{1}{2}$
40	8 $\frac{1}{2}$	1 $\frac{1}{2}$	11 $\frac{1}{2}$	2 $\frac{1}{2}$
50	6 $\frac{1}{2}$	1 $\frac{1}{4}$	9	2

*Two (2) milliliters or one-half (1/2) teaspoon of adjuvant added to 1 pint of water equals 3 pints of adjuvant per 100 gallons of water.



Arrosolo 3-3E

Emulsifiable Liquid Herbicide

TABLE 2
RATE TABLE FOR OTHER TANKMIX HERBICIDES

GALLONS OF SPRAY VOLUME TO BE APPLIED PER ACRE	AMOUNT OF DRY HERBICIDE TO ADD FOR EACH POUND TO BE APPLIED PER ACRE		AMOUNT OF LIQUID HERBICIDE TO ADD FOR EACH PINT TO BE APPLIED PER ACRE	
	TEAS.	ML	TEAS.	ML
10	4	1	2	1/2
15	7	1 1/2	3	3/4
20	11	2	4	1
25	14	2 1/2	5	1 1/4
30	17	3	6	1 1/2
40	23	4	8	2
50	30	5	10	2 1/2

STORAGE AND DISPOSAL: Do not use or store near heat or open flame. Do not store near seeds and fertilizer. Keep container closed when not in use. Open dumping is prohibited. Do not dump empty container (for non-bulk or non-bulk containers).

PESTICIDE DISPOSAL: Weeds resulting from the use of the product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Wash, triple rinse (or equivalent) then offer for recycling or reconditioning, or purchase and dispose of in a sanitary landfill or other procedures approved by State and local authorities.

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using the product.

CONDITIONS OF SALE AND LIMITED WARRANTY: The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use of this product (including claims based on contract, negligence, strict liability or other theory) Buyer or User must promptly notify in writing Seller of any claims to be asserted by either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER shall be the LIMIT OF LIABILITY OF Seller shall be the amount of product used. SELLER SHALL BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE REMEDY SHALL BE LIMITED TO THE PURCHASE PRICE.

This product is sold only for uses stated on its label. No express or implied warranty is made for uses other than those stated on the label. Patent No. 3,573,031.

ARROSOLO[®] and ORDAM[®] are trademarks of ICI Group Companies. Compent[™] is a trademark of Kato, Inc. Scento 164-D[™] is a trademark of Waco Corporation. Unile[™] is a trademark of HACO, Inc. ©1982 ICI Americas Inc.

Made in U.S.A.



Agricultural Pro

ICI Americas Inc.
Wilmington, Delaware 19897

BEST AVAILABLE COPY