Plesse read in specions on re			IFICATION Form Approve	11/4/97 od. OMB No. 2070	Page 1 3 5 2-0060. Approval expires 2-28-95				
SEPA	United S Environmental Pro Washington,	tection Age	ncy	on OPP Identifier Number 232871					
	Appl	ication for I	Pesticide - Section		. <u> </u>				
1. Company/Product Number			2. EPA Product Manager		3. Proposed Classification				
10182-280			James Tompkins		None Restricted				
4. Company/Product (Name) GRAMOXONE EXTRA Her	rbicide		PM#						
ZENECA Ag Products 1800 Concord Pike, PO	. Name and Address of Applicant (Include ZIP Code)			6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.					
Check if this	is a new address		Product Name						
		Sec	tion - II						
Amendment - Explain below. Resubmission in response to Agency letter dated			Final printed labels in response to NOTIFICATION Agency letter dated "Me Too" Application. NOV 4 - 1997						
Notification - Explain	-		Other - Explain		NOV 4 - 1997				
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Supplemental labeling for GRAMOXONE EXTRA for harvest aid use in corron, the addition of "For use only in the states of New Mexico, Oklahoma, and Texas. Rounding rates to even pint increments of 0.75 (for 0.8) and 1.5 (for 1.6)									
		Sec	tion - III		 				
1. Material This Product Will	T	1		T					
Yes*	No No No No nor		Water Soluble Packaging Yes No If "Yes" No. per		iteiner letal lastic lass aper				
tification must	Unit Packaging wgt. con	tainer Packa	ge wgt container		ther (Specify)				
3. Location of Net Contents Information 4. Size(s) Retailed Label Container			S. Location of Label Directions On Label On Labeling accompanying product						
6. Manner in Which Label is Affixed to Product Lithograph Other Other Stenciled									
Section - IV									
1. Contact Point Complete	items directly below for idea	ntification of indiv	ridual to be contacted, if n						
Name Deborah Touzell	_	Titte Regulato	ory Information Coord		ephone No. (Include Area Code) 302-886-1226				
•	ments I have made on this f ly knowingly false or mislead		ay be punishable by fine o		6. Date Application Received '''(Stamped)				
2. Signature Linal			tory Information Coor						
			5. Date October 17, 1997						
"PΔ Form 8570-1 (Bay 8.04	l) Previous aditions are obso	Language to the same of the sa	For post,	EDA ETI- O /	riginally Vallage Applicant Conv				

0 95

NOTIFICATION

NOV 4 - 1997

RESTRICTED USE PESTICIDE

Due to Acute Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABEL FOR USE ON COTTON IN THE STATES OF NEW MEXICO, OKLAHOMA AND TEXAS

GRAMOXONE® EXTRA HERBICIDE

EPA Reg. No. 10182-280


)

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Crop	Use Pattern	GRAMOXONE EXTRA Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
COTTON	Harvest			3	Harvest Aid Use Precautions (Applies to all sections) Do not pasture livestock in treated fields or feed treated foliage. Do not apply to cotton within 3 days before harvest. Repeat application if necessary. Do not exceed a total of 1.5 pts. per acre as a harvest aid. May be tankmixed with other cotton harvest aid materials known to be effective by the local expert. Unless otherwise instructed in this label, refer to tankmix product label for rates, directions, limitations and cautions. GRAMOXONE EXTRA can be applied in a tankmix with methylparathion and/or KARATE® insecticide. Nodes above cracked bolls (NACB) timing is for guidance and is not intended to restrict the local expert in their use of the product.

Crop	Use Pattern	GRAMOXONE EXTRA Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
SOUTHERN COTTON Harvest aid for boil opening and defoliation (Tank mix with phosphate and chlorate defoliants)	Broadcast	6.5 fl oz plus 1 pt phosphate or 1 gal chlorate	Ground: 10 gal Air: 5 gal	7	 Development of immature bolls will be inhibited. Apply when 80% or more of the bolls are open and the remaining bolls to be harvested are mature.
SOUTHERN COTTON Additional Tankmixes for boll opening and defoliation	Broadcast	2.5 - 4.0 oz	Ground: 10 gal Air: 5 gal	-	To aid in defoliation and opening of mature bolls, GRAMOXONE EXTRA may be tankmixed with the following products: ACCELERATE® defoliant DEF® defoliant DROPP® defoliant ETHEPHON plant growth regulant FOLEX® defoliant HARVADE® harvest growth regulant PREP™ PGR Apply when 60% or more of the bolls are open and the remaining bolls to be harvested are mature. Development of immature bolls will be inhibited. Refer to tank mix product label for rate, directions, limitations and cautions.
Post Defoliation- To aid in opening of mature bolls and to desiccate green weeds	Broadcast	0.75 - 1.5 pints	Ground: 10 gai Air: 5 gai	3	 Use higher rate if weed infestation is heavy or dense. Apply when 75% or more of the bolls are open and remaining bolls to be harvested are mature. Development of immature bolls will be inhibited. After a defoliation or conditioning application has been made, delay desication application of GRAMOXONE EXTRA approximately 3-7 days to minimize leaf sticking.



Сгор	Use Pattern	GRAMOXONE EXTRA Rate Per Acre	Minimum Total Spray Per Acre	Grazing or Preharvest Interval (Days)	Precautions, Restrictions and Comments
COTTON Stripper or Spindle Harvested Harvest aid for defoliation and boll opening	Broadcast	2.5 - 9 fl oz	Ground: 10 gal Air: 5 gal	3	IT IS ADVISABLE, BECAUSE OF EXTREMES IN ENVIRONMENTAL AND PLANT CONDITIONS, TO APPLY THE RANGE OF RATES ON A SMALL, BLOCK OF COTTON TO DETERMINE THE RATE THAT BEST FITS YOUR NEEDS. Apply when 75% of the bolls are open and the remaining bolls to be harvested are mature. DEVELOPMENT OF IMMATURE BOLLS WILL BE INHIBITED. SLICE BOLLS AND INSPECT THE SEED FOR MATURITY. GRAMOXONE EXTRA may be applied alone or tank mixed with the following cotton harvest aids: ACCELERATE® defoliant DEF® defoliant DROPP® defoliant ETHEPHON plant growth regulator PREP™ PGR May be applied as a split application. Do not exceed a total of 1.5 pts/Acre. To avoid leaf sticking, apply GRAMOXONE EXTRA as a desiccant approximately 3-7 days after defoliation or a conditioning application and 7-14 days before harvest. Cooler temperatures may cause a longer waiting period between application of GRAMOXONE EXTRA as a desiccant and defoliation/condition. Lower rates in the range may be necessary south of 1-10 in Texas where temperatures are typically higher during defoliation.



All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

GRAMOXONE® EXTRA HERBICIDE is a trademark of a ZENECA Group Company.