

SCIMITAR® WP Insecticide

Only for Sale to, Use, and Storage by Commercial Applicators

**For Use to Control Insect Pests On Ornamentals and Lawns in
Landscaped Areas Around Residential, Institutional, Public,
Commercial and Industrial Buildings, Parks, Recreational Areas and Athletic Fields**

ACTIVE INGREDIENT:

Lambda-cyhalothrin

[1 α (S*),3 α (Z)]-(\pm)-cyano(3-phenoxyphenyl)methyl-3-
(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-

dimethylcyclopropanecarboxylate 10.0%

INERT INGREDIENTS¹ 90.0%

Total 100.0%

¹Contains aromatic petroleum distillates.

EPA Reg. No. 10182-300

EPA Est. No.

Net Contents:

KEEP OUT OF REACH OF CHILDREN

WARNING


AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

See Side Panel for Statement of Practical Treatment and Precautionary Statements

BEST AVAILABLE COPY

Made in U.S.A.
ZENECA Professional Products
ZENECA Inc.
Wilmington, DE 19897



STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. If victim is alert and not convulsing, rinse mouth out and give 200-300 ml (1 cup) of water to dilute material. Immediately contact local Poison Control Center. Vomiting should only be induced under the direction of a physician or Poison Control Center. If spontaneous vomiting occurs, have victim lean forward with head down to avoid breathing in of vomitus, rinse mouth and administer more water. Immediately transport victim to an emergency facility.

IF IN EYES: Wash with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably by mouth-to-mouth.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

MAY BE FATAL IF SWALLOWED. HARMFUL IF ABSORBED THROUGH SKIN OR INHALED. CAUSES EYE IRRITATION. **PROLONGED OR FREQUENTLY REPEATED SKIN CONTACT MAY CAUSE ALLERGIC SKIN REACTION IN SOME INDIVIDUALS** Do not get in eyes, on skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.

Wear goggles and rubber gloves when handling or mixing wettable powder.

Skin exposure may also result in a sensation described as a tingling, itching, burning or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 to 30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label. Care should be used when spraying to avoid fish and reptile pets.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Notify State and/or Federal authorities and ZENECA Inc. immediately if you observe any adverse environmental effects due to the use of this product.

BEST AVAILABLE COPY

**CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not reuse empty container.

STORAGE: Mix as needed. Do not store diluted material. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep up and remove to chemical waste storage area until proper disposal can be made.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty jar or bag into application equipment.

Plastic Container: Triple rinse (or equivalent); then puncture and dispose of empty jar in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Paper or Plastic Bag: Dispose of bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

MIXING: SCIMITAR® WP insecticide is intended for dilution with water for spray application and may be used in all types of application equipment. Fill application tank with 1/2 volume of water and add SCIMITAR WP and then fill to desired volume. Close and shake or agitate before use in order to ensure proper suspension of the wettable powder. Shake or reagitate application equipment before use if application is interrupted. Make up only amount of application volume as required.

COMPATIBILITY: SCIMITAR WP has been found to be compatible with all commonly used fungicides, miticides, liquid fertilizers, and other insecticides. Check physical compatibility using the correct proportion of products in a small jar if local experience is unavailable.

APPLICATION: SCIMITAR WP insecticide mixes readily with water and may be used in all types of application equipment. Mix product with the required amount of water and apply as a dilute spray application to the point of runoff. When applying hard-to-wet foliage such as holly, pine, or ivy, the addition of a spreader sticker may enhance knockdown and residual activity. If concentrate or mist-type application equipment is used, an equivalent amount of product should be used on the area treated as would be used in a dilute application. Phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed. It is advised to pretreat a selection of ornamentals if local use experience is unavailable.

SCIMITAR WP may be used for applications to maintain indoor or outdoor areas where turf and ornamentals are grown such as residential landscaped areas and non-residential landscapes around institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

BEST AVAILABLE COPY

IMPORTANT: Time application to flowering plants during periods of time when pollinating insects are not present, such as early morning or late evening.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

DO NOT APPLY THIS PRODUCT TO EDIBLE CROPS.

USE	PEST	AMOUNT OF SCIMITAR WP	REMARKS
ORNAMENTALS IN RESIDENTIAL LANDSCAPED AREAS AND LANDSCAPED AREAS AROUND INSTITUTIONAL PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, AND ATHLETIC FIELDS (Including Trees, Shrubs, Flowers, Evergreens, Foliage Plants and Groundcovers)	Ants (Including Imported fire ants) ² Armyworms Azalea caterpillar Aphids Bagworms ³ Black vine weevil (adult) Boxelder bugs Budworms California oakworm Cankerworms Cockroaches Crickets Cutworms Eastern Tent caterpillar Elm leaf beetles European sawfly Fall webworm Flea beetles Forest tent caterpillar Gypsy moth larvae Japanese beetles (adults) June beetles (adults) Lace bugs Leaf-feeding caterpillars Leafhoppers Leafminers (adults) Leaf rollers Leaf skeletonizers Midges Mosquitoes Oleander moth larvae Pillbug Pine sawfly Pine shoot beetles Pine tip moths Plant bugs Root weevil Sawfly Scale insects (or others) Spiders Spittlebugs Striped beetles Striped oakworm Thrips Tip moth Tussock moth larvae Wasps	1.2 to 4.8 oz/ 100 gals. or 3 to 12 scoops/100 gals. ¹	Applications should be started prior to establishment of high pest population and reapplications made as necessary to keep populations under control, using higher rates as pest pressure increases. Good spray coverage is necessary to provide the most effective level of control. Addition of spreader/sticker at recommended rates may enhance control of insects on certain species of ornamentals having hard-to-wet foliage. Apply at 7-day intervals if retreatment is necessary. Do not apply more than 0.36 lbs of active ingredient per acre per year. Do not apply to any plants grown for food or feed. Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, avoid applying this pesticide within 25 feet of ponds, lakes, streams, estuaries, and marine habitats. NOTE: Certain cultivars may be sensitive to the final solution. A small number of plants should be treated prior to application of entire planting

BEST AVAILABLE COPY

USE	PEST	AMOUNT OF SCIMITAR WP	REMARKS
	Broadmites Brown softscale California redscale (crawler) ¹ Clover mites Mealybugs Pinenacledscale (crawler) ¹ Spider mites Whiteflies	2.4 to 4.8 oz/ 100 gals. or 6 to 12 scoops/100 gals. ¹	
LAWNS AROUND RESIDENTIAL, INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS AND ATHLETIC FIELDS	Ants (Including Imported fire ants) ² Armyworms Centipede Crickets Cutworms Earwigs Fleas (adult) Grasshopper Japanese beetle (Adult) Millipedes Mites Pillbugs Sod webworms Sow bugs Ticks (including species which transmit Lyme disease)	3 to 6 grams/1000 sq. ft. or 4.8 to 9.6 oz/Acre or 12 to 24 scoops/Acre	Apply when pests first appear. Use the 3 grams/1000 sq. ft. rate for light infestations and/or maintenance. Subsurface feeders such as grubs and mole crickets will only be controlled if exposed to the insecticide. The addition of a spread sticker may enhance control. Watering following application (1/4 - 1/2 inch) will also help control. Apply at 7-day intervals if retreatment is necessary. Do not make more than 3 applications per acre per year or apply more than 0.36 lbs active ingredient per acre per year.
	Bluegrass billbug (adult) Black turfgrass ateenius (adult) Chinch bugs Chiggers Fleas (adult) Grubs (suppression) Hyperodes weevil (adult) Mole crickets	6 grams/1000 sq. ft. or 4.8 to 9.6 oz/Acre or 24 scoops/Acre	Do not mow or water for 24 hours for optimum control. Delay watering or mowing for at least 24 hours for optimum control of surface feeding insect pests. Keep children and pets off treated areas until lawn spray has dried following the application. Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, avoid applying this pesticide within 25 feet of ponds, lakes, streams, rivers, estuaries, and other habitats. Do not make applications when wind speed is 15 miles per hour or greater.

1 Scoop = 0.4 oz; 3 grams = 1 teaspoon of SCIMITAR WP.

BEST AVAILABLE COPY

¹For spot treatments with 1 gallon sprayers, use 0.5 to 1 scoop per gallon.

²To control ants effectively, the mound must be treated with a drench application. Use 1 scoop or 0.4 oz of SCIMITAR WP per gallon of water and thoroughly cover and soak each mound ~~and treat a 4-foot diameter circle around the mound.~~ Apply mixture gently to avoid disturbing the mound. Disturbing the mound may disturb the ants and cause migration, thus reducing effectiveness of treatment. For best results apply in cool weather, such as in early morning or late evening hours, but not in the heat of the day. Make additional treatments as needed.

³Bagworm: Apply when larvae begin to hatch. Spray larvae directly. Applications made when larvae are young will be most effective.

⁴Scalecrawlers/twig borers: Treat trunks, stems and twigs in addition to plant foliage.

AMOUNT OF SCIMITAR WP SCOOPS/GALLON(S) OF WATER BASED ON GRAMS OF ACTIVE INGREDIENT PER 1000 SQ. FT.			
RATE	SPRAY VOLUME/GALLONS/1000 SQ. FT.		
	2.0 gals/1000 sq. ft.	3.0 gals/1000 sq. ft.	4.0 gals/1000 sq. ft.
3 grams	12 scoops/91 gals.	6 scoops/68 gals.	6 scoops/91 gals.
6 grams	24 scoops/91 gals.	15 scoops/85 gals.	12 scoops/91 gals.

NOTE: To calculate amount needed per 1000 sq. ft., multiply 0.006 (for 6 gram rate) or 0.003 (for 3 gram rate) by the number of square feet to be sprayed. Divide that figure by 11.3 (grams per scoop) to determine the number of scoops needed.

SCIMITAR® is a trademark of a ZENECA Group Company.

BEST AVAILABLE COPY