

PM03

10182-300

4/24/98

Page 1 of 11  
SCIMITAR WP Container  
SCI300.RSH - 10/15/97**SCIMITAR® WP Insecticide**

For use by individuals/firms licensed or registered by the state to apply insecticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the pest control regulatory agency of your state prior to use of this product.

For Use to Control Insect Pests On Ornamentals and Lawns in  
Landscaped Areas Around Residential, Institutional, Public,  
Commercial and Industrial Buildings, Parks, Recreational Areas and Athletic Fields

**ACTIVE INGREDIENT:**Lambda-cyhalothrin<sup>1</sup>[1 $\alpha$ (S\*),3 $\alpha$ (Z)]-( $\pm$ )-cyano(3-phenoxyphenyl)methyl-3-  
(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-

dimethylcyclopropanecarboxylate ..... 10.0%

**INERT INGREDIENTS<sup>2</sup>** ..... 90.0%**Total** ..... 100.0%<sup>1</sup> Synthetic pyrethroid<sup>2</sup> Contains aromatic petroleum distillates.EPA Reg. No. 10182-300  
EPA Est. No.


Net Contents:

**KEEP OUT OF REACH OF CHILDREN****WARNING****AVISO**

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**See Side Panel for Statement of Practical Treatment and Precautionary Statements**

Made in U.S.A.  
ZENECA Professional Products  
ZENECA Inc.  
Wilmington, DE 19850-5458



211

## STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** If victim is alert and not convulsing, rinse mouth out and give 200-300 mL (1 cup) of water to dilute material. Immediately contact local Poison Control Center. Vomiting should only be induced under the direction of a physician or Poison Control Center. If spontaneous vomiting occurs, have victim lean forward with head down to avoid breathing in of vomitus, rinse mouth and administer more water. Immediately transport victim to an emergency facility.

**IF IN EYES:** Wash with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF INHALED:** Remove victim to fresh air. If not breathing give artificial respiration, preferably by mouth-to-mouth.

---

**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).**

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.**

---

311

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### WARNING

MAY BE FATAL IF SWALLOWED. HARMFUL IF ABSORBED THROUGH SKIN OR INHALED. CAUSES EYE IRRITATION. PROLONGED OR FREQUENTLY REPEATED SKIN CONTACT MAY CAUSE ALLERGIC SKIN REACTION IN SOME INDIVIDUALS. Do not get in eyes, on skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.

Wear goggles and rubber gloves when handling or mixing wettable powder.

Skin exposure may also result in a sensation described as a tingling, itching, burning or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 to 30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

#### ENVIRONMENTAL HAZARDS


This product is extremely toxic to fish. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label. Care should be used when making applications to avoid household pets, particularly fish and reptile pets.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds. In the state of New York, do not apply within 100 feet of coastal marshes or streams that drain into coastal marshes.

Do not make applications when wind speed is 15 miles per hour or greater.

Notify State and/or Federal authorities and ZENECA Inc. immediately if you observe any adverse environmental effects due to the use of this product.



**CONDITIONS OF SALE  
AND LIMITATION OF WARRANTY AND LIABILITY**


**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.



5/11

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**IMPORTANT:** Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns or grounds.

SCIMITAR WP Insecticide may be used for applications to maintain indoor or outdoor areas where turf and ornamentals are grown such as residential landscaped areas and non-residential landscapes around institutional, public, commercial and industrial buildings, parks, recreational areas, golf courses and athletic fields.

**IMPORTANT:** Time application to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

This product is restricted for use in New York state. Only for sale to, use and storage by certified applicators.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

DO NOT APPLY THIS PRODUCT TO EDIBLE CROPS.

DO NOT APPLY THIS PRODUCT BY AERIAL APPLICATION.

---

## STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Do not reuse empty container.

**STORAGE:** Mix as needed. Do not store diluted material. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep up and remove to chemical waste storage area until proper disposal can be made.


**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty jar or bag into application equipment.

**Plastic Container:** Triple rinse (or equivalent); then puncture and dispose of empty jar in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Paper or Plastic Bag:** Dispose of bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

---



**MIXING:** SCIMITAR® WP insecticide is intended for dilution with water for application and may be used in all types of application equipment. Fill application tank with ½ volume of water and add SCIMITAR WP insecticide and then fill to desired volume. Close and shake or agitate before use in order to ensure proper suspension of the wettable powder. Shake or reagitae application equipment before use if application is interrupted. Make up only amount of application volume as required.

**COMPATIBILITY:** SCIMITAR WP Insecticide has been found to be compatible with all commonly used fungicides, miticides, liquid fertilizers, and other insecticides. Check physical compatibility using the correct proportion of products in a small jar if local experience is unavailable.

**APPLICATION:** SCIMITAR WP Insecticide insecticide mixes readily with water and may be used in all types of application equipment. Mix product with the required amount of water and apply as a dilute application to the point of runoff. When applying hard-to-wet foliage such as holly, pine, or ivy, the addition of a spreader sticker may enhance knockdown and residual activity. If concentrate or mist-type application equipment is used, an equivalent amount of product should be used on the area treated as would be used in a dilute application. Phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed. It is advised to pretreat a selection of ornamentals if local use experience is unavailable.

USE	PEST	AMOUNT OF SCIMITAR WP
ORNAMENTALS IN RESIDENTIAL LANDSCAPED AREAS AND LANDSCAPED AREAS AROUND INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, AND ATHLETIC FIELDS (Including Trees, Shrubs, Flowers, Evergreens, Foliage Plants and Groundcovers)	Ants (Including Imported fire ants) <sup>2</sup> Armyworms Azalea caterpillar Aphids Bagworms <sup>3</sup> Black vine weevil (adult) Boxelder bugs Budworms California oakworm Cankerworms Cockroaches Crickets Cutworms Eastern Tent caterpillar Elm leaf beetles European sawfly Fall webworm Flea beetles Forest tent caterpillar Gypsy moth larvae Japanese beetles (adults) June beetles (adults) Lace bugs Leaf-feeding caterpillars Leafhoppers Leafminers (adults) Leaf rollers Leaf skeletonizers Midges Mosquitoes Oleander moth larvae Pillbug Pine sawfly Pine shoot beetles Pinetip moths Plant bugs Root weevil Sawfly Scale insects (crawlers) Spiders Spittlebugs Striped beetles Striped oakworm Thrips Tip moth Tussock moth larvae Wasps	1.2 to 4.8 oz/ 100 gals. or 3 to 12 scoops/100 gals. <sup>1</sup>
	Broadmite Brown softscale California redscale (crawler) <sup>4</sup> Clover mites Mealybugs Pineneedlescale (crawler) <sup>4</sup> Spider mites Whiteflies	2.4 to 4.8 oz/ 100 gals. or 6 to 12 scoops/100 gals. <sup>1</sup>

1 Scoop = 0.4 oz; 3 grams = 1 teaspoon of SCIMITAR WP Insecticide.

8/11

- Application to ornamentals should be started prior to the establishment of high insect pest populations. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Apply at 7-day intervals if treatment is necessary.

Do not apply more than 0.36 pounds of the active ingredient (57.6 oz. of wettable powder) per acre per year.

Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at recommended rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage.

For spot treatments use 0.4 oz. SCIMITAR WP Insecticide per 1 to 2½ gallons of water.

Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.

NOTE: While phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed, certain cultivars may be sensitive to the final spray solution.


It is advised to prespray a selection of ornamental plants and observe them for 7 to 10 days prior to treating large areas if local use experience is unavailable.

**Scale.** Thoroughly cover the plant with SCIMITAR WP Insecticide spray, including trunks, stems, twigs and foliage for control of scale insects.

**Bagworm.** Apply SCIMITAR WP Insecticide when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young

SCIMITAR WP INSECTICIDE MIXING CHART FOR ORNAMENTAL INSECT PEST CONTROL  
(Scimitar WP to add per spray tank)

Rate of Scimitar WP	25 Gallons	50 Gallons	100 Gallons	200 Gallons	300 Gallons
1.2 oz	0.3 oz.	0.6 oz.	1.2 oz.	2.4 oz.	3.6 oz.
2.4 oz.	0.6 oz.	1.2 oz.	2.4 oz.	4.8 oz.	7.2 oz.
4.8 oz.	1.2 oz.	2.4 oz.	4.8 oz.	9.6 oz.	14.4 oz.





USE	PEST	AMOUNT OF SCIMITAR WP
LAWNS AROUND RESIDENTIAL, INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS AND ATHLETIC FIELDS	Ants (including imported fire ants) <sup>2</sup> Armyworms Centipede Crickets Cutworms Earwigs Fleas (adult) Grasshopper Japanese beetle (adult) Millipedes Mites Pillbugs Sod webworms Sow bugs Spittlebugs Ticks (including species which transmit Lyme disease)	3 to 6 grams/1000 sq. ft. or 4.8 to 9.6 oz/Acre or 12 to 24 scoops/Acre
	Bluegrass billbug(adult) Black turfgrass ataenius (adult) Chiggers Fleas (adult) Grubs (suppression) Hyperodes weevil (adult) Mole crickets (nymphs and young adults)	6 grams/1000 sq. ft. or 9.6 oz/Acre or 24 scoops/Acre
	Mole crickets (mature adults) Chinch bugs (Not for use on mature adult mole crickets and chinch bugs in New York state.)	12 grams/1000 sq. ft. or 19.2 oz/Acre or 48 scoops/Acre

19/11

- Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Apply at 7-day intervals if treatment is necessary.

Do not apply more than 0.36 pounds of active ingredient (57.6 oz. of wettable powder) per acre per year.


For spot treatments use 0.4 oz. of SCIMITAR WP Insecticide per 1 to 2½ gallons of water.

Do not apply when turfgrass are waterlogged or when soils are saturated with water (i.e. will not accept irrigation).

Keep children and pets off treated areas until spray has dried following the application.

**Surface Insect Control (armyworm, cutworms, fleas, etc.)** For best results apply SCIMITAR WP Insecticide at recommended rates in 2 to 5 gallons of water per 1,000 sq. ft. The use of a spreader-sticker maybe useful if high rainfall amounts are forecast, otherwise the addition of adjuvants is not necessary under normal conditions for surface insect control in turf. Delay watering or mowing for 12 to 24 hours for optimum control of surface-feeding insect pests.

**Thatch Inhabiting Insect Control (chinch bugs, billbugs, etc.)** For best results apply SCIMITAR WP Insecticide at recommended rates in 2 to 10 gallons of water per 1,000 sq. ft. The use of a non-ionic wetting agent, penetrant or similar adjuvant is recommended at label rates. Lightly irrigate after application with up to ½ inch of water to move the SCIMITAR WP Insecticide into the thatch layer. If irrigation is not available, then use high water application rates for optimum results.



**Subsurface Insect Control (mole crickets, grubs, etc.)** For best results apply SCIMITAR WP Insecticide at recommended rates in 4 to 10 gallons of water per 1,000 sq. ft. The use of a non-ionic wetting agent, penetrant or similar adjuvant is strongly recommended following label rates. Use the highest water application rates possible with your sprayer. Apply SCIMITAR WP Insecticide to turf wet with dew, rain or irrigation. Water-in immediately after application with ¼ to ½ inch of water for optimum results.

**Fire Ant Control** Treat individual mounds with a drench application using a watering can. Use 0.4 oz. of SCIMITAR WP Insecticide per 2½ gallons of water. Thoroughly soak each mound and a 3 foot diameter circle around each mound. Gently apply the mixture to avoid disturbing the mound. Disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. For best results, apply in early morning or late evening hours. Make additional treatments as needed.

**Mosquito Control** Apply as a general spray around landscape plantings, turf, and building foundations to control mosquitoes. For best results apply SCIMITAR WP Insecticide at recommended rates in 2 to 5 gallons of water per 1,000 sq. ft.

SCIMITAR WP Insecticide Mixing Chart for Turf Insect Pest Control  
 (Scimitar WP to add per 100 gallon spray tank)

Rate of Scimitar	Application Rate Per 1,000 Square Feet of Turf				
	2 gal.	4 gal.	6 gal.	8 gal.	10 gal.
4.8 oz./acre	5.7 oz.	2.9 oz.	1.9 oz.	1.4 oz.	1.2 oz.
9.6 oz./acre	11.5 oz.	5.7 oz.	3.8 oz.	2.9 oz.	2.3 oz.
19.2 oz./acre	23.0 oz.	11.5 oz.	7.7 oz.	5.7 oz.	4.6 oz.

Conversion Rate: 1 ounce equals 28.35 grams..

AMOUNT OF SCIMITAR WP SCOOPS/GALLON(S) OF WATER BASED ON GRAMS OF ACTIVE INGREDIENT PER 1000 SQ. FT.			
RATE	SPRAY VOLUME/GALLONS/1000 SQ. FT.		
	2.0 gals/1000 sq.ft.	3.0 gals/1000 sq. ft.	4.0 gals/1000 sq. ft.
3 grams	12 scoops/91 gals	6 scoops/68 gals	6 scoops/91 gals
6 grams	24 scoops/91 gals	15 scoops/85 gals	12 scoops/91 gals
12 grams	48 scoops/91 gals	30 scoops/85 gals	24 scoops/91 gals

NOTE: To calculate amount needed per 1000 sq. ft., multiply 0.006 (for 6 gram rate) or 0.003 (for 3 gram rate) by the number of square feet to be sprayed. Divide that figure by 11.3 (grams per scoop) to determine the number of scoops needed.

SCIMITAR® is a trademark of a ZENECA Group Company.