



TOUCHDOWN® 6
NONSELECTIVE FOLIAR SYSTEMIC HERBICIDE

**FOR WEED CONTROL IN NONCROP AREAS AROUND THE FARM AND
NONBEARING TREES AND VINES**

ACTIVE INGREDIENT

Sulfosate

Trimethylsulfonium carboxymethylaminomethylphosphonate 57.6%

INERT INGREDIENTS 42.4%

Total 100.0%

Contains 6 pounds active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If a known exposure occurs or is suspected, immediately initiate the recommended procedures below. If further treatment is required, contact a POISON CENTER, a physician, or the nearest hospital.

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth if victim is unconscious or convulsing.

IF ON SKIN: Flush with plenty of water for at least 15 minutes. Remove contaminated clothing and shoes. Get medical attention if skin irritation occurs. Wash clothing before re-use.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.


FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL 1-800-F-A-S-T-M-E-D (327-8633).
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

_____ Gal. Net _____ L

EPA Reg. No. 10182-
EPA Est. No.

Made in U.S.A. by
ICI Agricultural Products
ICI Americas Inc.
Wilmington, Delaware 19897

BEST AVAILABLE COPY



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

HARMFUL IF SWALLOWED, ABSORBED THROUGH THE SKIN OR INHALED. PROLONGED CONTACT MAY CAUSE SKIN IRRITATION. MAY CAUSE SKIN REACTIONS. Avoid contact with skin, eyes or clothing. Wear rubber gloves when handling. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before re-use.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds and streams.
Do not apply to any body of water, wetlands or other aquatic habitats. Do not contaminate water when disposing of equipment washwaters.

In case of spill, ISOLATE the spill. Absorb spill with inert absorbent material such as clay or Fuller's earth. Sweep up used absorbent and place in an appropriate chemical waste container. Flush spill area with water. Observe all local, State and Federal laws and regulations regarding disposal, spill, cleanup, removal or discharge.

OTHER PRECAUTIONS

Caution must be taken when applying TOUCHDOWN 6 herbicide to avoid drift or contact with nontarget plant species. Such contact may result in plant injury.

TOUCHDOWN 6 herbicide should not be mixed or stored in galvanized steel or unlined steel containers. Stainless steel, plastic-lined steel, plastic or fiberglass containers are acceptable for mixing and storing TOUCHDOWN 6 herbicide.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product without complying with all label directions and warnings.

- Apply this product only as specified on the label.
- Do not apply by aircraft.
- Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

TOUCHDOWN 6 herbicide is a nonselective systemic herbicide used to control a broad spectrum of emerged weeds. TOUCHDOWN 6 controls most grass and broadleaf weeds – both annual and perennial. TOUCHDOWN 6 may be used in certain noncrop areas around the farm and nonbearing groves, orchards or vineyards up to within 1 year of harvest to control unwanted vegetation.

TOUCHDOWN 6 herbicide is formulated as a liquid concentrate which contains six (6) lbs. of active ingredient per gallon.

TOUCHDOWN 6 herbicide requires a six hour rain-free period after application. Rain occurring within six hours of application may reduce weed control.

APPLICATION DIRECTIONS

Timing

TOUCHDOWN 6 herbicide should be applied to actively growing emerged weeds when they are small. Weeds 6" in height are the easiest to control. Larger weeds may be more difficult to control. The exception to this is Johnsongrass or bermudagrass. These grasses should be allowed to reach the seedhead stage before application.

When weeds have been mowed, thus removing much of the green foliage, allow the weeds to regrow to a height of at least 6 inches before spraying.

Rates

Follow recommended rates for TOUCHDOWN 6 listed in the Weeds Controlled Section. Use the higher label rates when weeds are dense or large. Also, use higher application volumes and pressures when weed vegetation is dense.

BEST AVAILABLE COPY

Tank Mixes

When perennial broadleaf weeds such as Canada thistle, field bindweed, or difficult to control annual broadleaf weeds such as common ragweed or morningglory spp. are present, tank mixes with 2,4-D or Banvel, where labeled, will help improve control. In general, labeled TOUCHDOWN 6 tank mixes with other products should be mixed as follows:

1. Fill spray tank 1/2 full with clean water.
2. Begin tank agitation and continue throughout mixing and spraying.
3. Add dry formulations (WP, DF, etc.) to tank.
4. Add liquid formulations (SC, EC, L, etc.) to tank.
5. Add TOUCHDOWN 6 and nonionic surfactant to tank.
6. Fill remainder of spray tank.

Always refer to labels of other pesticide products for mixing directions and precautions which may differ from those outlined here.

SPRAY ADDITIVES

Surfactants/Wetting Agents

A surfactant or wetting agent (approved for use on growing crops) such as Ortho X-77*, Triton AG-98, Activator 90, or Lo-Dose IS REQUIRED with TOUCHDOWN 6 to improve coverage of weed foliage. All surfactants or wetting agents should contain at least 50% active ingredient. The concentration of active ingredient in a particular surfactant will determine how much should be added to the tank. Refer to table below for specific recommendations:

Surfactant % Active Ingredient	Rate of Surfactant Per 1 Gallon of Water	Rate of Surfactant** Per 100 Gallons of Water
75% or greater	3/4 fl oz	2 quarts*
50-74%	1-1/2 fl oz	4 quarts
Less than 50%	DO NOT USE	DO NOT USE

*2 quarts = 0.5% V/V.

Ammonium Sulfate

Control of annual and perennial weeds with TOUCHDOWN 6 may be improved by adding dry ammonium sulfate at two (2) percent by weight or 17 lbs/100 gallons water in addition to nonionic surfactant. The addition of ammonium sulfate will generally improve control under conditions of adverse environmental stress. Do not reduce use rates of TOUCHDOWN 6 or nonionic surfactant when using ammonium sulfate.

APPLICATION EQUIPMENT & TECHNIQUES

REPRODUCTION OF THIS DOCUMENT IS PROHIBITED

Broadcast Application

Apply TOUCHDOWN 6 to actively growing weeds at the broadcast rates indicated in the Weeds Controlled section. Applications should be made in 10 to 30 gallons of water per acre with conventional spray equipment, or in 3 to 10 gallons per acre with low volume equipment. Increase volumes if weed foliage is dense.

The use of flat-fan nozzles will result in the most effective application of TOUCHDOWN 6. Flat fans should be used at 30-50 psi. Flood nozzles are generally not as good as flat fans since they produce large uneven droplets. The use of flood nozzles may result in reduced weed control due to inadequate coverage.

Spot Treatments

For control of the weeds listed use a 0.25% solution of TOUCHDOWN 6 plus nonionic surfactant in water to weeds less than 6 inches in height or runner length. For annual weeds over 6 inches, use a 0.5% to 1% solution. For best results, use a 1.5% to 2% solution on harder-to-control perennials, such as hemp dogbane, field bindweed, dock, bermudagrass, milkweed, and canada thistle. Refer to the Weeds Controlled section for a list of hard to control weeds. Use a 3% solution when using motorized spot spray equipment (ie. bean buggy or a rider bar). Spray the solution on actively growing foliage until uniformly wet but not to the point of runoff. Re-treat if necessary.

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

BEST AVAILABLE COPY

Spot Spray Recommendations

Amount of TOUCHDOWN 6 Herbicide for Hand-Directed Spot Spray*

Solution Strength	To Make This Volume				
	1 gallon	2 gallons	10 gallons	25 gallons	100 gallons
0.25%	1/3 fl oz	2/3 fl oz	3-1/3 fl oz	8 fl oz	2 pints
0.5%	2/3 fl oz	1-1/3 fl oz	6-2/3 fl oz	1 pint	4 pints
1%	1-1/3 fl oz	2-1/2 fl oz	12-1/2 fl oz	2 pints	1 gallon
1.5%	2 fl oz	4 fl oz	19-1/3 fl oz	3 pints	1-1/2 gallons
2%	2-2/3 fl oz	5-1/3 fl oz	1-1/2 pints	4 pints	2 gallons
3%	4 fl oz	8 fl oz	2-1/2 pints	6 pints	3 gallons
5%	6-1/2 fl oz	13 fl oz	4 pints	10 pints	5 gallons

* See SPRAY APPLICATION directions for surfactants and amounts which can be used.

1/3 fl. oz. = 2 teaspoons
 1/2 fl. oz. = 1 tablespoon
 1 fl. oz. = 2 tablespoons

8 fl. oz. or 1/2 pint = 1 cup
 16 fl. oz. = 1 pint
 1 gallon = 8 pints

Wiper Applications

TOUCHDOWN 6 herbicide may be applied using a wiper or "wick" applicator for selective control or suppression of annual and perennial weeds which become taller than the crop or desirable vegetation. To prevent injury to the crop or desirable vegetation, precautions should be taken to avoid contact with these plants. Equipment should be operated at speeds of 5 mph or less. Use slower speeds where weeds are dense. For improved control, make two applications in opposite directions.

Mixing Directions

Mix 2 quarts of TOUCHDOWN 6 herbicide in 2 gallons of water. Apply this mixture to target weeds, while avoiding contact with desirable vegetation.

BEST AVAILABLE COPY

GENERAL USE PRECAUTIONS

- TOUCHDOWN 6 requires actively growing green plant tissue to function. Application to drought-stressed weeds or weeds with little green foliage (i.e., mowed or cut weeds), may result in reduced weed control.
- DO NOT spray if conditions of thermal inversion exist, or if wind direction and speed may cause spray to drift onto adjacent areas. Drift minimization is the responsibility of the applicator. Consult with local and state agricultural authorities for information regarding avoiding or minimizing spray drift.
- Do not tank-mix TOUCHDOWN 6 herbicide with other pesticides or any other additives except as specified on this label or other supplemental labeling.
- Do not apply TOUCHDOWN 6 herbicide if rain is expected within six (6) hours.
- Thoroughly clean spray tank after each use.

CONSERVATION RESERVE OR FEDERAL SET-ASIDE PROGRAMS

(For use in compliance with the Federal Conservation Reserve or Federal set-aside programs)

- Apply to annual and perennial grass and broadleaf weeds according to the Weeds Controlled Section of this label.
- TOUCHDOWN 6 may be tank-mixed with other herbicides registered for this use.

FARMSTEAD (NONCROP)

- Applications can be made in noncrop areas around the farm such as:

Farmyards	Rights-of-way
Soil Bank Land	Barrier Strips
Farm Buildings	Equipment
Fuel Storage Areas	Fence Rows

- Apply to annual and perennial grass and broadleaf weeds according to the Weeds Controlled Section of this label.
- Avoid contact with the foliage of ornamentals or other desirable plants.
- Repeat applications may be necessary.

1997-12-10 COPY

- Do not make more than one application or exceed a total of 4 lbs (5.3 pints) of TOUCHDOWN 6 herbicide per acre per year.

NONBEARING CROPS

GENERAL INFORMATION

Touchdown 6 herbicide can be used to control annual and perennial weeds in nonbearing groves, orchards or vineyards. Use Touchdown 6 during site preparation prior to planting and up to within 1 year of harvest. Touchdown 6 is recommended on the following non-bearing crops:

Acerola (West Indian Cherry)	Kiwi
Almonds	Lemons
Asparagus	Limes
Avocados	Macadamia
Berries	Mango
Conifers	Olives
Seedling establishment	Oranges
Nursery Beds	Pears
Christmas tree plantings	Pecans
Dates	Pistachios
Figs	Pomegranates
Filberts	Tangelos
Florigraze perennial peanuts	Tangerines
Ginseng	Walnuts
Grapes	
Grapefruit	
Guava	
Johoba	

GENERAL USE PRECAUTIONS

BEST AVAILABLE COPY

Do not apply more than 1-1/2 gallons of Touchdown 6 Herbicide.

Do not allow the spray, spray drift, or mist to contact green foliage or green bark on the trunk, suckers, open wounds, or other green parts of trees and vines.

Contact with any tree or vine part other than mature brown woody bark may cause crop injury.

Avoid contact with stumps as injury to adjacent trees may occur from root grafting.

If annual or perennial weeds have been mowed or grazed, wait for 3-4 inches of new growth to appear before making an application of Touchdown 6 Herbicide. See the "Weeds Controlled"

section for specific rates of application.

DIRECTIONS FOR USE

Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, orchard sprayers, or with wiper application equipment. See the "Application Directions" section of this label for specific information on use of equipment, carriers, and nozzles.

Multiple applications may be necessary to control certain perennial weeds, and weeds originating from underground parts (rhizomes, stolons, tubers) that are initially untreated. Touchdown 6 does not provide residual weed control, and as such, weeds emerging after application will not be controlled. For residual weed control, tankmix Touchdown 6 with residual herbicides as prescribed in the "Tank Mixtures for Nonbearing Crops" section, or make multiple applications.

TANK MIXTURES FOR NONBEARING CROPS

Touchdown 6 can be tankmixed with the following herbicides for control or suppression of emerged annual and perennial weeds the following herbicides provide preemergence control of weeds listed on the respective labels.

Touchdown 6 plus:

- * Goal
- * Karmex
- * Krovar I
- * Krovar II
- * Simazine/Caliber 90
- * Simazine (80W, or 4L, or Caliber 90) plus Surflan (AS, or 75W)
- * Goal plus Surflan (AS, or 75W)
- * Goal plus Simazine (80W, or 4L, or Caliber 90)
- * Goal plus Surflan (AS, or 75W) plus Simazine (80W, or 4L, or Caliber 90)
- * Simazine 4L
- * Simazine 80W
- * Solicam 80WP
- * Surflan AS
- * Surflan 75W

Refer to the individual product labels for precautionary statements, restrictions, and a list of weeds controlled.

For control of emerged weeds in a tankmix:

- * Use Touchdown 6 at 1-1/3 pints (21 fl.oz.) for the control of annual weeds that are less than 12 inches tall. When annual weeds are taller than 12 inches use 2 pints (32 fl.oz.) of Touchdown 6 Herbicide.
- * Use Touchdown 6 at 2-2/3 pints (48 fl.oz.) to 6-2/3 pints (107 fl.oz.) for suppression of perennial

BEST AVAILABLE COPY

weeds. Perennial weeds may require multiple applications for control.
* Add an agriculturally approved non-ionic surfactant when tankmixing Touchdown 6 with residual herbicides. See the "Spray Additives" section of this label for recommendations.

PERENNIAL GRASS SUPPRESSION IN ORCHARD FLOORS BAHIA GRASS

Touchdown 6 herbicide can be used to inhibit seedhead emergence and suppress vegetative growth for approximately 40-50 days with a single application. By using a sequential application, suppression of vegetative growth and inhibition of seedhead emergence can be extended to 120 days. Applications must be made prior to seedhead emergence. Apply Touchdown 6 at 20-30 days after complete green-up or after mowing to 4 inches tall. When a single application is planned, use 4 fluid ounces of Touchdown 6 per acre plus 0.5% to 1.0% v/v nonionic surfactant in 10-25 gallons of water. When a sequential application is planned, use 3 fluid ounces per acre for the first application followed by another application of 3 fluid ounces per acre 40-50 days later.

BERMUDA GRASS

For Suppression Only:

1. East of the Rock Mountains

Apply 4 to 10 fluid ounces plus 0.5 to 1.0% v/v nonionic surfactant in 3-20 gallons of water per acre. Make the application 2 weeks after complete green-up or after 3-4 inches of regrowth following mowing. Use 4 to 6-1/2 fluid ounces per acre if a lesser degree of suppression is desired. A sequential application can be used when regrowth occurs if bermudagrass injury and stand reduction can be tolerated.

BEST AVAILABLE COPY

2. West of the Rocky Mountains

Apply 10 fluid ounces of Touchdown 6 plus 0.5% to 1.0% v/v nonionic surfactant per acre in 3 to 10 gallons of water per acre. Make the application 2 weeks after complete green-up or after 3-4 inches of regrowth following mowing. A sequential application can be used when regrowth occurs if bermudagrass injury and stand reduction can be tolerated.

Partial Control and Burndown:

Touchdown 6 herbicide can be used for burndown and partial control of bermudagrass at 1-1/3 pints (21 fl.oz.) to 2-2/3 pints (42 fl.oz.) plus 0.5 to 1.0% v/v nonionic surfactant in 3 to 20 gallons of water per acre. Use 1-1/3 pints east of the Rocky Mountains and 2-2/3 pints west of the Rocky Mountains.

Use this treatment only if reduction of the bermudagrass stand can be tolerated. Allow at least 21 days for complete burndown.

COOL SEASON GRASS COVERS

For suppression of orchardgrass, fine fescue, tall fescue, and quackgrass apply 5 fluid ounces of Touchdown 6 herbicide. Add a nonionic surfactant at 0.5 to 1.0% v/v in 10 to 20 gallons of water. Add ammonium sulfate at 2% by weight or 17 pounds per 100 gallons of spray mix. For suppression of Kentucky bluegrass, use 4 fluid ounces of Touchdown 6 plus 0.5% to 1.0% v/v nonionic surfactant. Do not add ammonium sulfate for suppression of Kentucky bluegrass.

BEST COPY AVAILABLE

LOW VOLUME APPLICATION (FLORIDA AND TEXAS ONLY)

To control or suppress the perennial weeds listed in the table below, apply the recommended rate of Touchdown 6 plus 0.5% to 1.0% v/v nonionic surfactant in 3-20 gallons of water per acre. Use higher gallonage if weed foliage is dense. Apply when the weeds have 3-4 inches of regrowth after mowing and are actively growing.

Weed Species	Rate of Touchdown 6 (Pints per Acre)			
	1-1/3	2-2/3	4	7 1/4
Bermudagrass	B	NR	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	NR	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	NR	PC	C

S = Suppression B = Burndown
 PC = Partial Control C = Control
 NR = Not Recommended

Weed Species	Type of Grass Response*			
	S	B	PC	C
Bermudagrass	.5	1.3	4	6
Guineagrass				
Texas and Florida Ridge	NR	1.3	NR	2.6
Florida Flatwoods	NR	2.6	NR	4
Paragrass	NR	1.3	NR	2.6
Torpedograss	1.3	NR	4	6

*Type of Grass Response

S = Suppression PC = Partial Control
 B = Burndown C = Control

BEST AVAILABLE COPY

Page 13 of 16
TD 6 Nonbearing
RS-121990B

WEEDS CONTROLLED

All weeds listed below are controlled by spray and hand-directed spot applications of TOUCHDOWN 6. For spray application use the indicated rates. See DIRECTIONS FOR USE for mixing and application instructions.

TABLE 1

The following weeds are controlled by 1/2 to 1-1/3 pints (0.375 - 1.0 lbs ai) TOUCHDOWN 6 per acre:

COMMON NAME	SCIENTIFIC NAME
Bluegrass, Annual	(<i>Poa annua</i>)
Chickweed	(<i>Stellaria media</i>)
Cupgrass, Southwestern	(<i>Eriochloa gracilis</i>)
Foxtail, Giant	(<i>Setaria faberii</i>)
Foxtail, Green	(<i>Setaria viridis</i>)
Jimsonweed	(<i>Datura stramonium</i>)
Kochia	(<i>Kochia scoparia</i>)
Oats, Wild	(<i>Avena fatua</i>)
Puncturevine	(<i>Tribulus terrestris</i>)
Ryegrass, Italian (Annual)	(<i>Lolium multiflorum</i>)
Signalgrass	(<i>Brachiaria spp.</i>)
Sowthistle	(<i>Sonchus spp.</i>)
Spurge, Spotted	(<i>Euphorbia maculata</i>)
Thistle, Russian	(<i>Salsola kali</i>)
Wheat, Volunteer	(<i>Triticum spp.</i>)
Witchgrass	(<i>Panicum capillare</i>)

BEST COPY AVAILABLE

BEST AVAILABLE COPYPage 14 of 16
TD 6 Nonbearing
RS-121990B**TABLE 2**

The following weeds are controlled by 1 to 2-1/4 pints (0.7 - 1.7 lbs a) TOUCHDOWN 6 per acre:

COMMON NAME	SCIENTIFIC NAME
Alfalfa, Volunteer	(Medicago sativa)
Barley, Little	(Hordeum pusillum)
Barnyardgrass	(Echinochloa crus-galli)
Corn, Volunteer	(Zea mays)
Crabgrass, Large	(Digitaria sanguinalis)
Eveningprimrose, Cutleaf	(Oenothera lacinolata)
Goosegrass	(Eleusine Indica)
Lambsquarters, Common	(Chenopodium album)
Mustard	(Brassica spp.)
Nightshade, Black	(Solanum nigrum)
Panic Grass	(Panicum spp.)
Pigweed, Redroot	(Amaranthus retroflexus)
Pineappleweed	(Matricaria matricarioides)
Quackgrass	(Agropyron repens)
Pigweed, Redroot	(Amaranthus retroflexus)
Ryegrass, Perennial	(Lolium perenne)
Sicklepod	(Cassia obtusifolia)
Smartweed, Pennsylvania	(Polygnum pennsylvanicum)
Timothy	(Phleum pratense)

TABLE 3

The following weeds are controlled by 1-1/3 to 2-2/3 pints (1 - 2.0 lbs a) TOUCHDOWN 6 per acre:

COMMON NAME	SCIENTIFIC NAME
Brome, Downy	(Bromus tectorum)
Cocklebur	(Xanthium spp.)
Dogbane, Hemp	(Apocynum cannabinum)
Hophornbeam, Copperleaf	(Acalypha ostryaefolia)
Johnsongrass, Rhizome	(Sorghum halepense)
Johnsongrass, Seedling	(Sorghum halepense)
Milkweed	(Asclepias spp.)
Purslane, Common	(Portulaca oleracea)
Radish, Wild	(Raphanus raphistrum)
Ragweed, Giant	(Ambrosia trifida)
Shepherdspurse	(Capella bursa-pastoris)

** To control regrowth of rhizome johnsongrass, apply TOUCHDOWN 6 at or near seedhead stage or later.

Page 15 of 16
TD 6 Nonbearing
RS-121990B

TABLE 4

The following weeds are controlled by 1.9 to 4 pints (1.4 - 3.0 lbs ai) TOUCHDOWN 6 per acre:

COMMON NAME	SCIENTIFIC NAME
Morningglory	(Ipomea spp.)
Ragweed, Common	(Ambrosia artemisiifolia)
Sida, Prickly	(Sida spinoza)
Thistle, Canada	(Cirsium arvense)

TABLE 5

The following weeds are controlled by 2-3/4 to 5-1/3 pints (2.1 - 4 lbs ai) TOUCHDOWN 6 per acre:

COMMON NAME	SCIENTIFIC NAME
Bermudagrass*	(Cynodon dactylon)
Bindweed, Field*	(Convolvulus arvensis)
Cogongrass	(Imperata cylindrica)
Fescue, Tall	(Fescue arundinacea)
Nutsedge, Yellow	(Cyperus esculentus)
Sesbania, Hemp	(Sesbania exaltata)
Velvetleaf	(Abutilon theophrasti)

- * A second application may be needed to control bermudagrass. To control regrowth, apply TOUCHDOWN 6 at or near seedhead stage or later.
- ** A second application may be needed to control field bindweed. To control regrowth, apply TOUCHDOWN 6 at or near flowering stage.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL DIRECTIONS

PROHIBITIONS: Do not contaminate water, food, feed, fertilizer or seeds by storage or disposal.

STORAGE: Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Rinse spray equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Plastic - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Metal - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or disposal in a sanitary landfill, or by other approved State and local procedures.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States of America. U.S. Patent No. 4,315,765.

TOU000B.RSB

8-57 121990B 1087