



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

246334

Application for Pesticide - Section I

1. Company/Product Number 10182-324	2. EPA Product Manager Robert J. Taylor	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) TOUCHDOWN Herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Zeneca Agricultural Products P.O. Box 15458 Wilmington, DE 19850-5458 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Please see cover letter for details of notification.

NOTIFICATION
JAN 2 1997

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 1 QT, 2.5, 30 and 110 gal.		5. Location of Label Directions <input type="checkbox"/> On Label <input checked="" type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Nancy Bockrath	Title Regulatory Assistant	Telephone No. (include Area Code) 302-886-1066
2. Signature 		6. Date Application Received (Stamped)
3. Title Regulatory Assistant		
4. Typed Name Nancy Bockrath		
5. Date December 16, 1996		

ZENECA
FEDERAL EXPRESS

December 17, 1996

Zeneca Ag Products
P.O. Box 15458
Wilmington, DE 19350-5458

Telephone (302) 886-1000
Fax (302) 886-1553

2/22

Mr. Robert J. Taylor
Product Manager Team (25)
Fungicide-Herbicide Branch
Registration Division (H7505C)
Office of Pesticide Programs
Document Processing Desk
U.S. Environmental Protection Agency
Crystal Mall 2, Room 266A
1921 Jefferson Davis Highway
Arlington, VA 22202

Dear Mr. Taylor:

RE: TOUCHDOWN® Herbicide
EPA Registration Number 10182-324
NOTIFICATION
Final Printed Labeling

Zeneca hereby submits one copy of final printed labeling for TOUCHDOWN Herbicide, EPA Reg. No. 10182-324. Label components include container labeling for a 1 quart, 2.5 gallon, 30 gallon and 110-gallon container; and a booklet.

Please be advised that this labeling is being submitted as a NOTIFICATION under the provisions of 40 CFR, Part 152.46(a) as the following changes were made and/or revisions incorporated:

- > The only crop added at this time is soybeans approved in the agency's letter dated March 27, 1996. This incorporates final printed supplemental labeling submitted to EPA on March 28, 1996 for use in soybeans (label code 09-3609-003). No other approved crops are being added at this time.
- > Addition of aerial application and revision in weed tables, approved by the Agency on October 1, 1996. The Fallowland use also approved October 1, 1996 will continue to be handled by Supplemental labeling submitted to EPA on October 21, 1996 (label code 09-2609-010)
- > The surfactant rate range approved in the Agency letter of 7/10/96 will continue to be handled by supplemental labeling submitted to EPA on August 6, 1996 (No label code) and has not been incorporated into this labeling.
- > Other minor wording changes incorporated for clarity are:



LISTENING • LEARNING • LEADING

A business unit of Zeneca Inc.

3/22

- removal of word "groves" after citrus throughout the label
- adding "/wetting agent" after surfactant in the "Air" section. Approved this way throughout rest of label.
- omitted word "crop" before canopy in Air section
- added "as a combination of broadcast, spot, and wiper applications" in second paragraph of Application Directions for Bearing Citrus and Nonbearing crops to clarify the rate is a total maximum by all application methods.

Should there be any questions concerning this submission, or specific questions on the labeling, please contact me at 423-982-8379.

Sincerely,

Becky Rhodes /llw

Becky Rhodes
Regulatory Product Manager

BR/llw

12179601.ber



ZENECA Ag Products



**NONSELECTIVE FOLIAR SYSTEMIC
HERBICIDE FOR WEED CONTROL**

**COMPLETE
DIRECTIONS FOR USE**

04004234
029604 Rev. 11/96

**CONDITIONS OF SALE AND
LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks, referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIA-

NOTIFICATION

JAN 21 1997




BILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America. U.S. Patent No. 4,315,765.

TOUCHDOWN[®], DEVRINOL[®], FUSION[®], SURPASS[®], TOPNOTCH[™] and WARRIOR[™] are trademarks of a ZENECA Group Company. Banvel[®], Frontier[®], Guardsman[®], Marksman[®], and Solicam[®] are trademarks of Sandoz Crop Protection Corporation. Bicep[®], Bicep II[®], Caliber[®], Dual[®] and Dual II[®] are trademarks of Ciba. Bladex[®], Canopy[®], Extrazine[®], Gemini[®], Karmex[®], Krovar[®], Loxone[®], Lorox[®] and Preview[®] are trademarks of E. I. du Pont de Nemours & Co., Inc. Broadstrike[®] and Surlflan[®] are trademarks of Dow-Elanco. Bullet[®], Harness[®], Lariat[®], Lasso[®], Micro-Tech[®] and Partner[®] are trademarks of Monsanto Company. Command[®] is a trademark of FMC Corporation. Direx[®] and Linex[®] are trademarks of Griffen Corporation. Goal[®] is a trademark of Rohm and Haas Company. Prowl[®], Pursuit[®], Scepter[®], and Squadron[®] are trademarks of American Cyanamid Company. Senor[®] and Turbo[®] are trademarks of Miles. Sim Trol[®] is a trademark of Oxon Italia.



4/00

TABLE OF CONTENTS

	Page
STATEMENT OF PRACTICAL TREATMENT	5
PRECAUTIONARY STATEMENTS	6
Personal Protective Equipment	6
User Safety Recommendations	6
Environmental Hazards	6
Other Precautions	7
DIRECTIONS FOR USE	7
AGRICULTURAL USE REQUIREMENTS	8
STORAGE AND DISPOSAL	9
GENERAL INFORMATION	10
APPLICATION DIRECTIONS	10
Timing	10
Rates	10
Tankmixes with Residual Herbicides	10
Spray Additives	12
Surfactants/Wetting Agents	12
Ammonium Sulfate	12
APPLICATION EQUIPMENT & TECHNIQUES	12
Ground	12
Air	13
Spot Treatment	14
Wiper Application	17
GENERAL USE PRECAUTIONS	17
CONSERVATION COMPLIANCE	18
POST HARVEST USE	18
CORN (Field, Popcorn, and Seed Corn)	18
General Use Precautions	18
Tank Mixtures for Corn	19
SOYBEANS	19
General Use Precautions	20
Tank Mixtures for Soybeans	20

	Page
CITRUS (BEARING)	21
NONBEARING CROPS	21
APPLICATION DIRECTIONS FOR BEARING CITRUS AND NONBEARING CROPS	22
TANK MIXTURES FOR BEARING CITRUS AND NONBEARING CROPS	22
PERENNIAL GRASS SUPPRESSION IN ORCHARD FLOORS	24
Bahagrass	24
Bermudagrass	24
Cool Season Grass Covers	25
LOW VOLUME APPLICATION (FLORIDA AND TEXAS ONLY)	25
GENERAL USE PRECAUTIONS FOR BEARING CITRUS AND NONBEARING CROPS	27
WEEDS CONTROLLED	27
Annual Weed Control	28
Perennial Weed Control	35
NONAGRICULTURAL USE REQUIREMENTS	43
Farmstead (Noncrop)	43

KEEP OUT OF REACH OF CHILDREN
CAUTION

ACTIVE INGREDIENT:

Sulfonium, trimethyl-salt with N-(phosphonomethyl)glycine (1:1)*	57.6%
INERT INGREDIENTS:	42.4%
Total	100.0%

*Sulfosate

Contains 6 pounds active ingredient per gallon.
EPA Reg. No. 10182-324STATEMENT OF
PRACTICAL TREATMENT

If a known exposure occurs or is suspected, immediately initiate the recommended procedures below. If further treatment is required, contact a POISON CONTROL CENTER, a physician, or the nearest hospital.

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth if victim is unconscious or convulsing.

IF ON SKIN: Flush with plenty of water for at least 15 minutes. Get medical attention if skin irritation occurs.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

6

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, absorbed through the skin, or inhaled. Prolonged contact may cause skin irritation. May cause allergic skin reactions. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds and streams. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

In case of spill, ISOLATE the spill. Absorb spill with inert absorbent material such as clay or Fuller's earth. Sweep up used absorbent and place in an appropriate chemical waste container. Flush spill area with water. Observe all local, State and Federal laws and regulations regarding disposal, spill, cleanup, removal or discharge.

OTHER PRECAUTIONS

Caution must be taken when applying TOUCHDOWN nonselective foliar systemic herbicide to avoid drift or contact with nontarget plant species. Such contact may result in plant injury.

Stainless steel, plastic-lined steel, plastic or fiberglass containers are acceptable for mixing and storing TOUCHDOWN. TOUCHDOWN should not be mixed or stored in galvanized steel or unlined steel containers. TOUCHDOWN is not compatible with carbon steel and neoprene or butyl rubber.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Apply this product only as specified on the label.
- Do not apply this product through any type of irrigation system.
- Do not exceed a total of 5% pints of TOUCHDOWN per acre per year.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

7

8

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

6/22

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Rinse spray equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Plastic—Triple rinse (or equivalent). Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

FOR RECYCLABLE/REFILLABLE CONTAINERS

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

10

GENERAL INFORMATION

TOUCHDOWN is a nonselective foliar systemic herbicide used to control a broad spectrum of emerged grass and broadleaf weeds, both annual and perennial, in

- conservation compliance;
- corn;
- soybeans;
- bearing orange, grapefruit, lemon, lime, citron, kumquat, tangelo, and tangerine;
- nonbearing groves, orchards, or vineyards up to 1 year of harvest; and
- certain noncrop areas around the farm

TOUCHDOWN is formulated as a liquid concentrate containing 6 pounds active ingredient per gallon.

Rainfall or irrigation within 4 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours may require retreatment.

APPLICATION DIRECTIONS**Timing**

TOUCHDOWN should be applied to actively growing emerged weeds. Annual weeds of 6 inches or less in height are typically the easiest to control. Generally, perennial weeds should be treated at the flowering or seedhead stage.

When annual weeds have been mowed or grazed, wait for 3-4 inches of new growth to appear prior to application. When perennial weeds have been mowed or grazed, allow new growth to reach the recommended stage prior to application.

Rates

Follow recommended rates for TOUCHDOWN listed in the WEEDS CONTROLLED section. Use the higher label rates when weeds are dense or large. Use higher application volumes and pressures when weed vegetation is dense.

Tankmixes with Residual Herbicides

Refer to crop sections for recommended tankmixes. It is recommended that the compatibility of any tankmix combination be tested on a small scale such as a jar test before actual tank

mixing. Tankmixes of TOUCHDOWN with other pesticides, fertilizers, or any other additives except as specified on this label or other approved Zeneca supplemental labeling may result in tankmix incompatibility or unsatisfactory performance.

For Control of Annual Weeds with a Tankmix:

Apply TOUCHDOWN at 1½ pints for the control of annual weeds that are less than 12 inches tall and actively growing. When annual weeds are taller than 12 inches or under stress, use 2 pints of TOUCHDOWN herbicide.

For Control of Perennial Weeds with a Tankmix:

Apply TOUCHDOWN at 2 to 5½ pints for the control or suppression of perennial weeds. Refer to the WEEDS CONTROLLED section for application rates and timing for specific perennial weeds. Perennial weeds may require multiple applications for control. If multiple applications are required, then no more than 5½ pints of TOUCHDOWN should be applied per year.

Add an agriculturally approved nonionic surfactant/wetting agent when tank mixing TOUCHDOWN with residual herbicides. See the SPRAY ADDITIVES section of this label for recommendations.

In general, TOUCHDOWN should be tank mixed as follows:

1. Fill spray tank ½ full with clean water.
2. Begin tank agitation and continue throughout mixing and spraying.
3. Add dry formulations (WP, DF, etc.) to tank.
4. Add liquid formulations (SC, EC, L, etc.) to tank.
5. Add TOUCHDOWN and nonionic surfactant/wetting agent to tank.
6. Fill remainder of spray tank.

Always refer to labels of other pesticide products for mixing directions and precautions which may differ from those outlined here. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

9

11

7/92

SPRAY ADDITIVES

Surfactants/Wetting Agents

A nonionic surfactant or wetting agent (approved for use on growing crops) IS REQUIRED with TOUCHDOWN to improve coverage of weed foliage. All nonionic surfactants or wetting agents should contain at least 50% active ingredient unless otherwise indicated. The concentration of active ingredient in a particular surfactant/wetting agent will determine how much should be added to the tank. Refer to table below for specific recommendations:

Surfactant/ Wetting Agent % Active Ingredient	Rate of Surfactant/ Wetting Agent Per 1 Gallon of Water	Rate of Surfactant/ Wetting Agent Per 100 Gallons of Water
75% or greater	¼ fl oz	2 quarts*
50% - 74%	1½ fl oz	4 quarts
Less than 50%	DO NOT USE	DO NOT USE

*2 quarts = 0.5% V/V

Ammonium Sulfate (Not for use in California)
Control of annual and perennial weeds with TOUCHDOWN may be improved by adding dry ammonium sulfate 1-2% by weight or 8.5-17 pounds per 100 gallons water in addition to nonionic surfactant/wetting agent. Use ammonium sulfate in conservation compliance programs, in noncrop areas, in nonbearing crops, in citrus, and for preplant/preemergence applications. DO NOT reduce use rates of TOUCHDOWN or nonionic surfactant/wetting agent when using ammonium sulfate.

APPLICATION EQUIPMENT & TECHNIQUES

Broadcast Applications

Ground

Apply TOUCHDOWN to actively growing weeds at the broadcast rates indicated in the WEEDS CONTROLLED section. Applications should be made in 10 to 30 gallons of water per

acre with conventional spray equipment, or in 3 to 10 gallons per acre with low volume equipment. Increase volumes if weed foliage is dense.

Flat-fan nozzles will result in the most effective application of TOUCHDOWN. Flat fans should be used at 30 to 50 psi. Flood nozzles may result in reduced weed control due to inadequate coverage.

Air

Apply TOUCHDOWN plus surfactant/wetting agent to actively growing weeds by air equipment in a minimum of 3 gallons of water per acre. Refer to the WEEDS CONTROLLED Section for rates and timing of application.

All aerial equipment must be properly maintained and washed to remove product residues after use.

Spray should be released at the lowest height consistent with weed control and flight safety. Applications more than 10 feet above the canopy should be avoided.

Use the largest droplet size consistent with good weed control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Drift control additives may be used with TOUCHDOWN. Read and follow the manufacturer's directions for use.

Do not make aerial applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Do not apply when winds exceed 15 MPH or there are wind gusts as this could cause drift to adjacent nontarget vegetation.

Spot Treatment

For annual weeds less than 6 inches, use a 0.5% solution (TOUCHDOWN plus nonionic surfactant/wetting agent in water). For annual weeds over 6 inches, use a 1% solution. Use a 2% solution for most perennials except as noted in the SPOT TREATMENT OF HARD TO CONTROL WEEDS table. When using motorized spot spray equipment (rider bar), use a 3% solution. Spray the solution on actively growing weeds until uniformly wet but not to the point of runoff. Retreat 14-21 days later if regrowth occurs.

8/22

9/22

Wiper Application
 TOUCHDOWN may be applied using a wiper or "wick" applicator for selective control, or suppression of annual and perennial weeds which become taller than the crop or desirable vegetation. Mix 1 quart of TOUCHDOWN in 1 gallon of water. Add nonionic surfactant/wetting agent at 1% v/v. Precautions should be taken to avoid contact with crops or desirable vegetation. Equipment should be operated at speeds of 5 mph or less. Use slower speeds where weeds are dense. For improved control, make two applications in opposite directions. Do not use wiper applications in corn.

GENERAL USE PRECAUTIONS

- TOUCHDOWN requires actively growing green plant tissue to function. Application to drought-stressed weeds or weeds with little green foliage (i.e., mowed or cut weeds), may result in reduced weed control.
- Do not till or mow within 3 days following application.
- DO NOT spray if conditions of thermal inversion exist, or if wind direction and speed may cause spray to drift onto adjacent areas. Drift minimization is the responsibility of the applicator. Consult with local and State agricultural authorities for information regarding avoiding or minimizing spray drift.
- It is recommended that the spray system be thoroughly cleaned with water and a commercial tank cleaner after each use.
- Rotational crops may be planted back into TOUCHDOWN treated areas 35 days after application.
- Do not exceed a total of 5% pints of TOUCHDOWN per acre per year.
- Do not graze or harvest treated cover crops for feed.

17

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

TOUCHDOWN Herbicide Spot Spray Dilution Table*

Solution Strength	To Make This Volume				
	1 gallon	2 gallons	10 gallons	25 gallons	100 gallons
0.5%	1/2 fl oz	1 1/2 fl oz	6 1/2 fl oz	1 pint	4 pints
1%	1 1/2 fl oz	2 1/2 fl oz	12 1/2 fl oz	2 pints	1 gallon
2%	2 1/2 fl oz	5 1/2 fl oz	1 1/2 pints	4 pints	2 gallons
3%	4 fl oz	8 fl oz	2 1/2 pints	8 pints	3 gallons

*See SPRAY ADDITIVES directions for surfactants/wetting agents and amounts which can be used.
 1/2 fl. oz. = 2 teaspoons 8 fl. oz. or 1/2 pint = 1 cup
 1 fl. oz. = 1 tablespoon 16 fl. oz. = 1 pint
 1 fl. oz. = 2 tablespoons 1 gallon = 8 pints

SPOT TREATMENT OF HARD TO CONTROL WEEDS

WEED SPECIES	SCIENTIFIC NAMES	HANDGUN % SOLUTION	APPLICATION TIMING AND REMARKS
Balsamapple*	<i>Momordica charantia</i>	1%	Actively growing
Guineagrass*	<i>Panicum maximum</i>	1%	Actively growing, 7-leaf stage
Lantana-largeleaf*	<i>Lantana camara</i>	1%	Actively growing, at bloom
Paragrass*	<i>Brachiaria mutica</i>	1%	Actively growing, seedhead
Pampasgrass*	<i>Erianthus ravennae</i>	2%	Actively growing, at or beyond boot
Sweet Potato, Wild*	<i>Ipomoea pandurata</i>	2%	Actively growing, at or beyond bloom
Torpedograss*	<i>Panicum repens</i>	1%	Actively growing at or beyond seedhead
Vaseygrass*	<i>Paspalum urvillei</i>	1%	Actively growing, at seed-head

*Repeat treatment may be required to provide control.

16

18

18

CONSERVATION COMPLIANCE

(For use in compliance with the Federal Conservation Compliance Programs)

- Apply to annual and perennial grass and broadleaf weeds according to the WEEDS CONTROLLED section of this label.
- TOUCHDOWN may be tank mixed with other herbicides registered for this use.

POST HARVEST USE

TOUCHDOWN may be applied to fields following harvest to control annual and perennial weeds. Refer to WEEDS CONTROLLED Section for use rates. Repeat applications may be necessary to control weeds emerging after application. Use the higher rate on heavy or sodded infestations. Allow sufficient time for regrowth to occur after harvest prior to application. Avoid application after plants have been exposed to severe frost.

Rotational crops may be planted back into TOUCHDOWN treated areas 35 days after application.

CORN (Field Corn, Popcorn, and Seed Corn)**MINIMUM OR NO-TILLAGE SYSTEMS**

TOUCHDOWN is a nonselective systemic herbicide for use as a broadcast spray to control emerged annual and perennial weeds in fields planted to minimum or no-till corn. Apply as a broadcast application before, during, or after planting, but before crop emergence. TOUCHDOWN can also be applied to corn as a spot spray.

Follow the directions listed in the APPLICATION DIRECTIONS, SPRAY ADDITIVES, and APPLICATION EQUIPMENT & TECHNIQUES sections. Refer to the WEEDS CONTROLLED section for rates.

GENERAL USE PRECAUTIONS FOR CORN

- Do not apply within 90 days of harvest of corn grain or corn fodder.
- Do not use on sweet corn.
- Do not apply more than 5½ pints per acre per year as a combination of broadcast sprays prior to crop emergence and as spot spray applications.

TANK MIXTURES FOR CORN

TOUCHDOWN can be tank mixed with the following pesticides. Several of the herbicides listed provide preemergence control or suppression of annual weeds. WARRIOR™ is an insecticide which can be tank mixed with TOUCHDOWN for control of labeled pests.

TOUCHDOWN plus:

Atrazine	Lariat*
Banvel*	Lasso*
Blecp*	Linex*
Blecp II*	Lorox*
Bladex*	Marksman*
Broadstrike*	Micro-Tach*
Bullet*	Prowl*
Dual*	Simazine
Dual II*	SURPASS* EC
Extrazine*	SURPASS 100
Frontier*	TOPNOTCH™
Guardsman*	WARRIOR
Harness*	2,4D
Harness Xtra	

Refer to individual product labels for precautionary statements, restrictions, rates, and a list of weeds controlled. Refer to the WEEDS CONTROLLED section for rates of TOUCHDOWN to use in a tankmix.

SOYBEANS**MINIMUM OR NO-TILLAGE SYSTEMS**

TOUCHDOWN is a nonselective systemic herbicide for use as a broadcast spray to control emerged annual and perennial weeds in minimum or no-till planted soybeans. Apply as a broadcast application before, during or after planting, but before crop emergence. TOUCHDOWN can be applied to soybeans as a spot spray and by wiper or wick. Follow directions listed in the APPLICATION DIRECTIONS, SPRAY ADDITIVES, AND APPLICATION EQUIPMENT & TECHNIQUES sections. Refer to WEEDS CONTROLLED section for rates.

19

20

GENERAL USE PRECAUTIONS FOR SOYBEANS

- Spot application must be made at least 8 weeks before harvest.
- Wiper or wick application must be made at least 7 days before harvest.
- Avoid contact with soybean foliage.
- Do not apply more than 5½ pints per acre per year as a combination of broadcast sprays prior to crop emergence and as spot spray and wiper applications.

TANK MIXTURES FOR SOYBEANS

TOUCHDOWN can be tank mixed with the following pesticides. Several of the herbicides listed provide preemergence control or suppression of annual weeds. WARRIOR™ is an insecticide which can be tank mixed with TOUCHDOWN for control of labeled pests.

TOUCHDOWN plus:

2,4-D	Lorox*
Canopy*	Lorox Plus
Command*	Partner*
Dual*	Preview*
Dual II	Prowl*
Frontier*	Pursuit* Plus
FUSION*	Scepter*
Gemini*	Sencor*
Lasso*	Squadron*
Lexone*	Turbo*
Linex*	WARRIOR

Refer to individual product labels for precautionary statements, restrictions, rates, and a list of weeds controlled. Refer to the WEEDS CONTROLLED tables for rates of TOUCHDOWN to use in a tankmix.

10/20

CITRUS (BEARING) AND NONBEARING CROPS

TOUCHDOWN can be used to control annual and perennial weeds in the following citrus crops and in the following nonbearing groves, orchards, and vineyards:

CITRUS (BEARING)

Citrus	Lime
Grapefruit	Orange
Kumquat	Tangelo
Lemon	Tangerine

NONBEARING CROPS*

Acerola	Huckleberry
(West Indian Cherry)	Jojoba
Almonds	Kiwi
Apple	Lemons
Apricots	Limes
Artichokes	Loganberry
Asparagus	Macadamia
Avocados	Mango
Bananas	Nectarines
Blackberry	Olives
Blueberry	Oranges
Boysenberry	Papayas
Cherries	Passion Fruit
Coffee	Peaches
Crabapple	Pears
Cranberry	Pecans
Dates	Persimmons
Dewberry	Pistachios
Elderberry	Plantains
Figs	Plums
Filberts	Pomegranates
Ginseng	Prunes
Gooseberry	Raspberry
Grapes	Tangelos
Grapefruit	Tangerines
Guava	Walnuts

*TOUCHDOWN can be used during site preparation prior to planting and up to 1 year of harvest.

APPLICATION DIRECTIONS FOR BEARING CITRUS AND NONBEARING CROPS

Applications may be made with boom equipment, shielded sprayers, hand-held and high-volume wands, lances, orchard sprayers, or with wiper application equipment. Follow directions listed in the APPLICATION DIRECTIONS, SPRAY ADDITIVES, and APPLICATION EQUIPMENT & TECHNIQUES sections of this label. Refer to the WEEDS CONTROLLED section for application rates and timing. For hard to control weeds, refer to the SPOT TREATMENT section.

Multiple applications may be necessary to control certain perennial weeds, and weeds originating from underground parts (rhizomes, stolons, tubers) that are initially untreated, but do not apply more than 5K pints of TOUCHDOWN per acre per year as a combination of broadcast, spot, and wiper applications. TOUCHDOWN does not provide residual weed control, and as such, weeds emerging after application will not be controlled. For residual weed control, tank mix TOUCHDOWN with residual herbicides as prescribed in the TANK MIXTURES FOR BEARING CITRUS AND NONBEARING CROPS section, or make multiple applications.

If annual weeds have been mowed or grazed, allow 3-4 inches of new growth to appear prior to application. If perennial weeds have been mowed or grazed, allow new growth to reach recommended stage prior to application.

TANK MIXTURES WITH RESIDUAL HERBICIDES FOR BEARING CITRUS AND NONBEARING CROPS

TOUCHDOWN can be tank mixed with the following herbicides for control or suppression of emerged annual and perennial weeds. The following residual herbicides provide preemergence control of weeds listed on the individual product labels.

TOUCHDOWN plus:

- * DEVRINOL® 50-DF herbicide
- * Direx® 4L herbicide
- * Goal® 1.6E herbicide¹
- * Karmex® DF herbicide
- * Krovar® I herbicide
- * Krovar II
- * Prowl® herbicide
- * Prowl® 3.3 EC herbicide
- * Simazine/Caliber® 90 herbicide²
- * Simazine (80W, 4L, or Caliber 90)² plus Surflan® herbicide (AS or 75W)
- * Simazine 4L²
- * Simazine 80W²
- * Sim-Trol® 4L
- * Solicam® DF
- * Surflan AS
- * Surflan 75W
- * Goal 1.6E¹ plus Surflan (AS or 75W)
- * Goal 1.6E¹ plus Simazine (80W, 4L, or Caliber 90)²
- * Goal 1.6E¹ plus Surflan (AS or 75W) plus Simazine (80W, 4L, or Caliber 90)²

¹A tankmix of TOUCHDOWN and Goal cannot be used in bearing citrus.

²A tankmix of TOUCHDOWN and Simazine can only be used in bearing oranges, bearing grapefruit and in nonbearing crops.

Refer to the individual product labels for precautionary statements, restrictions, recommended rates and a list of weeds controlled.

Refer to the Tankmixes with Residual Herbicides section of the label for TOUCHDOWN rates and timing information.

11/22

PERENNIAL GRASS SUPPRESSION IN ORCHARD FLOORS

BAHIA GRASS

TOUCHDOWN can be used to inhibit seedhead emergence and suppress vegetative growth for approximately 40 to 50 days with a single application. By using a sequential application, suppression of vegetative growth and inhibition of seedhead emergence can be extended to 120 days. Applications must be made prior to seedhead emergence. Apply TOUCHDOWN at 20 to 30 days after complete green-up or after mowing to 4 inches tall. When a single application is planned, use 4 fluid ounces of TOUCHDOWN per acre plus 0.5 to 1.0% v/v nonionic surfactant/wetting agent in 10 to 25 gallons of water. When a sequential application is planned, use 3 fluid ounces per acre for the first application followed by another application of 3 fluid ounces per acre 40 to 50 days later.

BERMUDAGRASS

For Suppression Only:

East of the Rocky Mountains:
Apply 4 to 10 fluid ounces of TOUCHDOWN plus a nonionic surfactant/wetting agent in 3 to 20 gallons of water per acre. Make the application 2 weeks after complete green-up or after 3 to 4 inches of regrowth following mowing. Use 4 to 6% fluid ounces per acre if a lesser degree of suppression is desired. A sequential application can be used when regrowth occurs.

West of the Rocky Mountains:

Apply 10 fluid ounces of TOUCHDOWN plus a nonionic surfactant/wetting agent per acre in 3 to 10 gallons of water per acre. Make the application 2 weeks after complete green-up or after 3 to 4 inches of regrowth following mowing. A sequential application can be used when regrowth occurs.

For Partial Control and Burndown:

TOUCHDOWN can be used for burndown and partial control of bermudagrass at 1½ to 2½ pints plus a nonionic surfactant/wetting agent in 3 to 20 gallons of water per acre. Use 1½ pints east of the Rocky Mountains and 2½ pints west of the Rocky Mountains.

Use this treatment only if reduction of the bermudagrass stand can be tolerated. Allow at least 21 days for complete burndown.

COOL SEASON GRASS COVERS

For suppression of orchardgrass, fine fescue, tall fescue and quackgrass, apply 5 fluid ounces of TOUCHDOWN plus a nonionic surfactant/wetting agent in 10 to 20 gallons of water per acre. Add ammonium sulfate at 2% by weight or 17 pounds per 100 gallons of spray mix.

For suppression of Kentucky bluegrass, use 4 fluid ounces of TOUCHDOWN plus a nonionic surfactant/wetting agent. Do not add ammonium sulfate for suppression of Kentucky bluegrass.

LOW VOLUME APPLICATION (FLORIDA AND TEXAS ONLY)

To control or suppress the perennial weeds listed in the following table, apply the recommended rate of TOUCHDOWN plus a nonionic surfactant/wetting agent in 3 to 20 gallons of water per acre. Use 10-20 gallons per acre if weed foliage is dense. Apply when weeds are actively growing. Refer to the WEEDS CONTROLLED section for application timing for specific perennial weeds. If perennial weeds have been mowed or grazed, allow new growth to reach recommended growth stage prior to application.

USES FOR TOUCHDOWN IN LOW VOLUME APPLICATIONS IN FLORIDA AND TEXAS ONLY

Weed Species	Rate of TOUCHDOWN (Pints per Acre)		
	1½	2½	4
Bermudagrass	B	B	PC
Guineagrass Texas and Florida Ridge Florida Flatwoods	B	C	C
Paragrass	NR	C	C
Torpedograss	NR	S	PC

S = Suppression
PC = Partial Control
NR = Not Recommended

B = Burndown
C = Control

12/92

13/02

TABLE 1 (cont'd)
ANNUAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	MAXIMUM HEIGHT/ LENGTH (INCHES)	PINTS PER ACRE
Bluegrass, annual	<i>Poa annua</i>	6	%
Bluegrass, bulbous	<i>Poa bulbosa</i>		
Brome, downy	<i>Bromus tectorum</i>		
Cheat	<i>Bromus secalinus</i>		
Chickweed, common	<i>Stellaria media</i>		
Chickweed, mouseear	<i>Cerastium vulgatum</i>		
Cocklebur, common	<i>Xanthium strumarium</i>		
Corn	<i>Zea mays</i>		
Goatgrass, jointed	<i>Aegilops cylindrica</i>		
Groundsel, common	<i>Senecio vulgaris</i>		
Henbit	<i>Lamium amplexicaule</i>		
Horseweed/Marestail	<i>Conyza canadensis</i>		
Kochia	<i>Kochia scoparia</i>		
Lambsquarters, common	<i>Chenopodium album</i>		

29

TABLE 1
ANNUAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	MAXIMUM HEIGHT/ LENGTH (INCHES)	PINTS PER ACRE
Foxtails	<i>Setaria</i> sp.	12	%
Mustard, blue	<i>Chorispora tenella</i>	6	%
Mustard, tansy	<i>Descurainia pinnata</i>		
Mustard, tumble	<i>Sisymbrium altissimum</i>		
Mustard, wild	<i>Brassica kaber</i>		
Barley	<i>Hordeum vulgare</i>	12	%
Rye	<i>Secale cereale</i>		
Sandbur, field	<i>Cenchrus incertus</i>		
Shattercane	<i>Sorghum bicolor</i>		
Stinkgrass	<i>Eragrostis ciliaris</i>		
Morningglory	<i>Ipomoea</i> sp.	2	%
Sicklepod	<i>Cassia obtusifolia</i>		

28

27

GENERAL USE PRECAUTIONS FOR BEARING CITRUS AND NONBEARING CROPS

- Do not allow the spray, spray drift, or mist to contact green foliage or green bark on the trunk, suckers, open wounds, or other green parts of trees and vines. Contact with any tree or vine part other than mature brown woody bark may cause crop injury.
- Allow at least 13 days from the last application to harvest of citrus fruit.
- Avoid contact with stumps as injury to adjacent trees may occur from root grafting.

WEEDS CONTROLLED

All weeds listed are controlled by spray and hand-directed spot applications of TOUCHDOWN. For spray application, use the indicated rates. See DIRECTIONS FOR USE for mixing and application instructions.

14/00

TABLE 1 (cont'd)
ANNUAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	MAXIMUM HEIGHT/ LENGTH (INCHES)	PINTS PER ACRE
Sowthistle, annual Sprangletop Sunflower, common Thistle, Russian	<i>Sonchus oleraceus</i> <i>Leptochloa</i> sp. <i>Helianthus annuus</i> <i>Salsola iberica</i>	6	1½
Goosegrass Sicklepod Spanishneedles Spurge, annual	<i>Eleusine indica</i> <i>Cassia obtusifolia</i> <i>Biddens bipinnata</i> <i>Euphorbia</i> sp.	6-12	1½
Cutleaf eveningprimrose Carolina geranium	<i>Oenothera laciniata</i> <i>Geranium dessectum</i>	12	1½
Wheat	<i>Triticum aestivum</i>	12	1½
Velvetleaf	<i>Abutilon theophrasti</i>	6	2

32

TABLE 1 (cont'd)
ANNUAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	MAXIMUM HEIGHT/ LENGTH (INCHES)	PINTS PER ACRE
Sheperdspurse Sicklepod Signalgrass, broadleaf Wheat	<i>Capsella bursa-pastoris</i> <i>Cassia obusifolia</i> <i>Brachiaria platyphylla</i> <i>Triticum aestivum</i>	6	1
Amaranth, livid Amaranth, slender Amaranth, spiny Horseweed/Marestail Lambsquarters, common	<i>Amaranthus lividus</i> <i>Amaranthus viridus</i> <i>Amaranthus spinosus</i> <i>Conyza canadensis</i> <i>Chenopodium album</i>	12	1
Red rice Teaweed	<i>Oryza sativa</i> <i>Sida spinosus</i>	4	1½
Fleabane, rough Panicum, fall Pusley, Florida	<i>Erigeron strigosus</i> <i>Panicum dichotomiflorum</i> <i>Richardia scabra</i>	6	1½

31

TABLE 1 (cont'd)
ANNUAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	MAXIMUM HEIGHT/ LENGTH (INCHES)	PINTS PER ACRE
Pennycress, field Rocket, London Spurge, annual Woolly cupgrass	<i>Thlaspi arvense</i> <i>Sisymbrium irio</i> <i>Euphorbia</i> sp. <i>Eriochloa villosa</i>	6	¾
Crabgrass Johnsongrass, seedling Oats, wild Panicum, Texas Pigweed, redroot Pigweed, smooth Witchgrass	<i>Digitaria</i> sp. <i>Sorghum halepense</i> <i>Avena fatua</i> <i>Panicum texanum</i> <i>Amaranthus retroflexus</i> <i>Amaranthus hybridus</i> <i>Panicum capillare</i>	12	¾
Barnyardgrass Black nightshade Ryegrass, Italian	<i>Echinochloa crus-galli</i> <i>Solanum</i> sp. <i>Lolium multiflorum</i>	6	1

30

TABLE 3
PERENNIAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	PINTS PER ACRE	APPLICATION TIMING AND REMARKS
Alfalfa	<i>Medicago sativa</i>	2	At 6 to 8" or more after final cutting in fall. Deep till after 7 days
Artichoke, Jerusalem*	<i>Helianthus tuberosus</i>	5%	Actively growing, at or after flowering
Bahiagrass	<i>Paspalum notatum</i>	4	Actively growing, seed head stage
Bermudagrass	<i>Cynodon dactylon</i>	5%	Actively growing, seed heads present
Bindweed, field	<i>Convolvulus arvensis</i>	5%	Actively growing, beyond full bloom
Bluegrass, Kentucky	<i>Poa pratensis</i>	2%	Actively growing, boot to early seed head

35

*Partial Control

TABLE 2
ANNUAL WEED CONTROL - TOUCHDOWN RATES IN A TANKMIX WITH
0.25 LBS AI/A OF BANVEL OR 0.5 LBS AI/A OF 2,4-D

WEED SPECIES	SCIENTIFIC NAMES	MAXIMUM HEIGHT/LENGTH (INCHES)	PINTS PER ACRE
Lettuce, prickly Morningglory Ragweed, common Ragweed, giant Smartweed, Pennsylvania Velvetleaf	<i>Lactuca scariola</i> <i>Ipomoea</i> sp. <i>Ambrosia artemisiifolia</i> <i>Ambrosia trifida</i> <i>Polygonum pennsylvanicum</i> <i>Abutilon theophrasti</i>	6	1/2
Fleabane, rough Horseweed/Marestail Lambsquarters, common	<i>Erigeron strigosus</i> <i>Conyza canadensis</i> <i>Chenopodium album</i>	6-12	
Cocklebur, common Kochia Thistle, Russian	<i>Xanthium strumarium</i> <i>Kochia scoparia</i> <i>Salsola iberica</i>	12	

Read and follow Banvel and 2,4-D labels

TABLE 1 (cont'd)
ANNUAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	MAXIMUM HEIGHT/LENGTH (INCHES)	PINTS PER ACRE
Fleabane, rough Kochia Sowthistle, annual Sunflower, common Thistle, Russian	<i>Erigeron strigosus</i> <i>Kochia scoparia</i> <i>Sonchus oleraceus</i> <i>Helianthus annuus</i> <i>Salsola iberica</i>	6-12	2
Filaree Panicum, fall Sprangletop	<i>Erodium</i> sp. <i>Panicum dichotomiflorum</i> <i>Leptochloa</i> sp.	12	2

33

TABLE 3 (cont'd)
PERENNIAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	PINTS PER ACRE	APPLICATION TIMING AND REMARKS
Guineagrass	<i>Panicum maximum</i>	4	Actively growing, 7 leaf
Horsenettle	<i>Solanum carolinense</i>	5%	Actively growing, early head to early bud
Johnsongrass	<i>Sorghum halepense</i>	2	Actively growing, boot to head or prior to frost
Milkweed, common	<i>Asclepias syriaca</i>	4	Actively growing, late bud to flower
Muhly, wirestem	<i>Muhlenbergia frondosa</i>	2%	Actively growing, 8" in height or more, actively growing
Mullein, common	<i>Verbascum thapsus</i>	5%	Early head to early bud

38

TABLE 3 (cont'd)
PERENNIAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	PINTS PER ACRE	APPLICATION TIMING AND REMARKS
Cogongrass*	<i>Imperata cylindrica</i>	5%	Actively growing, late summer/ fall greater than 18"
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2	Actively growing, less than 6"
Dandelion	<i>Taraxacum officinale</i>	3	Actively growing, early bud
Dock, curly	<i>Rumex crispus</i>	3	Actively growing, early bud
Dogbane, hemp	<i>Apocynum cannabinum</i>	5%	Actively growing, late bud to flower
Fescue	<i>Festuca</i> sp.	4	Actively growing, boot to early seedhead
Goatweed	<i>Scoparia dulcis</i>	4	Actively growing, less than 7" tall

37

*Partial Control

TABLE 3 (cont'd)
PERENNIAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	PINTS PER ACRE	APPLICATION TIMING AND REMARKS
Brackenfern	<i>Pteridium aquilinum</i>	5%	Fronds fully expanded and at least 18" long
Bromegrass, smooth	<i>Bromus inermis</i>	2% - 4	Actively growing, boot to early seed head
Canarygrass, reed	<i>Phalaris arundinacea</i>	4	Actively growing, boot to head
Wheatgrass, Western	<i>Agropyron smithii</i>	4	Actively growing, boot to head
Cattail	<i>Typha</i> sp.	5%	Actively growing, early head to early bud
Clover, red	<i>Trifolium pratense</i>	5%	Actively growing, early head to early bud
Clover, white	<i>Trifolium repens</i>	5%	Actively growing, early head to early bud

36

17/02

TABLE 3 (cont'd)
PERENNIAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	PINTS PER ACRE	APPLICATION TIMING AND REMARKS
Thistle, Canada	<i>Cirsium arvense</i>	4	Actively growing beyond bud stage
Timothy	<i>Phleum pratense</i>	4	Actively growing boot to head
Torpedograss*	<i>Panicum repens</i>	5%	Actively growing at or beyond seedhead
Trumpet creeper*	<i>Campsis radicans</i>	2%	Actively growing late September/October
Vaseygrass	<i>Paspalum urvillei</i>	2	Actively growing, less than 12 inches in height

41

*Partial Control

TABLE 3 (cont'd)
PERENNIAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	PINTS PER ACRE	APPLICATION TIMING AND REMARKS
Quackgrass	<i>Agropyron repens</i>	1%	Actively growing, less than 5" tall - retreatment may be required
		3%	Actively growing, greater than 8" tall
Redvine*	<i>Brunnichia ovata</i>	2	Actively growing, greater than 18" tall in September, October
Ryegrass, perennial	<i>Lolium perenne</i>	2	Actively growing, boot to head or prior to frost
Smartweed, swamp	<i>Polygonum coccleum</i>	2	Actively growing, early bud
Spurge, leafy	<i>Euphorbia esula</i>	3	Actively growing, greater than 12" tall in late summer or fall

40

*Partial Control

TABLE 3 (cont'd)
PERENNIAL WEED CONTROL - TOUCHDOWN RATES

WEED SPECIES	SCIENTIFIC NAMES	PINTS PER ACRE	APPLICATION TIMING AND REMARKS
Nutsedge, purple Nutsedge, yellow	<i>Cyperus rotundus</i> <i>Cyperus esculentus</i>	4	Actively growing, flower
		2+2	Make sequential applications when actively growing, 3-5 leaf, less than 6" tall
Orchardgrass	<i>Dactylis glomerata</i>	2% - 4	Actively growing, minimum height of 12" in spring, 6" in fall
Paragrass	<i>Brachiaria mutica</i>	2%	Actively growing, seedhead stage
Phaseybean*	<i>Phaseolus lathyroides</i>	5%	Actively growing, less than 8" tall

39

*Partial Control

04004234/R4 Touchdown bk1c 12/6/96 11:07 PM Page 40

18/22

NONAGRICULTURAL USE REQUIREMENTS
 The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.
 Keep people and pets off treated areas until spray solution has dried.

- FARMSTEAD (NONCROP)**
 Applications can be made in noncrop areas around the farm such as:
- Farmyards
 - Soil Bank Land
 - Farm Buildings
 - Fuel Storage Areas
 - Apply to annual and perennial grass and broadleaf weeds according to the WEEDS CONTROLLED section of this label.
 - Rights-of-way
 - Barrier Strips
 - Equipment Areas
 - Fence Rows
- Avoid contact with the foliage of ornamentals or other desirable plants.
 Repeat applications may be necessary.

Made in U.S.A.
ZENECA Ag Products
 ZENECA, Inc.
 Wilmington, Delaware 19850-5458

04004234
 029604 Rev. 11/96

TABLE 4
PERENNIAL WEED CONTROL - TOUCHDOWN RATES IN A TANKMIX
WITH 0.25 LBS AI/A OF BANVEL OR 0.5 LBS AI/A OF 2,4-D

WEED SPECIES	SCIENTIFIC NAMES	PINTS PER ACRE	APPLICATION TIMING AND REMARKS
Dandelion	<i>Taraxacum officinale</i>	½	Actively growing, early bud
Dock, curly	<i>Rumex crispus</i>	½	Actively growing, early bud
Dogbane, hemp	<i>Apocynum cannabinum</i>	1½	Actively growing, late bud to flower
Milkweed, common	<i>Asclepias syriaca</i>	1½	Actively growing, late bud to flower
Milkweed, honeyvine	<i>Ampelamus albidus</i>	2%	Actively growing, late bud to flower
Smartweed, swamp	<i>Polygonum coccineum</i>	½	Actively growing, early bud
Spurge, leafy*	<i>Euphorbia esula</i>	½	Actively growing, greater than 12" tall in late summer or fall

*Partial control
 Read and follow Banvel and 2,4-D labels

19/22

Touchdown[®]

HERBICIDE

NONSELECTIVE FOLIAR SYSTEMIC HERBICIDE FOR WEED CONTROL

ACTIVE INGREDIENT:	Sulfonium, trimethyl-salt with N-(phosphonomethyl) glycine (1:1)*	57.6%
INERT INGREDIENTS:		42.4%
Total		100.0%

*Sulfosac
 Contains 6 pounds active ingredient per gallon.
 EPA Reg. No. 10182-324

EPA Est. No. _____

NOTIFICATION

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If a known exposure occurs or is suspected, immediately initiate the recommended procedures below. If further treatment is required, contact a POISON CONTROL CENTER, a physician, or the nearest hospital.

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth if victim is unconscious or convulsing.

IF ON SKIN: Flush with plenty of water for at least 15 minutes. Get medical attention if skin irritation occurs.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F.A.-S-T-M-E-D (327-8633).
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through the skin, or inhaled. Prolonged contact may cause skin irritation. May cause allergic skin reactions. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds and streams. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

In case of spill, ISOLATE the spill. Absorb spill with inert absorbent material such as clay or Fuller's earth. Sweep up used absorbent and place in an appropriate chemical waste container. Flush spill area with water. Observe all local, State and Federal laws and regulations regarding disposal, spill, cleanup, removal or discharge.

OTHER PRECAUTIONS

Caution must be taken when applying TOUCHDOWN nonselective foliar systemic herbicide to avoid drift or contact with nontarget plant species. Such contact may result in plant injury.

Stainless steel, plastic-lined steel, plastic or fiberglass containers are acceptable for mixing and storing TOUCHDOWN. TOUCHDOWN should not be mixed or stored in galvanized steel or unlined steel containers. TOUCHDOWN is not compatible with carbon steel and neoprene or butyl rubber.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability found in the attached booklet before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Apply this product only as specified on the label.
- Do not apply this product through any type of irrigation system.
- Do not exceed a total of 5 1/3 pints of TOUCHDOWN per acre per year.

REFER TO ATTACHED BOOKLET FOR COMPLETE DIRECTIONS FOR USE.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling (i.e., attached booklet) under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Rinse spray equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America. U.S. Patent No. 4,315,765.

TOUCHDOWN[®] is a trademark of a ZENECA Group Company.
©1996 ZENECA Inc.

Contents: _____

Made in U.S.A.

ZENECA Ag Products

ZENECA Inc.
Wilmington, Delaware 19850-5458

9404231
079604
Rev. 11/96

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

If a known exposure occurs or is suspected, immediately initiate the recommended procedures below. If further treatment is required, contact a POISON CONTROL CENTER, a physician, or the nearest hospital.

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth if victim is unconscious or convulsing.

IF ON SKIN: Flush with plenty of water for at least 15 minutes. Get medical attention if skin irritation occurs.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL
1-800-F-A-S-T-M-E-D (327-8633).
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call
CHEMTREC 1-800-424-9300.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through the skin, or inhaled. Prolonged contact may cause skin irritation. May cause allergic skin reactions. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds and streams. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Touchdown®
HERBICIDE

**NONSELECTIVE FOLIAR SYSTEMIC
HERBICIDE FOR WEED CONTROL**

ACTIVE INGREDIENT:

Sulfonilum, trimethyl-salt with N-(phosphonomethyl) glycine (1:1)*57.6%

INERT INGREDIENTS:42.4%

Total100.0%

*Sulfosate

Contains 6 pounds active ingredient per gallon.

EPA Reg. No. 10182-324

EPA Est. No. 10182-AR-1 10182-NB-1

Superscript corresponds to first letter of lot number.

CONTENTS: 1 QUART

Made in U.S.A.

ZENECA Ag Products

ZENECA Inc.
Wilmington, Delaware 19850-5458

In case of spill, ISOLATE the spill. Absorb spill with inert absorbent material such as clay or Fuller's earth. Sweep up used absorbent and place in an appropriate chemical waste container. Flush spill area with water. Observe all local, State and Federal laws and regulations regarding disposal, spill, cleanup, removal or discharge.

OTHER PRECAUTIONS

Caution must be taken when applying TOUCHDOWN nonselective foliar systemic herbicide to avoid drift or contact with nontarget plant species. Such contact may result in plant injury.

Stainless steel, plastic-lined steel, plastic or fiberglass containers are acceptable for mixing and storing TOUCHDOWN. TOUCHDOWN should not be mixed or stored in galvanized steel or unlined steel containers. TOUCHDOWN is not compatible with carbon steel and neoprene or butyl rubber.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability found in the attached booklet before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Apply this product only as specified on the label.
- Do not apply this product through any type of irrigation system.
- Do not exceed a total of 5 1/3 pints of TOUCHDOWN per acre per year.

REFER TO ATTACHED BOOKLET FOR COMPLETE DIRECTIONS FOR USE.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling (i.e., attached booklet) under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Rinse spray equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Plastic-Triple rinse (or equivalent). Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America, U.S. Patent No. 4,315,765.

04004253 TOUCHDOWN® is a trademark of a ZENECA Group Company.
029604 ©1996 ZENECA Inc.

copy

2/1/22

ZENECA Ag Products NOTIFICATION

Touchdown[®]

HERBICIDE

NONSELECTIVE FOLIAR SYSTEMIC HERBICIDE FOR WEED CONTROL

ACTIVE INGREDIENT:

Sulfonium, trimethyl-salt with N-(phosphonomethyl) glycine (1:1)* 57.6%

INERT INGREDIENTS: 42.4%

Total 100.0%

*Sulfosate

Contains 6 pounds active ingredient per gallon.

EPA Reg. No. 10182-324

EPA Est. No. 10182-AR-1 10182-NB-1

Superscript corresponds to first letter of lot number.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If a known exposure occurs or is suspected, immediately initiate the recommended procedures below. If further treatment is required, contact a POISON CONTROL CENTER, a physician, or the nearest hospital.

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth if victim is unconscious or convulsing.


IF ON SKIN: Flush with plenty of water for at least 15 minutes. Get medical attention if skin irritation occurs.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

Net Contents: 2.5 gal

94003807
029504
Rev. 11/96



22/22

Touchdown

HERBICIDE

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through the skin, or inhaled. Prolonged contact may cause skin irritation. May cause allergic skin reactions. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds and streams. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

In case of spill, ISOLATE the spill. Absorb spill with inert absorbent material such as clay or Fuller's earth. Sweep up used absorbent and place in an appropriate chemical waste container. Flush spill area with water. Observe all local, State and Federal laws and regulations regarding disposal, spill, cleanup, removal or discharge.

OTHER PRECAUTIONS

Caution must be taken when applying TOUCHDOWN® nonselective foliar systemic herbicide to avoid drift or contact with nontarget plant species. Such contact may result in plant injury.

Stainless steel, plastic-lined steel, plastic or fiberglass containers are acceptable for mixing and storing TOUCHDOWN. TOUCHDOWN should not be mixed or stored in galvanized steel or unlined steel containers. TOUCHDOWN is not compatible with carbon steel and neoprene or butyl rubber.

Net Contents: 2.5 gal

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability found in the attached booklet before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Apply this product only as specified on the label.
- Do not apply this product through any type of irrigation system.
- Do not exceed a total of 5 1/2 pints of TOUCHDOWN per acre per year.

REFER TO LABEL BOOKLET ATTACHED TO CONTAINER FOR COMPLETE DIRECTIONS FOR USE.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling (i.e., attached booklet) under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Rinse spray equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Plastic—Triple rinse (or equivalent). Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America. U.S. Patent No. 4,315,765. TOUCHDOWN® is a trademark of a ZENECA Group Company. ©1996 ZENECA Inc.

Made in U.S.A.

ZENECA Ag Products

ZENECA Inc.
Wilmington, Delaware 19850-5458

04003808
029604
Rev. 11/96