

10/18/97 10:00:00 PM 04 10182-370 7/11/97 Page 1 of 2

SUPPLEMENTAL LABEL FOR USE ON ALFALFA GROWN FOR SEED IN THE STATE OF WYOMING ONLY

PIRIMOR[®] 50-DF INSECTICIDE

EPA Reg. No. 10182-370

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.


ALFALFA GROWN FOR SEED: For control of aphids. Apply 2 to 6 oz. of PIRIMOR 50-DF per acre by ground or aerial application. The higher rate is recommended for instances of heavy infestations of aphids. Spray to wet foliage completely. Minimal application spray volume is 10 gal. per acre by ground and 5 gal. per acre by air.

For the most effective and economic control, begin to apply PIRIMOR 50-DF as soon as aphids appear.

PRE-HARVEST INTERVAL (PHI): A PHI of 1 day is required after application of PIRIMOR.

GENERAL USE PRECAUTIONS

- PIRIMOR 50-DF is to be applied at a rate not to exceed 6 ounces of product per acre per season.
- PIRIMOR 50-DF may be applied a maximum of twice per season not to exceed 6 ounces of product per acre per season (i.e., two applications at 3 oz. each). For a second application, reapply in no less than 7 to 10 days.
- Do not apply when weather conditions favor drift from treated areas.
- Make ground applications when the wind velocity favors on target product deposition, (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph.
- Do not apply by ground equipment within 25 feet, or by air within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Applicators and alfalfa seed growers are required to report to EPA Headquarters, the EPA Regional Office and the State Department of Agriculture any adverse effects to birds resulting from the use of this pesticide.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.
- Ultra Low Volume (ULV) applications are prohibited.



- The largest droplet size consistent with good pest control is required. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Spray should be released at the lowest height consistent with pest control and flight safety.
- Aerial applications should be made when the wind velocity favors on-target product deposition. Do not apply when wind velocity exceeds 10 mph. Avoid applications when wind gusts approach 10 mph.
- To avoid risk of exposure to sensitive aquatic areas, aerial applicators should avoid application when wind direction is toward aquatic areas.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.
- Do not make aerial applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.
- For use only on fields in production of alfalfa seed. Not for use on fields producing alfalfa for livestock feed. No portion of the treated field, including seed, seed screenings, hay forage or stubble, may be used for human or animal feed.
- Do not cut current years' treated alfalfa seed crop for hay or forage. Do not graze current years' treated alfalfa seed crops.
- Screening from alfalfa seed processing are prohibited from feed channels. All PIRIMOR 50-DF treated alfalfa seed screening must be removed from the feed market.
- Treated alfalfa seed is not to be used for sprouting. All alfalfa seed treated with PIRIMOR 50-DF is to be tagged at processing plants, NOT FOR HUMAN CONSUMPTION. It shall be the grower's responsibility to notify the processing plants of any seed crop treated with PIRIMOR 50-DF.
- Do not apply this product through any type of irrigation system.

COMPATIBILITY: Do not use with oxidizing agents.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

PIRIMOR® is a trademark of a ZENECA Group Company.

09-2584-011 07/87