

PM 22

10182-382

10/22



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

10182-382

Date of Issuance:

APR 7 1994

Term of Issuance: Conditional

Name of Pesticide Product:

Bonsi II Ornamental Growth Regulator

NOTICE OF PESTICIDE:
 X Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Zeneca, Inc.
Zeneca Ag Products
1800 Concord Pike
Wilmington, DE 19897

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his own motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label changes:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 10182-382".
 - b. You must group together the precautionary statements on the front panel of the label or place a referral statement on the front panel of the label.
 - c. Under the "AGRICULTURAL USE REQUIREMENTS" section, revise the restricted entry interval (REI) to 24 hours, not 12 hours. In addition, delete "Waterproof gloves" from the PPE required for early entry and replace it with "Chemical-resistant gloves".
3. Submit two copies of the revised final printed label before you release your product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Date:

APR 7 1994

20/00

BONZI® II
Ornamental Growth Regulator

ACTIVE INGREDIENT:

Paclobutrazol

(±)-(R*,R*)-beta-((4-Chlorophenyl)methyl)-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol 00.13%

INERT INGREDIENTS: 99.87%

Total 100.0%

BONZI 2 contains 0.036 grams active ingredient per fluid ounce (1280 ppm).

EPA Reg. No.: 10182-
EPA Est. No.:

Contents:

KEEP OUT OF REACH OF CHILDREN

CAUTION

Made in U.S.A.
ZENECA Professional Products
ZENECA Inc.
Wilmington, DE 19897

ACCEPTED
with COMMENTS
In EPA Letter Dated:

APR 7 1994

Federal Insecticide
and Fungicide Act
for the pesticide
EPA Reg. No.

10182-382

30/02

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL 1-800-F-A-S-T-M-E-D (327-8633)

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

HARMFUL IF ABSORBED THROUGH SKIN. AVOID CONTACT WITH SKIN OR CLOTHING.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber or Nitrile Rubber or Neoprene Rubber or Polyvinyl Chloride or Viton.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
-

BEST AVAILABLE COPY


48/22

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flames.



5022

**CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

BEST AVAILABLE COPY

60/100

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use. Refer to label booklet attached to container for complete directions for use.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling (i.e., attached booklet) under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

STORAGE AND DISPOSAL


PROHIBITIONS: Do not reuse empty containers. Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep container closed when not in use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Plastic containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BONZI® is a trademark of a ZENECA Group Company.



70/22

BONZI® II
Ornamental Growth Regulator

COMPLETE DIRECTIONS FOR USE

**CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

BEST AVAILABLE COPY

80/25

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

Made in U.S.A.
ZENECA Professional Products
ZENECA Inc.
Wilmington, DE 19897

9 5

9/2/22

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACTIVE INGREDIENT:

Paclobutrazol

(±)-(R*,R*)-beta-((4-Chlorophenyl)methyl)-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol

00.13%

INERT INGREDIENTS:

99.87%

Total

100.0%

BONZI 2 contains 0.036 grams active ingredient per fluid ounce (1280 ppm).

EPA Reg. No.: 10182-

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL 1-800-F-A-S-T-M-E-D (327-8633)

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

HARMFUL IF ABSORBED THROUGH SKIN. AVOID CONTACT WITH SKIN OR CLOTHING.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

BEST AVAILABLE COPY

10/1/22

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber or Nitrile Rubber or Neoprene Rubber or Polyvinyl Chloride or Viton.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
-

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flames.

1/10/02

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

24 hours
Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not reuse empty containers. Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep container closed when not in use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

BEST AVAILABLE COPY

Daf

CONTAINER DISPOSAL: Plastic containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

BONZI® II is a plant growth regulator for use on ornamental plants grown in containers in greenhouses, shadehouses, and interiorscapes. Use of BONZI II effectively reduces internode elongation, resulting in more desirable compact plants. When used as directed, BONZI II produces no phytotoxic effects.

MIXING INSTRUCTIONS

Be sure the sprayer is clean and not contaminated with any material. Fill the spray tank with half the required amount of water. Using the dilution table (Table 1), determine the amount of BONZI II needed for the required concentration. Measure the desired volume accurately and add it to the tank.

Fill tank with the remaining amount of required water.

Agitate the mixture of BONZI II and water frequently to assure uniform distribution during application.

134/02

**TABLE 1
BONZI II DILUTION TABLE**

ppm BONZI II Desired	Fl. Oz. Per Gallon	mL Per Gallon
1	0.1	3
2	0.2	6
3	0.3	9
4	0.4	12
5	0.5	15
10	1.0	30
20	2.0	60
25	2.5	75
30	3.0	90
40	4.0	120
50	5.0	150
100	10.0	300
200	20.0	600

APPLICATION TECHNIQUES

Desired height control with BONZI II can be obtained with three different types of applications: sprays, drenches or bulb soaks.

Under certain conditions, sequential spray applications may be desirable.

Frequent agitation of the BONZI II solution and proper application techniques are critical in order to achieve desired results. Be sure of your calculations, volume measurements and sprayer calibration. When in doubt, recalculate.

1. SPRAY APPLICATIONS

In spray applications, BONZI II penetrates into plant stems and is translocated to the terminal where it reduces internode elongation.

When applying a spray application, it is important that:

- adequate spray volume is used to thoroughly wet plant stems. The misting technique used for some other growth retardants, where only upper leaves are covered with a light spray, will not produce desired results with BONZI II;

BEST AVAILABLE COPY

- sprays are not applied to the point of excessive runoff into the potting media. The spray volume which drips down into the media may be desirable as it will be taken up by the roots and increase the effectiveness of BONZI II. However, too much runoff into the media may result in excessive height control;
- the spray technique provide thorough, consistent, uniform coverage of all plants. Failure to do so may result in non-uniform height control..

BONZI II may be applied at any time of the day without danger of burning leaves or causing chlorosis.

Overhead irrigation or rain 30 minutes after spray applications does not reduce the effectiveness of BONZI II.

Addition of wetting agents for spray applications is not necessary.

There are two methods of applications for insuring uniform coverage of BONZI II sprays:

Bench Spray Method: The recommended spray volume for small plants in small containers which are closely spaced is 2 qts./100 sq. ft. of bench space. For larger plants with a well-developed canopy, a spray volume of 3 qts./100 sq. ft. of bench space is recommended.

Individual Plant Method: Apply sprays to individual plants, insuring that all plants of the same species/cultivar and size are treated with an equivalent amount of spray volume. Spray volume requirements will vary with plant size and type: For poinsettias in 6-inch pots with laterals from 1½ to 2 inches, the recommended spray volume is about 0.7 fl. oz. (20 ml.)

Sequential Applications: Using sequential applications may provide more uniform growth regulation and safety against over application. In general, sequential spray applications are to be applied using 50-100% of the lower recommended rate. Growers in cooler climates may have to use lower rates.

With some plant species, particularly chrysanthemums, hibiscus and azaleas, individual lateral shoots will outgrow the other laterals, causing non-uniform plant appearance. This results when individual laterals do not receive enough chemical when spray is applied. The use of sequential applications will reduce this problem.

2. DRENCH APPLICATIONS

Application of BONZI II to the growing media will provide good control of plant height. BONZI II is readily absorbed by plant roots and translocated to the terminals.

Drench applications generally provide a longer-lasting, more uniform height control than spray applications. Drench applications should be made to moist potting media. This may be achieved by watering plants the day before treatments. Drench applications to dry media will result in poor distribution.

BEST AVAILABLE COPY

15 of 22

Multiple plants growing in the same pot require a more uniform distribution of drench solution to achieve uniform height control.

Drench Rates and Volumes: The rates recommended for soil drench applications are based on a drench volume of 4 fl. oz. of final solution for an average 6-inch 'azalea' pot. Based on this recommendation, one gallon of solution will treat 32 6-inch pots. For smaller or larger pots, a suitable drench volume is enough final solution applied to achieve total run through of no more than 10%, providing that the potting media is properly moist before treatment. Table 2 may be used as a guide in determining appropriate drench volume needed for the specified pot sizes. For the grower who likes to apply BONZI II as a known amount of active ingredient per pot, Table 2 also shows the amount of active ingredient found in a specific volume at a known concentration.

**TABLE 2
DRENCH VOLUME GUIDELINES AND CONVERSIONS**

Pot Diameter (inches)	Drench Volume (fl. oz./pot)	Mg. ai. BONZI II/Pot			
		1 PPM	2 PPM	3 PPM	4 PPM
4"	2	0.063	0.125	0.188	0.25
5"	3	0.094	0.188	0.282	0.375
6"	4	0.125	0.25	0.375	0.50
8"	10	0.313	0.625	0.938	1.25
10"	25	0.783	1.56	2.35	3.125
12"	40	1.25	2.5	3.75	5.0

NOTE: The recommended drench volumes were based on the soil capacity of a common 6-inch 'azalea' type pot. Extrapolating the recommendation for this 6-inch 'azalea' type pot to smaller or larger containers may not be correct for total drench volume but should only be used as a guideline. The user must determine the appropriate rate and drench volume needed to achieve the desired result, based on both pot size and potting media used.

3. PREPLANT BULB SOAKS

Soaking of bulbs in solutions of BONZI II is also a very effective way to attain height control. The rates used and length of soaking time will vary, depending on the species. See the section on BULB CROPS for specific recommendations.

FACTORS AFFECTING PLANT RESPONSE TO BONZI II

IN ADDITION TO PROPER APPLICATION TECHNIQUE, THERE ARE SEVERAL ENVIRONMENTAL AND CULTURAL FACTORS WHICH CAN AFFECT A PLANT'S RESPONSE TO TREATMENT WITH BONZI II. These factors may cause a variation in the amount of BONZI II needed to provide desired plant height.

BEST AVAILABLE COPY

160/22

Cultural practices may affect the plant's response to BONZI II. Plants which are grown at close spacing or in smaller pots and using high water and fertilizer levels may require an increase in the amount of BONZI II needed.

For drench applications, plants grown in media with pine bark or a high organic content may require higher rates of BONZI II than those grown in media without pine bark or with a low organic content.

Different varieties or cultivars within a given plant species may require a higher or lower rate of BONZI II. The taller, more vigorous varieties generally require more chemical than do the naturally short, less vigorous varieties.

Growers should consult with plant and seed suppliers for vigor and other growth characteristics for newly released varieties.

Temperature can be the overriding factor in determining the amount of BONZI II needed. Stem elongation increases with increased temperatures. Growers in warm climates need to use higher rates and/or more applications compared to those in cooler climates.

The amount of BONZI II needed and number of applications may also vary depending on the time of year, with higher rates and/or more applications needed during warmer months.

DETERMINING OPTIMUM RATES

Optimum BONZI II rates will vary with different growers and will depend on their individual desired final plant height, growing conditions, and applications techniques. Different varieties or cultivars of the same species may respond differently to BONZI II. Growers should conduct trials with small numbers of plants using the recommended rates to determine the optimum rates for their situations.

Before BONZI II is applied to large numbers of plants, it is important that the user determine their own optimum rate for each species.

The rates recommended on this label are rate ranges and should be used only as a guideline.

The user should conduct trials on a small number of plants, adjusting the rate of BONZI II to achieve the desired height and length of control. For preplant bulk soak trials, it may be necessary to adjust both the rate and length of soak time in order to achieve desired results.

For plant species listed on the label, the user should run initial trials using the lowest recommended rates.

For plant species not specifically listed on the label, the user should run initial trials using the rates recommended in Table 3.

BEST AVAILABLE COPY

17 of 22

TABLE 3
RECOMMENDED TRIAL RATES (ppm) BY GENERAL PLANT TYPE*

Plant Type	Spray	Drench	Bulb Soak
Bedding Plants	30	1	N/A
Bedding Plant Plugs	5	NR	N/A
Flowering/Foliage Plants (Annual or perennial)			
- Herbaceous Species	30	1	N/A
- Woody Species	50	2	N/A
Woody Landscape Plants	100	4	N/A
Bulb Crops	100	10	20 (@ 15 min.)

NR = Use is not recommended
N/A = Use is not applicable

* The recommended trial rates are based on information developed in the Sunbelt Region. Growers in regions north of the Sunbelt should run initial tests using 1/2 X the recommended trial rates listed in Table 3.

USE AND RATE RECOMMENDATIONS BY CROP

BONZI II is effective in controlling the height of most ornamental crops. Be sure to read and fully understand the section on DETERMINING OPTIMUM RATES before applying to large numbers of plants.

A. AZALEAS (FLORIST)



BONZI II can be used to control plant height, reduce bypass shoot elongation and promote flower bud initiation.

Spray applications are effective in the rate range of 100 to 200 ppm.

Drench Applications are effective in the rate range of 5 to 15 ppm. To control plant height and promote flower bud initiation, applications should begin when new growth, following final shaping, is 1 1/2 to 2 inches long.

To reduce bypass shoot development, applications should be made after bud set when bypass shoots are barely visible, or about 5 to 7 weeks prior to cooling.

BEST AVAILABLE COPY

80722

B. BEDDING PLANTS

Spray applications of BONZI II will provide height control of most bedding plants at a wide rate range of 5 to 90 ppm. The rate ranges for some specific bedding plants are:

Plant	Rate Range (ppm)	Plant	Rate Range (ppm)
Ageratum	15 - 45	Marigold (French)	15 - 45
Celosia	15 - 45	Pansy	5 - 15
Coleus	15 - 45	Petunia	30 - 60
Dahlia	15 - 45	Salvia	20 - 60
Dianthus	20 - 60	Snapdragon	45 - 90
Impatiens	15 - 45	Zinnia	15 - 45
Marigold (African)	30 - 60		

- Do not use on fibrous begonias as they are very sensitive to BONZI II. Overly stunted plants can result if they receive spray drift from applications made to surrounding species.
- Do not use on annual Vinca (periwinkle) as BONZI II may cause spotting of foliage, especially at high temperatures.

For bedding plants not specifically listed above, the user should determine optimum rates starting with a rate of 30 ppm in the Sunbelt Region and 15 ppm in the Northern Belt Region. Time of application should begin when new growth in height or width reaches 2 inches or when plants reach desired size to hold them at a marketable stage.


Late application timings and/or excessive rates may slow the growth of plants when transplanted.

High rates of BONZI II may delay flowering, especially of impatiens.

Drench applications are effective on bedding plants, but are recommended only for those plants in containers 6 inches or larger. The user should determine optimum rates, starting at 1 ppm.

C. BEDDING PLANT PLUGS

Spray applications of BONZI II can also be used to effectively control height of bedding plant plugs. The recommended rate range of 1 to 20 ppm is much lower than the rate range for older bedding plants.



190/02

Plant	Rate Range (ppm)	Plant	Rate Range (ppm)
Ageratum	5 - 10	Marigold	10 - 20
Celosia	5 - 10	Pansy	1 - 5
Coleus	5 - 10	Petunia	5 - 10
Dahlia	5 - 10	Salvia	5 - 10
Dianthus	5 - 10	Snapdragon	5 - 10
Impatiens	5 - 20	Zinnia	5 - 10

For bedding plant plugs not specifically listed above, the user should determine optimum rates starting with a rate of 5 ppm. Timing of application should begin at the 1 to 2 true leaf stage.

Drench applications are not recommended for bedding plant plugs due to the sensitivity and extremely low rates needed.

D. BULB CROPS

Height control can be achieved on a variety of bulb crops by any of the three application types.

Spray applications, although moderately effective, are the least desirable method for controlling height. Sequential applications are recommended in order to achieve desired uniformity. Applications should begin when plants are 2 to 4 inches tall.

Drench applications are very effective in the rate range of 8 to 160 ppm. Optimum rates vary widely, depending on species. Timing of application will also vary, depending on species. For bulbs which require a cold period, BONZI II is generally applied 1 to 5 days after removal from the cooler. For most other bulb types, application should be made when newly emerged shoots are 1 to 2 inches tall.

Preplant bulb soaks are also very effective. Effective rates for most species are in the range of 5 to 25 ppm, with a soaking time of 15 minutes to 1 hour. In general, lower use rates will require longer soaking times.

The following table gives recommended rate ranges and length of soaking time for a variety of bulb species:

BEST AVAILABLE COPY

200100

Bulb Type	Spray Rate (ppm)	Drench Rate (ppm)	Preplant Bulb Soak Rate (ppm) / Soak Time
Amaryllis	ND	200	ND
Caladium	100 - 200	10 - 20	ND
Calla Lily	ND	10 - 30	20 / 15 min.
Daffodil	ND	20 - 40	ND
Freesia	ND	ND	100 - 300 / 1 hr.
Hybrid Lily	250 - 500	10 - 20	20 - 30 / 15 min.
Montbretia	ND	ND	20 - 30 / 15 min.
Tulip	ND	5 - 40	2 - 5 / 1 hr.

ND = Rates for this particular use have not been determined. For these applications the user should run trials using the rates recommended in Table 3.

For species not specifically listed, trials should be conducted using rates outlined in the section on DETERMINING OPTIMUM RATES.

E. CHRYSANTHEMUMS (POT)

BONZI II is effective in controlling the height of pot chrysanthemums when applied as either a spray or drench.

Spray applications are effective at rates of 50 to 200 ppm. Applications should begin when axillary shoots are 2 to 3 inches long. BONZI II can be applied earlier to vigorous varieties if additional control is desired.

Sequential applications of lower rates generally provide more uniformly shaped plants than single-spray applications.

BONZI II may be applied at time of disbud to reduce late stretch without reducing flower size or delaying flowering.

Drench applications of BONZI II are effective at recommended rates of 1 to 4 ppm. Application timing is when axillary shoots are 2 to 3 inches long.

Because pot chrysanthemums are usually planted with multiple cuttings per pot, uniform application is critical to achieving desired results.

F. FLOWERING PLANTS/FOLIAGE PLANTS (not specifically listed)

BONZI II is effective as a spray or drench application in controlling height on a wide variety of other flowering plants and foliage plants. It can be used as either a holding agent to stop growth (e.g., interiorscape) or a toning agent to slow growth. In general, herbaceous

BEST AVAILABLE COPY

2/10/22

species will require lower rates than woody species. Trials should be conducted using rates outlined in the section on DETERMINING OPTIMUM RATES.

G. GERANIUMS

Geraniums are particularly sensitive to BONZI II. The user must determine optimum rates before applying to large numbers of plants.

Spray applications of BONZI II at recommended rates of 10 to 30 ppm will effectively control growth of geraniums. Early applications may require lower rates to avoid overstretching. Time of application for zonal geraniums is when growth is 1½ to 2 inches long; for seed geraniums, 2 to 4 weeks after transplanting or when needed.

BONZI II will reduce late stretch when applied as the flower stems begin to elongate.

Drench applications, although effective, should be made with caution due to the extreme sensitivity of geraniums to BONZI II. Trials should be conducted to determine optimum rates.

H. HIBISCUS

Spray applications at 30 to 150 ppm will effectively reduce shoot elongation. Application should be made when laterals are 1 to 4 inches long, depending on desired final plant size.

Single applications will control lateral growth for 3 to 6 weeks. Sequential applications may provide more uniform plant shape. BONZI II can be applied 1 to 2 weeks prior to flowering to prevent late stretch.

Drench applications will also effectively reduce shoot elongation. Trials should be conducted using recommended rates outlined in the section on DETERMINING OPTIMUM RATES.

I. POINSETTIAS

Spray applications of BONZI II will effectively control height of poinsettias. Recommended rates at 10 to 30 ppm for most areas of the U.S. In southern Florida, higher rates of 15 to 45 ppm are recommended.

Single applications may be made using the higher recommended rates. However, sequential applications initially using lower rates will provide better safety against overstretching retarded plants. For subsequent applications use 50 to 100% of the initial rate, depending on plant vigor at the time of reapplication.

Applications to slower growing varieties in cool climates should begin when axillary shoots are 2 to 3 inches long. For vigorous growing varieties in warm climates, applications should begin when axillary shoots are 1½ to 2 inches long. Sequential applications may be applied 1 to 3 times, at 7 to 14-day intervals, depending on plant vigor/growth.

BEST AVAILABLE COPY

22 of 22

Seasonably late applications of BONZI II will reduce plant height, but, like most PGR's may also reduce bract size. For growers scheduling early December flowering, BONZI II should not be applied after initiation of short days. As a guide, do not apply BONZI II after October 1 for areas outside of Florida, or after October 25 in Florida.

Drench applications generally have less of an effect on bract size than do sprays. Recommended rates are in the range of 2 to 4 ppm, based on a drench volume of 4 fl. oz./6-inch pot. Application should be made when axillary shoots are 2 to 3 inches long.

NOTE: Optimum BONZI II rates and timings for both spray and drench applications to poinsettias will vary depending on the variety.

J. WOODY LANDSCAPE PLANTS (Container-grown in greenhouses/shadehouses)

BONZI II is effective in controlling height on a wide variety of woody landscape plants using both spray or drench applications. Rate ranges for different species vary greatly. Trials should be conducted using rates outlined in the section on DETERMINING OPTIMUM RATES.

Examples of woody plants on which the product can be applied are:

- | | |
|---------------|--------------|
| Azalea | Juniper |
| Bougainvillea | Ligustrum |
| Camellia | Magnolia |
| Euonymus | Photinia |
| Hydrangea | Pine |
| Ilex | Rhododendron |

BONZI® is a trademark of a ZENECA Group Company.