

PM-23

Reg # 10182-390 NOTIFICATION

F4C

1920

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number
227318

Application for Pesticide - Section I

1. Company/Product Number 10182-390	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) FLEXSTAR Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) ZENECA Ag Products 1800 Concord Pike, PO Box 15458 Wilmington, DE 19850-5458 <input type="checkbox"/> Check if this is a new address.	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
 Notifying the Agency that based on the approved stamped Flexstar Label, we have redefined our production label in a manner that more precisely outlines geographic and marketing needs. Required for Zeneca State Registration Specialists to facilitate the registration of Flexstar in approved states. (North and South Regions have been precisely defined.)

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Saundra M. O'Bryan	Title Regulatory Product Lead	Telephone No. (Include Area Code) 302-886-8789
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Saundra M. O'Bryan</i>	3. Title Regulatory Product Lead	
4. Typed Name Saundra M. O'Bryan	5. Date February 7, 1996	

2/20

**FLEXSTAR™
Herbicide**

For Postemergence Control of Broadleaf Weeds in Soybeans

COMPLETE DIRECTIONS FOR USE

**CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

5 y 20

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ACTIVE INGREDIENT:

Sodium salt of fomesafen	
5-[2-chloro-4-(trifluoromethyl)phenoxy]-N-(methylsulfonyl)-2-nitrobenzamide*	12.8%
INERT INGREDIENTS	87.2%
TOTAL	100.0%

*Equivalent to 12.2% fomesafen or 1 pound fomesafen active ingredient per gallon.

EPA Reg. No. 10182-390

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

IF INHALED: Removed victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-I-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

This product contains fomesafen which has been determined to cause tumors in laboratory animals (mice). Risks can be reduced by closely following use directions and precautions and by wearing the protective clothing specified elsewhere on this label.

CAUSES SKIN IRRITATION. HARMFUL IF INHALED OR ABSORBED THROUGH SKIN. Do not get on skin or on clothing. Avoid breathing vapor or spray mist. Avoid contact with eyes. Prolonged or repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants.
- Chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton.
- Chemical resistant footwear plus socks.
- Chemical resistant apron when cleaning equipment, mixing or loading.


User Safety Recommendations

Users should:

- - Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.



7920

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants.
- Chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton.
- -Chemical resistant footwear plus socks.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Store above 45°F in original containers only. If product solidifies, return to room temperature and agitate to reconstitute. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

8925

CONTAINER DISPOSAL:

Metal Containers: Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Plastic Containers: Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Glass Containers: Triple rinse (or equivalent); then dispose of in a sanitary landfill or by other approved state and local procedures.

FOR BULK AND MINI-BULK CONTAINERS:

Container Disposal: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

Container Precautions: Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.

REFILL ONLY WITH FLEXSTAR . The contents of this container cannot be completely removed by cleaning. Refilling with materials other than FLEXSTAR will result in contamination and may weaken container.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

GENERAL INFORMATION

Read all label directions before using.

FLEXSTAR is a selective postemergence herbicide for control of broadleaf weeds in soybeans. Soybean plants are tolerant to FLEXSTAR when applied at recommended rates. Some bronzing, crinkling or spotting of soybean leaves may occur, but soybeans soon outgrow these effects and develop normally.

FLEXSTAR is effective through contact action, therefore weeds must be thoroughly covered with spray. Optimum weed control is achieved when young actively growing weeds are treated that are not under stress from moisture, temperature, low soil fertility, mechanical or chemical injury.

Secondary flushes of certain weeds may be controlled by soil residual activity if rainfall occurs shortly after application. The extent of soil activity is dependent upon soil type, ground cover at time of application, amount of rainfall and the rate of FLEXSTAR used.

FLEXSTAR is rainfast in 4 hours.

APPLICATION DIRECTIONS

TIMING: Best control of susceptible broadleaf weeds is obtained when FLEXSTAR is applied early to actively growing weeds. This usually occurs 14 to 28 days after planting. Refer to the weed tables for specific recommendations on weed growth stages, rates, and regions.

SPRAY ADDITIVES: FLEXSTAR is formulated with an optimized surfactant system for herbicidal performance.

In the Northern region (see Regional Use Map), FLEXSTAR should be used with a minimum of 1% of liquid nitrogen (28% or similar) or a minimum of 4 pounds ammonium sulfate per 100 gallons of spray volume.

The use of FLEXSTAR with additional adjuvants, such as crop oil concentrate or nonionic surfactant can improve performance on larger and/or more difficult to control weeds or during drought stress or when spray volume exceeds 20 gallons per acre.. Under certain conditions temporary crop injury may result.

GUIDELINES FOR USING ADDITIONAL ADJUVANTS: Only spray additives cleared for use on growing crops under 40 CFR 180.1001 may be used.

Nonionic Surfactant (NIS) - Use NIS containing at least 75% surface active agent at 0.25% (1/2 pint per 25 gallons) of the finished spray volume.

Crop Oil Concentrate (COC) - Use a nonphytotoxic COC or a once refined vegetable oil concentrate (VOC, MSO) containing 15-20% approved emulsifier at 0.5% (1 pint per 25 gallons) of finished spray volume.

Other Adjuvants - Adjuvants other than COC or NIS may be used providing the product meets the following criteria:

- 1. Contains only EPA exempt ingredients.
- 2. Is nonphytotoxic to the target crop.
- 3. Is compatible in mixture. (May be established through a jar test.)
- 4. Is supported locally for use with FLEXSTAR on the target crop through proven field trials and through university and extension recommendations.

Tankmixes: If a product tankmixed with FLEXSTAR requires a cropoil concentrate or nonionic surfactant, use the lowest recommended adjuvant rate for the tankmix product. (See Tankmix Application section for further details.)

Recommended Mixing Order:

- 1. Half required amount of water, begin agitation.*
- 2. Dry pesticide formulations.
- 3. FLEXSTAR herbicide.
- 4. Liquid pesticide formulation.
- 5. Adjuvant (COC or NIS) and fertilizer.

*Compatibility agent, 1 gallon/500 gallons of water or 0.2% v/v, may be added as needed.

10 9 20

GROUND APPLICATION: Use sufficient spray volume and pressure to ensure complete coverage of the target weeds. A spray volume of 10-20 gallons per acre and 30-60 psi at the nozzle tip is recommended. When weed foliage is dense, use 60 psi and a minimum of 20 gallons per acre to ensure coverage of weed foliage.

Use only hollow cone or flat fan nozzles. The sprayer must be calibrated to provide the proper volume and rate per acre. In addition, the boom and nozzle height must be adjusted to provide complete coverage of all weeds.

DO NOT USE FLOOD TYPE OR OTHER SPRAY NOZZLES, WHICH DELIVER COARSE, LARGE DROPLET SPRAYS.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

BAND APPLICATIONS: Thorough weed coverage is important for control. Best coverage is obtained with a minimum of two nozzles, one directed to each side of the planted row. Application with a single nozzle directed over the top of the row is not recommended. Cultivation of untreated areas may be needed following band applications. When making band applications and cultivating in the same operation, position nozzles in advance of the cultivation device. This will reduce dust in the spray area. Dust can intercept spray, reducing weed coverage, resulting in less than adequate weed control.

Calculate the amount of herbicide and water volume needed for band treatment by the following formulas:

$$\frac{\text{Band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{Band herbicide rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{row width in inches}} \times \text{broadcast volume per acre} = \text{Band water volume per acre}$$

AERIAL APPLICATION: Use sufficient spray volume and pressure to ensure complete coverage of the target weeds. A minimum of 5 gallons per acre of spray mixture should be applied with a maximum of 40 PSI pressure. When broadleaf weed foliage is dense, use a minimum of 10 gallons per acre to ensure coverage of weed foliage.


CULTIVATION: Cultivation prior to application is not recommended. Cultivation may put weeds under stress, reducing weed control. Timely cultivation 1-3 weeks after applying FLEXSTAR may assist weed control.

GENERAL USE PRECAUTIONS

- A maximum of 3 pints/A of FLEXSTAR herbicide may be used per season in Region 1 (see Regional Use Map).
- A maximum of 3 pints/A of FLEXSTAR herbicide may be applied in alternate years in Region 2 (see Regional Use Map).
- A maximum of 2.5 pints/A of FLEXSTAR herbicide may be applied in alternate years in Region 3 (see Regional Use Map).

11 9 20
7

- A maximum of 2.0 pints/A of FLEXSTAR herbicide may be applied in alternate years in Region 4 (see Regional Use Map).
- Do not apply FLEXSTAR or any other fomesafen containing product to any field in Regions 2, 3 or 4 more than once every two years.
- Apply FLEXSTAR herbicide before soybeans bloom.
- It is mandatory that the spray system be thoroughly cleaned with water and a commercial tank cleaner before and after each use.
- Tankmixes of FLEXSTAR herbicide with other pesticides, fertilizers or any other additives except as specified on this label or other approved ZENECA supplemental labels may result in tankmix incompatibility, unsatisfactory performance and/or unsatisfactory crop injury.
- FLEXSTAR herbicide should not be mixed with fertilizers or any other additive except as specified on this label or other approved ZENECA supplemental labeling.
- FLEXSTAR herbicide requires a 4-hour rain-free period for best results.
- Apply to actively growing weeds. Avoid applying FLEXSTAR to weeds or soybeans which are under stress from moisture, temperature, low soil fertility, mechanical or chemical injury, as reduced weed control and/or increased crop injury may result.
- Avoid overlapping spray swaths, as injury may occur to rotational crops.
- Avoid drift to all other crops and nontarget areas. Crops other than soybeans may be severely injured by drift. Application is prohibited when wind velocity exceeds 10 mph. Do not make ground or aerial application during temperature inversions.
- To provide adequate coverage, it is recommended that ground speed not exceed 10 MPH during application.
- Do not graze treated areas or harvest for forage or hay.
- Do not apply this product through any type of irrigation system.



12 of 20

ROTATIONAL CROP RESTRICTIONS


The following rotational crops may be planted after applying FLEXSTAR at recommended rates in soybeans:

<u>Crop To Be Planted</u>	<u>Minimum Rotation Interval (Months After Last REFLEX Application)</u>
Small grains such as wheat, barley, rye	4
Alfalfa, beans & peas, corn*, cotton, peanuts, rice	10
To avoid crop injury do not plant sunflowers, sugar beets, sorghum** or any other crop within	18

Do not graze rotated small grain crops or harvest forage or straw for livestock. In the event of a crop loss due to weather conditions soybeans can be replanted.

* Use 18 month minimum rotation interval for sweet corn in the states of Maine, Massachusetts, New Hampshire, New York, Rhode Island and Vermont.

** Sorghum may be planted back after 10 months in Region 1.



FLEXSTAR IN SOYBEANS - USE RATES AND WEEDS CONTROLLED

**REFER TO MAP FOR DEFINITION OF SPECIFIED
GEOGRAPHIC REGIONS**

REGION 1

REGION 1 - Includes the following states or portion of states where FLEXSTAR may be applied: Alabama, Arkansas, Georgia, Louisiana, Mississippi, Missouri (counties of Butler, Dunklin, Mississippi, New Madrid, Pemiscot, Scott and Stoddard), North Carolina, Oklahoma (east of U.S. Highway 75 and east of Indian Nation Parkway), South Carolina, Tennessee, and Texas (all areas east of Highway 77 to State Road 239, including all of Calhoun county).

(Insert Region 1 Use Map Here)

REGION 2

REGION 2 - Includes the following states where FLEXSTAR may be applied: Delaware, Kentucky, Maryland, and Virginia.

(Insert Region 2 Use Map Here)

REGION 3


REGION 3 - Includes the following states or portion of states where FLEXSTAR may be applied: Indiana, Illinois, Iowa, Maine, Massachusetts, Missouri (all counties except for those listed in Region 1), New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, West Virginia, and Wisconsin (South of U.S. Highway 18 between Prairie Du Chien and Madison, and South of Interstate 94 between Madison and Milwaukee).

(Insert Region 3 Use Map Here)

REGION 4

REGION 4 - Includes the following states or portion of states where FLEXSTAR may be applied: Kansas (all counties east of U.S. Highway 75), Michigan (all counties south of U.S. Highway 55), Minnesota (all counties south of U.S. Highway 212), Nebraska (all counties east of U.S. Highway 81), and Wisconsin (North of U.S. Highway 18 between Prairie Du Chien and Madison, and South of Interstate 94 from Hudson to Madison).

(Insert Region 4 Use Map Here)



14920

APPLICATION RATES FOR WEED GROWTH STAGES

Weed	FLEXSTAR Rate (Pts/A)		
	Maximum Growth Stage Controlled At		
	2 Pts/A # of True Leaves	2.5 Pts/A # of True Leaves	3 Pts/A # of True Leaves
Anoda, Spurred	—	2	2
Carpetweed	6" Diameter Size	Multi-leaf 6" Diameter	Unlimited Size
Citron (Wild Watermelon)	2	2	4
Cocklebur, Common	—	6	8
Copperleaf, Hophornbeam	2	2	4
Copperleaf, Virginia	2	2	4
Crotalaria, Showy	4	4	6
Croton, Tropic	2	2	4
Cucumber, Volunteer	4	4	6
Eclipta	2	2	4
Groundcherry, Cutleaf	2	4	6
Jimsonweed	4	6	8
Ladysthumb	2	4	6
Lambsquarters, Common	2*	2*	2*
Mexicanweed	—	2*	2
Morningglory			
Cypressvine	4	4	6
Entireleaf var.	2	3	5
Ivyleaf	2	3	5
Purple Moonflower	2	4	6
Red (Scarlet)	3	6	6
Smallflower	2	4	6
Pitted (Smallwhite)	3	6	6
Tail (Common)	2	3	5
Palmleaf (Willowleaf)	3	6	6
Mustard, Wild	4	6	8
Nightshade, Black	4	4	4

Weed	FLEXSTAR Rate (Pts/A)		
	Maximum Growth Stage Controlled At		
	2 Pts/A # of True Leaves	2.5 Pts/A # of True Leaves	3 Pts/A # of True Leaves
Nutsedge, Yellow	-	-	*
Pigweed, spp.			
Amaranth, Palmer	4	6	6
Amaranth, Spiny	-	4	4
Redroot	4	6	8
Smooth	4	4	6
Waterhemp, Common	4	4	6
Waterhemp, Tall	-	4	6
Poinsettia, Wild	2	4	6
Purslane, Common	Multi-Leaf 6" Diameter	Multi-Leaf 6" Diameter	Multi-Leaf 8" Diameter
Pusley, Florida	-	2	4
Ragweed, Common	4	6	8
Ragweed, Giant	4	6	8
Redweed	-	-	3*
Sesbania, Hemp	6	6	12
Sicklepod	-	-	Cotyledon*
Sida, Prickly	-	2	2
Smartweed, Pennsylvania	4	6	6
Smellmelon	2	2	4
Spurge, Prostrate	-	1" Diameter*	1" Diameter*
Spurge, Spotted	-	2*	2*
Starbur, Bristly	2	4	6
Sunflower, Common**	-	4	6
Velvetleaf	-	4	6
Venice Mallow	4	4	6
Witchweed	-	Multi-leaf Up to 7"	Multi-leaf Up to 10"
Yellow Rocket	4	6	6

*Suppression Only

** Regions 2 and 3 only


Under certain conditions, the mixture of FLEXSTAR with one of the above mentioned broadleaf herbicides may cause a reduction in activity of any postemergence grass herbicide in the mixture.

It is recommended no additional adjuvants be added to post grass herbicides plus FLEXSTAR tankmixtures for control of perennial grasses.

For sequential applications allow 2-3 days after the application of the grass herbicide before applying FLEXSTAR or FLEXSTAR mixtures. In case FLEXSTAR or the FLEXSTAR mixture is applied first apply the grass herbicide when the grass weeds begin to develop new leaves (generally around 7 days).

- Tankmix applications can result in increases in crop injury as compared to either product used alone.
- It is recommended no additional adjuvants be added to FLEXSTAR plus Pinnacle or Pinnacle containing premixes.
- Do not exceed 1 fluid ounce of Butyrac per acre in mixture with FLEXSTAR.

Always read and follow the restrictions and limitations for all products whether used alone, sequentially or in a tankmix. The most restrictive labelling of any product used applies.



APPENDIX

Scientific names are listed for those weeds referred to in the FLEXSTAR label.

COMMON NAME	SCIENTIFIC NAME
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Spiny	<i>Amaranthus spinosus</i>
Anoda, Spurred	<i>Anoda cristata</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bindweed, Field	<i>Convolvulus arvensis</i>
Bindweed, Hedge	<i>Calystegia sepium</i>
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>
Carpetweed	<i>Mollugo verticillata</i>
Citron (Wild Watermelon)	<i>Citrullus vulgaris</i>
Cocklebur, Common	<i>Xanthium strumarium</i>
Copperleaf, Hophornbeam	<i>Acalypha ostryiifolia</i>
Copperleaf, Virginia	<i>Acalypha virginica</i>
Crabgrass	<i>Digitaria spp.</i>
Crotalaria, Showy	<i>Crotalaria spectabilis</i>
Croton, Tropic	<i>Croton glandulosus</i>
Cucumber, Volunteer	<i>Cucumis sativas</i>
Eclipta	<i>Eclipta prostrata</i>
Foxtail, Giant	<i>Setaria faberi</i>
Foxtail, Green	<i>Setaria viridis</i>
Foxtail, Yellow	<i>Setaria glauca</i>
Goosegrass	<i>Eleusine indica</i>
Groundcherry, Cutleaf	<i>Physalis angulata</i>
Jimsonweed	<i>Datura stramonium</i>
Johnsongrass, Seedling	<i>Sorghum halepense</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, Common	<i>Chenopodium album</i>

19 9 20

COMMON NAME	SCIENTIFIC NAME
Mexicanweed	<i>Caperonia castaniifolia</i>
Milkweed, Climbing	<i>Sarcostemma cyanchoides</i>
Milkweed, Honeyvine	<i>Ampelamus albidus</i>
Morningglory, Cypressvine	<i>Ipomoea quamoclit</i>
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integruscula</i>
Ivyleaf	<i>Ipomoea hederacea</i> var. <i>hederacea</i>
Purple Moonflower	<i>Ipomoea turbinata</i>
Red (Scarlet)	<i>Ipomoea coccinea</i>
Smallflower	<i>Jacquemontia tamnifolia</i>
Pitted (Smallwhite)	<i>Ipomoea lacunosa</i>
Tall (Common)	<i>Ipomoea purpurea</i>
Palmleaf (Willowleaf)	<i>Ipomoea wrightii</i>
Mustard, Wild	<i>Brassica kaber</i>
Nightshade, Black	<i>Solanum nigrum</i>
Nutsedge, Yellow	<i>Cyperus esculentus</i>
Panicum, Fall	<i>Panicum dichotomiflorum</i>
Panicum, Texas	<i>Panicum texanum</i>
Pigweed, Redroot	<i>Amaranthus retroflexus</i>
Pigweed, Smooth	<i>Amaranthus hybridus</i>
Poinsettia, Wild	<i>Euphorbia heterophylla</i>
Purslane, Common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Ragweed, Common	<i>Ambrosia artemisiifolia</i>
Ragweed, Giant	<i>Ambrosia trifida</i>
Redweed	<i>Melochia corchorifolia</i>
Sesbania, Hemp	<i>Sesbania exaltata</i>
Sicklepod	<i>Cassia obtusifolia</i>
Sida, Prickly	<i>Sida spinosa</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Smellmelon	<i>Cucumis melo</i>
Spurge, Prostrate	<i>Euphorbia humistrata</i>
Spurge, Spotted	<i>Euphorbia maculata</i>

COMMON NAME	SCIENTIFIC NAME
Starbur, Bristly	<i>Acanthospermum hispidum</i>
Sunflower, Common	<i>Helianthus annuus</i>
Trumpetcreeper	<i>Campsis radicans</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Venice Mallow	<i>Hibiscus trionum</i>
Waterhemp, Common	<i>Amaranthus rudis</i>
Waterhemp, Tall	<i>Amaranthus tuberculatos</i>
Witchweed	<i>Striga asiatica</i>
Yellow Rocket	<i>Barbarea vulgaris</i>

FUSILADE[®], FUSION[®], and FLEXSTAR[™] are trademarks of ZENECA Group Companies.
 Basagran[®], Poast[®] and Poast Plus[™] are trademarks of BASF Corporation.
 Assure II[®], Classic[®], Concert[®], Pinnacle[®] and Synchrony STS[®] are trademarks of E.I.
 DuPont de Nemours & Co., Inc.
 Pursuit[®] and Scepter[®] are trademarks of American Cyanamid Company.
 Select[®] is a trademark of Valent Chemical Co.
 Bugle[™] and Option[®] II are trademarks of AgrEvo.
 Butyrac[®] is a trademark of Rhone Poulenc Ag Company.