



010182-00400-061099

Printed: 15:47:37 Tuesday, 20 Jul, 1999 # 38 / 3613

**Systems Integration Group, Inc.**

PM 03

10182-400

06/10/99

JUN 10 1999

Mr. S. K. Theodorakis  
Regulatory Product Lead  
Zeneca Ag Products  
1800 Concord Pike  
P.O. Box 15458  
Wilmington, DE 19850-5458

Subject: Amendment: Add use on ornamentals in greenhouses  
Scimitar GC Insecticide  
EPA Registration No. 10182-400  
Final Label Dated January 18, 1999

Dear: Mr. Theodorakis:

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data.
2. Move the highlighted statements "Do not apply...for pesticide regulation." found on page 6 of your label to a position directly under the DIRECTIONS FOR USE section.
3. Add the following statement to the AGRICULTURAL USE REQUIREMENTS section: "Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas."
4. Under the 3<sup>rd</sup> paragraph of the Environmental Hazards section, add the spray drift statement that appears on page 9 of the label "When treating tall trees..."
5. Submit two (2) copies of your final printed labeling before you release the product for shipment.

		CONCURRENCES					
SYMBOL	7505C						
SURNAME	SFRANT						
DATE	6/10/99						

A stamped copy of the label is enclosed for your records. If you have any questions, please call Dr. William Sproat of my team at 703-308-8587.

Sincerely,

George T. LaRocca  
Product Manager 13  
Insecticide Branch  
Registration Division (7505C)

Enclosure

)

)

3 8 14

**RESTRICTED USE PESTICIDE**  
**Due to Toxicity to Fish and Aquatic Organisms**

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**SCIMITAR® GC Insecticide**

For Use to Control Insect Pests On Golf Courses and Athletic Field Turf; Sod Farms; Ornamentals Grown in Commercial Greenhouses, Shadehouses and Nurseries; and on Ornamentals and Lawns in Landscaped Areas Around Residential, Institutional, Public, Commercial and Industrial Buildings, Parks, Recreational Areas, Golf Courses and Athletic Fields

**ACTIVE INGREDIENT:**

Lambda-cyhalothrin*	
[1α(S*),3α(Z)]-(±)-cyano(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate	9.7%
<b>INERT INGREDIENTS:</b>	<u>90.3%</u>
<b>Total</b>	<b>100.0%</b>

\*Synthetic pyrethroid.  
Contains 0.88 lbs active ingredient per gallon or 0.22 lbs ai per quart.

EPA Reg. No. 10182-400  
EPA Est. No.

Net Contents:  
ACCEPTED  
with COMMENTS  
in EPA Letter Dated

**KEEP OUT OF REACH OF CHILDREN**

**JUN 10 1999**

**CAUTION**

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 10182-400

See Back Panel for Statement of Practical Treatment and Precautionary Statements

Made in UK; Packaged in U.S.A.  
ZENECA Professional Products  
ZENECA Inc.  
Wilmington, DE 19850-5458

**CONDITIONS OF SALE  
AND LIMITATION OF WARRANTY AND LIABILITY**

---

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

---

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

**STATEMENT OF PRACTICAL TREATMENT**

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF SWALLOWED:** If victim is alert and not convulsing, rinse mouth out and give 200-300 mL (1 cup) of water to dilute material. Immediately contact local Poison Control Center. Vomiting should only be induced under the direction of a physician or Poison Control Center. If spontaneous vomiting occurs, have victim lean forward with head down to avoid breathing in of vomitus, rinse mouth and administer more water. Immediately transport victim to an emergency facility.

**IF IN EYES:** Flush with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

---

**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).**

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.**

---

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

HARMFUL IF ABSORBED THROUGH SKIN. PROLONGED OR FREQUENTLY REPEATED SKIN CONTACT MAY CAUSE ALLERGIC SKIN REACTIONS IN SOME INDIVIDUALS. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist or vapors. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Skin exposure may also result in a sensation described as a tingling, itching, burning or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 to 30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

**Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or viton
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

---

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
  - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
  - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- 

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label. Care should be used when making applications to avoid household pets, particularly fish and reptile pets.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds. In the state of New York, do not apply within 100 feet of coastal marshes or streams that drain into marshes.

~~Do not make applications when wind speed is 15 miles per hour or greater.~~

Notify State and/or Federal authorities and ZENECA Inc. immediately if you observe any adverse environmental effects due to the use of this product.

### PHYSICAL AND CHEMICAL HAZARDS

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

~~IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns or grounds.~~

---

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
  - Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or viton
  - Shoes plus socks
- 

SCIMITAR® GC may be used for applications to ornamentals grown in commercial greenhouses, shadehouses and nurseries, and turf grown on sod farms or for commercial seed production.

---

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep adults, children and pets off treated areas until spray has dried following the application.

---



---

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**STORAGE:** Keep container closed when not in use. Do not store near food or feed. Shake well before use. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste storage area until proper disposal can be made.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Completely empty container into application equipment, then offer for recycling or reconditioning, or puncture and dispose of empty container in a sanitary landfill, by incineration, or by other procedures approved by state and local authorities.

---

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

SCIMITAR GC may be used for applications to maintain indoor or outdoor areas where turf and ornamentals are grown such as residential landscaped areas and non-residential landscapes around institutional, public, commercial and industrial buildings, parks, recreational areas, golf courses and athletic fields.

SCIMITAR GC may also be used for applications to golf course fairways, greens, greens aprons and tee areas. ~~Not for use on sod farms or grass grown for seed.~~

**IMPORTANT:** Time application to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

DO NOT APPLY THIS PRODUCT TO EDIBLE CROPS OR CROPS GROWN FOR FOOD/FEED.

DO NOT APPLY THIS PRODUCT BY AERIAL APPLICATION.

**SPRAY DRIFT PRECAUTIONS:** Do not make applications when wind speed is 15 miles per hour or greater. Low humidity and high temperatures increase the likelihood of spray drift to sensitive areas. Avoid spraying during conditions of low humidity and/or high temperature. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when the wind direction is towards the aquatic area. Do not make outdoor applications during temperature inversions. Inversions are characterized by stable air and increasing temperature with height above ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**MIXING:** SCIMITAR GC insecticide is intended for dilution with water for spray application and may be used in all types of application equipment. Fill application tank with  $\frac{1}{2}$  to  $\frac{3}{4}$  volume of water. It is suggested that the water be 5 to 7 pH. Adjust water pH with a buffering agent if necessary. Slowly add SCIMITAR GC to applicator tank water with maximum agitation. If other chemicals are added to the applicator tank, SCIMITAR GC should be added last. Do not mix if mixed with EC formulations or oils, use within 24 hours. Fill tank to desired volume and continue to agitate while making applications. Shake or reagitae application equipment before use if application is interrupted. Make up only amount of application volume as required.

**COMPATIBILITY:** SCIMITAR GC has been found to be compatible with all commonly used fungicides, miticides, liquid fertilizers, and other insecticides. Check physical compatibility using the correct proportion of products in a small jar if local experience is unavailable.

**APPLICATION:** SCIMITAR GC insecticide mixes readily with water and may be used in all types of application equipment. Mix product with the required amount of water and apply as a dilute application to the point of runoff. Apply product using spray nozzles which produce a coarse droplet size. Formation of very small droplets may be minimized by appropriate nozzle selection and by avoiding excessive spray pressure. When applying to hard-to-wet foliage such as holly, pine, or ivy, the addition of a spreader sticker may enhance knockdown and residual activity. If concentrate or mist-type application equipment is used, an equivalent amount of product should be used on the area treated as would be used in a dilute application. Phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed. It is advised to pretreat a selection of ornamentals if local use experience is unavailable.

**INSECT PESTS CONTROLLED IN ORNAMENTALS BY SCIMITAR GC INSECTICIDE**

USE	PEST	AMOUNT OF SCIMITAR GC
<p><b>AGRICULTURAL</b>            ORNAMENTALS IN GREENHOUSES, SHADEHOUSES AND NURSERIES</p> <p><b>NON-AGRICULTURAL</b>            ORNAMENTALS (Including Trees, Shrubs, Flowers, Evergreens, Foliage Plants and Groundcovers) IN RESIDENTIAL LANDSCAPED AREAS AND LANDSCAPED AREAS AROUND INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, GOLF COURSES, AND ATHLETIC FIELDS</p>	<p>Ants (Including Imported fire ants)            Armyworms            Azalea caterpillars            Aphids            Bagworms            Black vine weevils (adult)            Boxelder bugs            Budworms            California oakworms            Cankerworms            Cockroaches            Crickets            Cutworms            Eastern Tent caterpillars            Elm leaf beetles            European sawflies            Fall webworms            Flea beetles            Forest tent caterpillars            Gypsy moth larvae            Japanese beetles (adults)            June beetles (adults)            Lace bugs            Leaf-feeding caterpillars            Leafhoppers            Leafminers (adults)            Leaf rollers            Leaf skeletonizers            Midges            Mosquitoes            Oleander moth larvae            Pillbugs            Pine sawflies            Pine shoot beetles            Pinetip moths            Plant bugs            Root weevils            Sawflies            Scale insects (crawlers)            Spiders            Spittlebugs            Striped beetles            Striped oakworms            Thrips            Tip moths            Tussock moth larvae            Wasps</p>	<p>1.5 to 5 fl oz/            100 gals            or            44 to 148 mL/            100 gals</p>
	<p>Broadmites            Brown softscales            California redscales (crawler)            Clover mites            Mealybugs            Pineneedlescales (crawler)            Spider mites            Whiteflies</p>	<p>3 to 5 fl oz/            100 gals            or            88 to 148 mL/            100 gals</p>

Application to ornamentals should be started prior to the establishment of high insect pest populations. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Apply at 7-day intervals if treatment is necessary.

Do not apply more than 0.36 pounds of the active ingredient (52.4 fl oz of concentrate) per acre per year.

When treating tall trees (> 15 feet from the ground) with high pressure sprays or during any application with air assisted equipment (mist blowers), do not apply within 150 feet of aquatic areas.

Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at recommended rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage.

For spot treatments use 0.5 fl oz SCIMITAR GC per 1 to 2 ½ gallons of water.

Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.

**NOTE:** While phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed, certain cultivars may be sensitive to the final spray solution.

It is advised to prespray a selection of ornamental plants and observe them for 7 to 10 days prior to treating large areas if local use experience is unavailable.

**Scale.** Thoroughly cover the plant with SCIMITAR GC spray, including trunks, stems, twigs and foliage for control of scale insects.

**Bagworm.** Apply SCIMITAR GC when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.

**SCIMITAR GC MIXING CHART FOR ORNAMENTAL INSECT PEST CONTROL**  
(Scimitar GC to add per spray tank)

Rate of Scimitar GC	25 Gallons	50 Gallons	100 Gallons	200 Gallons	300 Gallons
1.5 oz	0.4 oz	0.8 oz	1.5 oz	3.0 oz	4.5 oz
3.0 oz	0.8 oz	1.5 oz	3.0 oz	6.0 oz	9.0 oz
5.0 oz	1.3 oz	2.5 oz	5.0 oz	10.0 oz	15.0 oz

**INSECT PESTS CONTROLLED ON TURFGRASS BY SCIMITAR GC INSECTICIDE**

USE	PEST	AMOUNT OF SCIMITAR GC
<b>AGRICULTURAL</b> SOD FARMS AND GRASS SEED FARMS  <b>NON-AGRICULTURAL</b> LAWNS AROUND RESIDENTIAL, INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, GOLF COURSES AND ATHLETIC FIELDS  GOLF COURSE AND ATHLETIC FIELD TURF	Ants (Including Imported fire ants) Armyworms Centipedes Crickets Cutworms Earwigs Fleas (adult) Grasshoppers Japanese beetles (adult) Millipedes Mites Pillbugs Sod webworms Sow bugs Ticks (including species which transmit Lyme disease)	3.4 to 7 mL/1000 sq ft or 5 to 10 fl oz/acre
	Bluegrass billbugs (adult) Black turfgrass ataenius (adult) Chiggers Fleas (adult) Grub (suppression) Hyperodes weevils (adult) Mole crickets (nymphs and young adults)	7 mL/1000 sq ft or 10 fl oz/acre
	Chinch bugs Mole Crickets (mature adults) (Not for use on mature adult mole crickets and chinch bugs in New York state.)	14 mL/1000 sq ft or 20 fl oz/acre

Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Apply at 7-day intervals if treatment is necessary.

Do not apply more than 0.36 pounds of active ingredient (52.4 fl oz of concentrate) per acre per year.

For spot treatments use 0.5 fl oz of SCIMITAR GC per 1 to 2½ gallons of water.

Do not apply when turfgrass is waterlogged or when soils are saturated with water (i.e. will not accept irrigation).

Keep children and pets off treated areas until spray has dried following the application.

**Surface Insect Control (armyworm, cutworms, fleas, etc.)** For best results apply SCIMITAR GC at recommended rates in 2 to 5 gallons of water per 1,000 sq. ft. The use of a spreader-sticker may be useful if high rainfall amounts are forecast, otherwise the addition of adjuvants is not necessary under normal conditions for surface insect control in turf. Delay watering or mowing for 12 to 24 hours for optimum control of surface-feeding insect pests.

**Thatch Inhabiting Insect Control (chinch bugs, billbugs, etc.)** For best results apply SCIMITAR GC at recommended rates in 2 to 10 gallons of water per 1,000 sq. ft. The use of a non-ionic wetting agent, penetrant or similar adjuvant is recommended at label rates. Lightly irrigate after application with up to ½ inch of water to move the SCIMITAR GC into the thatch layer. If irrigation is not available, then use high water application rates for optimum results.

**Subsurface Insect Control (mole crickets, grubs, etc.)** For best results apply SCIMITAR GC at recommended rates in 4 to 10 gallons of water per 1,000 sq. ft. The use of a non-ionic wetting agent, penetrant or similar adjuvant is strongly recommended following label rates. Use the highest water application rates possible with your sprayer. Apply SCIMITAR GC to turf wet with dew, rain or irrigation. Water-in immediately after application with ¼ to ½ inch of water for optimum results.

**Fire Ant Control** Treat individual mounds with a drench application using a watering can. Use 0.5 fl oz of SCIMITAR GC per 2.5 gallons of water. Thoroughly soak each mound and a 3 foot diameter circle around each mound. Gently apply the mixture to avoid disturbing the mound. Disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. For best results, apply in early morning or late evening hours. Make additional treatments as needed.

**Mosquito Control** Apply as a general spray around landscape plantings, turf, and building foundations to control mosquitoes. For best results apply SCIMITAR GC at recommended rates in 2 to 5 gallons of water per 1,000 sq. ft.

**SCIMITAR GC Mixing Chart for Turf Insect Pest Control**  
(Scimitar GC to add per 100 gallon spray tank)

Rate of Scimitar GC	Application Rate Per 1,000 Square Feet of Turf				
	2 Gallons	4 Gallons	6 Gallons	8 Gallons	10 Gallons.
5 oz/acre	5.7 oz	2.9 oz	1.9 oz	1.4 oz	1.2 oz
10 oz/acre	11.5 oz	5.7 oz	3.8 oz	2.9 oz	2.3 oz
20 oz/acre	23.0 oz	11.5 oz	7.7 oz	5.7 oz	4.6 oz

Conversion Rate: 1 fluid ounce (fl oz) equals 29 milliliters (mL).

SCIMITAR® is a trademark of a ZENECA Group Company.