

10182-404

02/18/2000

1/32



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
10182-404

Date of Issuance:
FEB 18 2000

NOTICE OF PESTICIDE:
Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:
Unconditional

Name of Pesticide Product:
Reward Landscape and Aquatic Herbicide

Name and Address of Registrant (include ZIP Code):

Zeneca Ag Products
P.O. Box 15458
Wilmington, DE 19850-5458

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above. The following revision must be incorporated into the labeling before the product is released for shipment:

Revise the second sentence of the dermal statement in the Statement of Practical Treatment to read "Get medical attention."

Enclosed is a copy of your label stamped "Accepted with Comments". This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain registration of your product.

Signature of Approving Official:

Date:

FEB 18 2000

EPA Form 8570-8
7505C
D. KENNY
2/18/2000
PK:305-7546:HB/PM23

REWARD[®]
Landscape and Aquatic Herbicide

**TO PREVENT ACCIDENTAL POISONING, NEVER PUT INTO FOOD, DRINK OR
OTHER CONTAINERS AND USE STRICTLY IN ACCORDANCE WITH ENTIRE LABEL**

DO NOT USE THIS PRODUCT FOR REFORMULATION

ACTIVE INGREDIENT:

Diquat dibromide [6,7-dihydrodipyrido (1,2-a:2',1'-c) pyrazinediium dibromide]	37.3%
INERT INGREDIENTS	<u>62.7%</u>
TOTAL	100.0%

Contains 2 lbs. diquat cation per gal. as 3.73 lbs. salt per gal.

EPA Reg. No. 10182-404
EPA Est. No.

Net Contents:

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See (inside, back or side panels) for
Statement of Practical Treatment and Precautionary Statements

**ACCEPTED
with COMMENTS
In EPA Letter Dated**
FEB 18 2000
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
10182-404

Made in U.S.A.
ZENECA Professional Products
ZENECA Ag Products Inc.
Wilmington, DE 19850-5458

**CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.


ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

TABLE OF CONTENTS

	Page
STATEMENT OF PRACTICAL TREATMENT	
PRECAUTIONARY STATEMENTS	
HAZARDS TO HUMANS AND DOMESTIC ANIMALS	
PERSONAL PROTECTIVE EQUIPMENT	
ENVIRONMENTAL HAZARDS	
DIRECTIONS FOR USE	
DIRECTIONS	
AGRICULTURAL USE REQUIREMENTS	
NON-AGRICULTURAL USE REQUIREMENTS	
AQUATIC USE DIRECTIONS	
STORAGE AND DISPOSAL	



STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: IMMEDIATELY give water or milk to drink and induce vomiting by inserting finger in throat. Do not induce vomiting or give anything by mouth to an unconscious person. Take person and product container to the nearest hospital or physician fast. **PROMPT TREATMENT IS ESSENTIAL TO COUNTERACT POISONING** and should be initiated before signs and symptoms of injury appear.

IF ON SKIN: IMMEDIATELY wash with plenty of soap and water. See a doctor if diquat contacts a skin cut, abrasion or area of irritation.

IF IN EYES: IMMEDIATELY wash eyes with water for at least hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. ~~and g~~ Get medical attention.

IF INHALED: IMMEDIATELY remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention. ~~get away from spray mist. Stop and check spray procedure. See a doctor if irritation persists.~~

NOTE TO PHYSICIANS: Call ZENECA Medical Emergency Information Network 1-800-F-A-S-T-M-E-D (327-8633) at any hour to obtain toxicology information and a diquat analysis. To be effective, treatment for diquat poisoning must begin **IMMEDIATELY**. Treatment consists of binding diquat in the gut with suspensions of activated charcoal or bentonite clay, administration of cathartics to enhance elimination, and removal of diquat from the blood by charcoal hemoperfusion or continuous hemodialysis.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING

~~MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.~~
~~HARMFUL IF SWALLOWED OR INHALED.~~ CAUSES SUBSTANTIAL, BUT TEMPORARY, EYE INJURY. CAUSES SKIN IRRITATION. CONTACT WITH IRRITATED SKIN, OR A CUT, OR REPEATED CONTACT WITH INTACT SKIN MAY RESULT IN POISONING. Do not get in eyes, on skin or on clothing. ~~Do not breathe spray mist.~~ Avoid breathing vapor or spray mist. Do not feed forage from treated crops to livestock. Keep livestock and pets out of treated fields and crop areas.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants or coveralls over long-sleeved shirt and long pants.
- Waterproof gloves.
- Chemical-resistant footwear plus socks.
- Protective eyewear.
- Chemical-resistant headgear for overhead exposure.
- Chemical-resistant apron when cleaning equipment, mixing or loading.

EXCEPTION: After this product has been diluted with at least 50 gallons of water, applicators for AQUATIC SURFACE APPLICATIONS must, at a minimum, wear (Note - Mixers and Loaders for this application method must still wear the personal protective equipment (PPE) as described in the above section):

- Long-sleeved shirt and long pants.
- Shoes plus socks.
- Waterproof gloves.
- Protective eyewear.

EXCEPTION: At a minimum, applicators for AQUATIC SUBSURFACE APPLICATIONS must wear (Note - Mixers and Loaders for this application method must still wear the personal protective equipment (PPE) as described in the above section):

- Short-sleeved shirt and short pants.
- Waterproof gloves.
- Chemical-resistant footwear plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Mixers, loaders, and applicators using closed systems who meet these requirements may wear: long-sleeved shirt and long pants, protective eyewear, waterproof gloves, shoes plus socks, and a chemical-resistant apron when mixing, loading or cleaning equipment. If handling tasks are performed from inside an enclosed cab or aircraft with enclosed cockpits that meet these requirements may wear: long-sleeved shirt, long pants, shoes and socks for the labeling-specified PPE. All labeling-specified PPE must be immediately available for use in an emergency. All applicable requirements as specified in 40 CFR 170.240(d) (4-6) must be followed.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS (TERRESTRIAL AND AQUATIC USES)

This pesticide is toxic to aquatic invertebrates, wildlife. For Terrestrial Uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. For Aquatic Uses, do not apply directly to water except as specified on this label. Treatment of dense weed areas may result in oxygen loss from decomposition of dead weeds. This loss of oxygen may cause fish suffocation. Therefore, treat only 1/3 to 1/2 of the water body area at one time, especially if dense areas of weeds and/or algae exist, and wait 14 days between treatments.

Necessary approval and/or Permits should be obtained prior to application if required. Consult the responsible State Agencies (i.e., Fish and Game agencies or Department of Natural Resources) before making applications to public waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product through any type of irrigation system.

DIRECTIONS

REWARD® Landscape and Aquatic Herbicide is a nonvolatile herbicidal chemical for use as a general herbicide to control weeds in noncrop and aquatic areas. Absorption and herbicidal action is usually quite rapid with effects visible in a few days. REWARD Landscape and Aquatic Herbicide controls weeds by interfering with photosynthesis within green plant tissue. Weed plants should be succulent and actively growing for best results. Rinse all spray equipment thoroughly with water after use. AVOID SPRAY DRIFT to crops, ornamentals, and other desirable plants during application as injury may result. Application to muddy water may result in reduced control. Minimize creating muddy water during application. Use of dirty or muddy water for REWARD Landscape and Aquatic Herbicide dilution may result in reduced herbicidal activity. Avoid applying under conditions of high wind, water flow or wave action.

SPRAY DRIFT MANAGEMENT

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

- The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature inversions).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces large droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops).

AGRICULTURAL USES:

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants, or coveralls over long-sleeved shirt and long pants.
 - Waterproof gloves.
 - Chemical-resistant footwear plus socks.
 - Protective eyewear.
 - Chemical-resistant headgear for overhead exposure.
-

COMMERCIAL GREENHOUSES AND NURSERIES: For general weed control in commercial greenhouses (beneath benches), (field grown and container stock), and other similar areas. REWARD Landscape and Aquatic Herbicide may be applied preplant or postplant preemergence in field grown ornamental nursery plantings or postemergence as a directed spray. REWARD Landscape and Aquatic Herbicide may also be applied preemergence in ornamental seed crops. Avoid contact with desirable foliage as injury may occur. Do not use on food or feed crops.

Spot spray: 1 to 2 qts. REWARD Landscape and Aquatic Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 100 gallons water or 0.75 oz. (22 mls) REWARD Landscape and Aquatic Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon water.

Broadcast: 1 to 2 pints REWARD Landscape and Aquatic Herbicide in a minimum of 15 gallons water per acre. Add the labeled rate of a 75% or greater nonionic surfactant per 100 gallons spray mixture. Use an adequate spray volume to insure good coverage.

ORNAMENTAL SEED CROPS (FLOWERS, BULBS, ETC.) For preharvest desiccation of ornamental seed crops. **NOT FOR FOOD OR FIBER CROPS.** **Broadcast (Air or Ground):** 1.5 to 2 pints REWARD Landscape and Aquatic Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per acre in sufficient water (minimum of 5 gallons by air; 15 gallons by ground) for desiccation and weed burndown. Repeat as needed at no less than 5-day intervals up to three applications. Do not use seed, screenings, or waste as feed or for consumption.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas or vicinity where there may be drift.

For terrestrial uses, do not enter or allow entry of maintenance workers into treated areas, or allow contact with treated vegetation wet with spray, dew or rain, without appropriate protective clothing until spray has dried.

For aquatic uses, do not enter treated areas while treatments are in progress.

DIRECTIONS FOR LANDSCAPE, INDUSTRIAL, RECREATIONAL, COMMERCIAL, RESIDENTIAL AND PUBLIC AREAS:

Reward provides fast control of broadleaf and grassy weeds in industrial, recreational, golf course, commercial, residential, and public areas.

Reward is a non-selective herbicide that rapidly kills undesirable aboveground weed growth in 24 to 36 hours. Avoid application of Reward to desirable plants.

Reward is a contact/desiccant herbicide, it is essential to obtain complete coverage of the target weeds to get good control. Improper application technique and/or application to stressed weeds may result in unacceptable weed control. For best results apply to actively growing, young weeds.

Difficult weeds (such as perennial or deeply rooted weeds) can often be controlled by tank mixing Reward with other systemic-type herbicides. Refer to other product labels for specific application directions.

For residual weed control, tank mix Reward with a pre-emergent herbicide labeled for the intended use site. When mixing Reward with another herbicide it is recommended to mix just a small amount first to determine if the mixture is physically compatible before proceeding with larger volumes.

Zeneca has not tested all possible tank mixtures with other herbicides for compatibility, efficacy or other adverse effects. Before mixing with other herbicides Zeneca recommends you first consult your state experimental station, state university or extension agent.

Grounds maintenance weed control:

Reward can be used as a spot or broadcast spray to control weeds in public, commercial and residential landscapes, including landscape beds, lawns, golf courses and roadsides. Reward can also be used for weed control around the edges and nonflooded portions of ponds, lakes and ditches.

Trim and Edge weed control:

Reward can be used to eliminate undesired grass and broadleaf plant growth in a narrow band along driveways, walkways, patios, cart paths, fence lines, and around trees, ornamental gardens, buildings, other structures and beneath noncommercial greenhouse benches. Vegetation control with Reward is limited to the spray application width. Do not exceed the labeled rate of Reward as excessive rates may result in staining of concrete-based materials.

Reward, since it does not translocate systemically, can be used as an edging or pruning tool when precisely applied to select areas of grass or to undesirable growth on desirable ornamental bedding plants, ground covers, etc.

Industrial weed control:

Reward can be used as a spot or broadcast spray either alone or in combination with other herbicides as a fast burn-down or control weeds in rights-of-ways, railroads beds/yards, highways, roads, dividers and medians, parking lots, pipelines, pumping stations, public utility lines, transformer stations and substations, electric utilities, storage yards and other non-crop areas.

Spot spray: 1 to 2 qts. REWARD Landscape and Aquatic Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 100 gallons water or 0.75 oz. (22 mls) REWARD Landscape and Aquatic Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon water.

Broadcast: 1 to 2 pints REWARD Landscape and Aquatic Herbicide in a minimum of 15 gallons water per acre. Add the labeled rate of 75% or greater nonionic surfactant per 100 gallons spray mixture. Use an adequate spray volume to insure good coverage. Greater water volumes are necessary if the target plants are tall and/or dense. It is recommended that 60 gallons or greater water volume be used to obtain good coverage of dense weeds.

14/32

TURF RENOVATION (All Turf Areas Except Commercial Sod Farms)

To desiccate golf course turf and other turf areas prior to renovation, apply 1 to 2 pints of REWARD Landscape and Aquatic Herbicide per acre plus the labeled rate of a 75% or greater nonionic surfactant in 20 to 100 gallons of water (4 teaspoons of REWARD Landscape and Aquatic Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon of water) using ground spray equipment. Apply for full coverage and thorough contact with the turfgrass. Apply only when the turf is dry, free from dew and incidental moisture. For enhanced turf desiccation, especially in the case of thick turfgrass, water volumes should approach 100 gallons of water per acre.

For **suppression** of regrowth and quick desiccation of treated turfgrass, REWARD Landscape and Aquatic Herbicide may be mixed with other systemic non-selective or systemic postemergence grassy weed herbicides. Refer to other product labels for specific application directions and restrictions.

Avoid spray contact with, or spray drift to, foliage of ornamental plants or food crops.

Do not graze livestock on treated turf or feed treated thatch to livestock.

DORMANT ESTABLISHED TURFGRASS (Bermudagrass, zoysiagrass) NONFOOD OR FEED CROP

For control of emerged annual broadleaf and grass weeds, including Little Barley*, Annual Bluegrass, Bromes including Rescuegrass, Sixweeks fescue, Henbit, Buttercup, and Carolina Geranium in established dormant bermudagrass lawns, parks, golf courses, etc.

Apply 1 to 2 pts. REWARD Landscape and Aquatic Herbicide per acre in 20 to 100 gallons spray mix by ground as a broadcast application. Add the labeled rate of a 75% or greater nonionic surfactant per 100 gallons spray mixture.

Bermudagrass must be dormant at application. Application to actively growing bermudagrass may cause delay or permanent injury. Users in the extreme Southern areas should be attentive to the extent of dormancy at the time of application.

*For control of Little Barley, apply REWARD Landscape and Aquatic Herbicide prior to the mid-boot stage.

AQUATIC USE DIRECTIONS

New York - Not for Sale or Use in New York State without Supplemental Special Local Needs Labeling.

Necessary approval and/or permits should be obtained prior to application if required. Consult the responsible State Agencies (i.e., Fish and Game agencies or Department of Natural Resources). Treatment of dense weed areas may result in oxygen loss from decomposition of dead weeds. This loss of oxygen may cause fish suffocation. Therefore, treat only 1/3 to 1/2 of the water body area at one time and wait 14 days between treatment.

For application only to **still water** (i.e. ponds, lakes, and drainage ditches) where there is minimal or no outflow to public waters.

and/or

For applications to **public waters** in ponds, lakes, reservoirs, marshes, bayous, drainage ditches, canals, streams, rivers and other slow moving or quiescent bodies of water for control of aquatic weeds. For use by:

- Corps of Engineers; or
- Federal or State Public Agencies (i.e., Water Management District personnel, municipal officials); or
- Applicators and/or Licensees (Certified for aquatic pest control) that are authorized by the State or Local government .

Treated water may be used according to the following table or until such time as an approved assay (example: PAM II Spectromatic Method) shows that the water does not contain more than the designated maximum contaminant level goal (MCLG) of 0.02 mg/l. (ppm) of diquat dibromide (calculated as the cation):

16/32

**WATER USE RESTRICTIONS FOLLOWING APPLICATIONS
WITH REWARD Landscape and Aquatic Herbicide (Days)**

Application Rate	Drinking	Fishing and Swimming	Livestock Consumption	Spray Tank Applications** and Irrigation to Turf and Ornamentals	Spray Tank Applications** and Irrigation to Food Crops
2 gal/surface acre	3 days	0	1 day	3 days	5 days
1 gal/surface acre	2 days	0	1 day	2 days	5 days
0.75 gal/surface acre	2 days	0	1 day	2 days	5 days
0.50 gal/surface acre	1 day	0	1 day	1 day	5 days
Spot Spray* (< 0.5 gal/surface acre)	1 day	0	1 day	1 day	5 days

* Rates refer to total surface area.

**For preparing agricultural sprays for food crops, turf or ornamentals (to prevent phytotoxicity), do not use water treated with REWARD before the specified time period.

When the contents of more than one spray tank is necessary to complete a single aquatic application, no water holding restrictions apply between the consecutive spray tanks.

No applications are to be made in areas where commercial processing of fish, resulting in the production of fish protein concentrate or fish meal, is practiced. Before application, coordination and approval of local and/or State authorities must be obtained.

17/32

Apply REWARD Landscape and Aquatic Herbicide in accordance with the following table:

WEED SPECIES	SUBSURFACE OR BOTTOM PLACEMENT GALS/SURFACE ACRE*	SURFACE GALS/SURFACE ACRE*
Bladderwort (<i>Utricularia</i> spp.)	1-2	2
Coontail (<i>Ceratophyllum demersum</i>)	2	2
Elodea (<i>Elodea</i> spp.)	2	2
Naiad (<i>Najas</i> spp.)	1-2	2
Pondweeds ¹ (<i>Potamogeton</i> spp.)	2	2
Watermilfoils (<i>Myriophyllum</i> spp.)	1-2	2
Hydrilla (<i>Hydrilla verticillata</i>)	2	- 2
Waterlettuce ² (<i>Pistia Stratiotes</i>)	NA	0.5 - 0.75
Waterhyacinth ² (<i>Eichhornia crassipes</i>)	NA	0.5 - 0.75
Pennywort ³ (<i>Hydrocotyle</i> spp.)	NA	0.5 - 0.75
Frog's Bit ³ (<i>Limnobium spongia</i>)	NA	0.5 - 0.75
Salvinia ² (<i>Salvinia</i> spp.)	NA	0.5 - 0.75
Duckweed ⁴ (<i>Lemna</i> spp.)	NA	1
Cattails ³ (<i>Typha</i> spp.)	NA	1-2
Algae ⁵ (<i>Spirogyra</i> spp. & <i>Pithophora</i> spp.)	1-2	2

* For water less than or equal to 2 feet in average depth of treatment area, use a maximum of 1 gallon REWARD Landscape and Aquatic Herbicide per surface acre. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment area, for example, shallow shoreline areas. At water temperatures below 50°-60°F efficacy and immediacy of results may be reduced.

- ¹ REWARD Landscape and Aquatic Herbicide controls *Potamogeton* species except Richardson's pondweed (*P. richardsonii*). For control of *P. robbinsii*, applications must be made when the plants are in the early stages of growth such as in Spring and early Summer.
- ² For salvinia, waterlettuce and water hyacinth, use the labeled rate of REWARD Landscape and Aquatic Herbicide in 75 to 200 gallons water plus the labeled rate of a 75% or greater nonionic surfactant per acre for surface sprays and for aerial application for waterlettuce and water hyacinth control, apply the labeled rate of REWARD Landscape and Aquatic Herbicide in 10 to 24 gallons water plus the labeled rate of a 75% or greater nonionic surfactant per acre.
- ³ For Pennywort and cattail control, apply in 50-150 gallons of water plus the labeled rate of a 75% or greater nonionic surfactant per acre for full coverage and thorough weed contact. Repeat treatments may be necessary to control regrowth. For best results, apply before flowering (cattail).
- ⁴ For duckweed control, apply as an overall spray in 50-150 gallons of water plus the labeled rate of a 75% or greater nonionic surfactant per acre. Retreatment may be necessary for plants missed in previous applications and regrowth.
- ⁵ For suppression of certain filamentous algae species including *Spirogyra* and *Pithophora*, apply according to the submersed use directions.
- ⁶ Not for use in California.

APPLICATION: In mixed weed populations, use the high rate of application as indicated by weeds present.

SUBSURFACE APPLICATIONS: Where the submersed weed growth, especially Hydrilla, has reached the water surface, apply either in a water carrier or an invert emulsion through boom trailing hoses carrying nozzle tips to apply the dilute spray below the water surface to insure adequate coverage.

BOTTOM PLACEMENT: Where the submersed weeds, especially Hydrilla, Bladderwort, and Coontail growth have reached the water surface or where water is slowly moving through the submersed weed growth that has reached the water surface, especially Hydrilla, Bladderwort, and Coontail, control may be enhanced when applied in an invert emulsion carrier injecting diluted REWARD Landscape and Aquatic Herbicide near the bottom with weighted hoses. The addition of a copper-based algaecide will improve control. Where algae are present along with the submersed weeds, pre-treatment with copper-based algaecide at recommended rates is advised for best results.

SURFACE APPLICATION: For submerged aquatic weeds, apply REWARD Landscape and Aquatic Herbicide either as concentrate slowly poured directly from the container in strips or as a spray in sufficient carrier. Applications should be made to ensure complete coverage of the weed areas. In mixed weed populations, use the high rate of application as indicated by weeds present.

IF POSTING IS REQUIRED BY YOUR STATE OR TRIBE -- CONSULT THE AGENCY RESPONSIBLE FOR PESTICIDE REGULATIONS FOR SPECIFIC DETAILS.

GENERAL RECOMMENDATIONS FOR "POSTING NOTIFICATION" *

- **Flowing water:** "post" the restricted area (within/at 1600 feet downstream of treatment) for the duration of the water use restriction.
- **Standing water:** "post" the restricted area (within/at 1/4 mile of treatment) for the duration of the water use restriction.
- **No "posting" is necessary where water use is greater than 1600 feet downstream of treated water in flowing water bodies or where water use is greater than 1/4 mile from treated water in standing water bodies.**

*"Posting" should be removed at the end of the restriction period.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Do not contaminate feed, foodstuffs or drinking water. Do not store or transport near feed or food. Store at temperature above 32°F. For help with any spill, leak, fire or exposure involving this material, call CHEMTREC (1-800-424-9300).

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

~~CONTAINER DISPOSAL: Do not reuse container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse (or equivalent). Do not reuse container. Incinerate, burn, or puncture and dispose of in a sanitary landfill, or dispose of by other procedures allowed by State and local authorities. If burned, stay out of smoke.~~

FOR BULK AND MINI-BULK CONTAINERS:

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.

REFILL ONLY WITH REWARD Landscape and Aquatic Herbicide. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than REWARD Landscape and Aquatic Herbicide will result in contamination and may weaken container.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER!

20/32

REWARD® LS
Landscape Herbicide

Not for sale to, use and storage by individual homeowners unless licensed by the state.

A non-selective herbicide for fast control of broadleaf and grassy weeds in landscape around recreational, industrial, commercial, residential and public areas; and in commercial nurseries and greenhouses.

ACTIVE INGREDIENT:

Diquat dibromide [6,7-dihydrodipyrido (1,2-a:2',1'-c) pyrazinedium dibromide]	37.3%
INERT INGREDIENTS	<u>62.7%</u>
TOTAL	100.0%

Contains 2 lbs. diquat cation per gal. as 3.73 lbs. salt per gal.

EPA Reg. No. 10182-404
EPA Est. No.

Net Contents:

TO PREVENT ACCIDENTAL POISONING, NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS AND USE STRICTLY IN ACCORDANCE WITH ENTIRE LABEL

DO NOT USE THIS PRODUCT FOR REFORMULATION

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

FEB 18 2000

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
10182-404**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See (inside, back or side panels) for
Statement of Practical Treatment and Precautionary Statements

Made in U.S.A.
ZENECA Professional Products
ZENECA Ag Products Inc.
Wilmington, DE 19850-5458

COMPLETE DIRECTIONS FOR USE
CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

22/32

TABLE OF CONTENTS

	Page
STATEMENT OF PRACTICAL TREATMENT	
PRECAUTIONARY STATEMENTS	
HAZARDS TO HUMANS AND DOMESTIC ANIMALS	
PERSONAL PROTECTIVE EQUIPMENT	
ENVIRONMENTAL HAZARDS	
DIRECTIONS FOR USE	
DIRECTIONS	
AGRICULTURAL USE REQUIREMENTS	
NON-AGRICULTURAL USE REQUIREMENTS	
STORAGE AND DISPOSAL	

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: IMMEDIATELY give water or milk to drink and induce vomiting by inserting finger in throat. Do not induce vomiting or give anything by mouth to an unconscious person. Take person and product container to the nearest hospital or physician fast. **PROMPT TREATMENT IS ESSENTIAL TO COUNTERACT POISONING** and should be initiated before signs and symptoms of injury appear.

IF ON SKIN: IMMEDIATELY wash with plenty of soap and water. See a doctor if diquat contacts a skin cut, abrasion or area of irritation.

IF IN EYES: IMMEDIATELY ~~wash eyes with water for at least~~ hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. ~~and g~~ Get medical attention.

IF INHALED: IMMEDIATELY remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention. ~~get away from spray mist. Stop and check spray procedure. See a doctor if irritation persists.~~

NOTE TO PHYSICIANS: Call ZENECA Medical Emergency Information Network 1-800-F-A-S-T-M-E-D (327-8633) at any hour to obtain toxicology information and a diquat analysis. To be effective, treatment for diquat poisoning must begin **IMMEDIATELY**. Treatment consists of binding diquat in the gut with suspensions of activated charcoal or bentonite clay, administration of cathartics to enhance elimination, and removal of diquat from the blood by charcoal hemoperfusion or continuous hemodialysis.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL 1-800-F-A-S-T-M-E-D (327-8633)

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING

MAY BE FATAL IF ~~SWALLOWED, INHALED OR~~ ABSORBED THROUGH THE SKIN. HARMFUL IF SWALLOWED OR INHALED. CAUSES SUBSTANTIAL, BUT TEMPORARY, EYE INJURY. CAUSES SKIN IRRITATION. CONTACT WITH IRRITATED SKIN, OR A CUT, OR REPEATED CONTACT WITH INTACT SKIN MAY RESULT IN POISONING. Do not get in eyes, on skin or on clothing. ~~Do not breathe spray mist.~~ Avoid breathing vapor or spray mist. Do not feed forage from treated crops to livestock. Keep livestock and pets out of treated fields and crop areas.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants or coveralls over long-sleeved shirt and long pants.
- Waterproof gloves.
- Chemical-resistant footwear plus socks.
- Protective eyewear.
- Chemical-resistant headgear for overhead exposure.
- Chemical-resistant apron when cleaning equipment, mixing or loading.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Mixers, loaders, and applicators using closed systems who meet these requirements may wear: long-sleeved shirt and long pants, protective eyewear, waterproof gloves, shoes plus socks, and a chemical-resistant apron when mixing, loading or cleaning equipment. If handling tasks are performed from inside an enclosed cab or aircraft with enclosed cockpits that meet these requirements may wear: long-sleeved shirt, long pants, shoes and socks for the labeling-specified PPE. All labeling-specified PPE must be immediately available for use in an emergency. All applicable requirements as specified in 40 CFR 170.240(d) (4-6) must be followed.


User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates, wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

DIRECTIONS

REWARD® LS Landscape Herbicide is a nonvolatile herbicidal chemical for use as a general herbicide to control weeds in noncrop areas. Absorption and herbicidal action is usually quite rapid with effects visible in a few days. REWARD LS Landscape Herbicide controls weeds by interfering with photosynthesis within green plant tissue. Weed plants should be succulent and actively growing for best results. Rinse all spray equipment thoroughly with water after use. AVOID SPRAY DRIFT to crops, ornamentals, and other desirable plants during application as injury may result. Use of dirty or muddy water for REWARD LS Landscape Herbicide dilution may result in reduced herbicidal activity.

Do not apply this product through any type of irrigation system.

SPRAY DRIFT MANAGEMENT


AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

- The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.



DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature inversions).

Controlling Droplet Size

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces large droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

27/32

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops).

AGRICULTURAL USES:

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.


PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants, or coveralls over long-sleeved shirt and long pants.
 - Waterproof gloves.
 - Chemical-resistant footwear plus socks.
 - Protective eyewear.
 - Chemical-resistant headgear for overhead exposure.
-

COMMERCIAL GREENHOUSES AND NURSERIES: For general weed control in commercial greenhouses (beneath benches), (field grown and container stock), and other similar areas. REWARD LS Landscape Herbicide may be applied preplant or postplant preemergence in field grown ornamental nursery plantings or postemergence as a directed spray. REWARD LS Landscape Herbicide may also be applied preemergence in ornamental seed crops. Avoid contact with desirable foliage as injury may occur. Do not use on food or feed crops.

Spot spray: 1 to 2 qts. REWARD LS Landscape Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 100 gallons water or 0.75 oz. (22 mls) REWARD LS Landscape Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon water.

Broadcast: 1 to 2 pints REWARD LS Landscape Herbicide in a minimum of 15 gallons water per acre. Add the labeled rate of a 75% or greater nonionic surfactant per 100, gallons spray mixture. Use an adequate spray volume to insure good coverage.



ORNAMENTAL SEED CROPS (FLOWERS, BULBS, ETC.) For preharvest desiccation of ornamental seed crops. **NOT FOR FOOD OR FIBER CROPS.** **Broadcast (Air or Ground):** 1.5 to 2 pints REWARD LS Landscape Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per acre in sufficient water (minimum of 5 gallons by air; 15 gallons by ground) for desiccation and weed burndown. Repeat as needed at no less than 5-day intervals up to three applications. Do not use seed, screenings, or waste as feed or for consumption.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas or vicinity where there may be drift.

Do not enter or allow entry of maintenance workers into treated areas, or allow contact with treated vegetation wet with spray, dew or rain, without appropriate protective clothing until spray has dried.

DIRECTIONS FOR LANDSCAPE, INDUSTRIAL, RECREATIONAL, COMMERCIAL, RESIDENTIAL AND PUBLIC AREAS:

Reward LS provides fast control of broadleaf and grassy weeds in industrial, recreational, golf course, commercial, residential, and public areas.

Reward LS is a non-selective herbicide that rapidly kills undesirable aboveground weed growth in 24 to 36 hours. Avoid application of Reward to desirable plants.

Reward LS is a contact/desiccant herbicide, it is essential to obtain complete coverage of the target weeds to get good control. Improper application technique and/or application to stressed weeds may result in unacceptable weed control. For best results apply to actively growing, young weeds.

Difficult weeds (such as perennial or deeply rooted weeds) can often be controlled by tank mixing Reward LS with other systemic-type herbicides. Refer to other product labels for specific application directions.

For residual weed control, tank mix Reward LS with a pre-emergent herbicide labeled for the intended use site. When mixing Reward LS with another herbicide it is recommended to mix just a small amount first to determine if the mixture is physically compatible before proceeding with larger volumes.

Zeneca has not tested all possible tank mixtures with other herbicides for compatibility, efficacy or other adverse effects. Before mixing with other herbicides Zeneca recommends you first consult your state experimental station, state university or extension agent.

Grounds maintenance weed control:

Reward LS can be used as a spot or broadcast spray to control weeds in public, commercial and residential landscapes, including landscape beds, lawns, golf courses and roadsides. Reward LS can also be used for weed control around the edges and nonflooded portions of ponds, lakes and ditches.

Trim and Edge weed control:

Reward LS can be used to eliminate undesired grass and broadleaf plant growth in a narrow band along driveways, walkways, patios, cart paths, fence lines, and around trees, ornamental gardens, buildings, other structures and beneath noncommercial greenhouse benches. Vegetation control with Reward LS is limited to the spray application width. Do not exceed the labeled rate of Reward LS as excessive rates may result in staining of concrete-based materials.

Reward LS, since it does not translocate systemically, can be used as an edging or pruning tool when precisely applied to select areas of grass or to undesirable growth on desirable ornamental bedding plants, ground covers, etc.

Industrial weed control:

Reward LS can be used as a spot or broadcast spray either alone or in combination with other herbicides as a fast burn-down or control weeds in rights-of-ways, railroads beds/yards, highways, roads, dividers and medians, parking lots, pipelines, pumping stations, public utility lines, transformer stations and substations, electric utilities, storage yards and other non-crop areas.

Spot spray: 1 to 2 qts. REWARD LS Landscape Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 100 gallons water or 0.75 oz. (22 mls) REWARD LS Landscape Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon water.

Broadcast: 1 to 2 pints REWARD LS Landscape Herbicide in a minimum of 15 gallons water per acre. Add the labeled rate of 75% or greater nonionic surfactant per 100 gallons spray mixture. Use an adequate spray volume to insure good coverage. Greater water volumes are necessary if the target plants are tall and/or dense. It is recommended that 60 gallons or greater water volume be used to obtain good coverage of dense weeds.

TURF RENOVATION (All Turf Areas Except Commercial Sod Farms)

To desiccate golf course turf and other turf areas prior to renovation, apply 1 to 2 pints of REWARD LS Landscape Herbicide per acre plus the labeled rate of a 75% or greater nonionic surfactant in 20 to 100 gallons of water (4 teaspoons of REWARD LS Landscape Herbicide plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon of water) using ground spray equipment. Apply for full coverage and thorough contact with the turf grass. Apply only when the turf is dry, free from dew and incidental moisture. For enhanced turf desiccation, especially in the case of thick turfgrass, water volumes should approach 100 gallons of water per acre.

For **suppression** of regrowth and quick desiccation of treated turfgrass, REWARD LS Landscape Herbicide may be mixed with other systemic non-selective or systemic postemergence grassy weed herbicides. Refer to other product labels for specific application directions and restrictions.

Avoid spray contact with, or spray drift to, foliage of ornamental plants or food crops.

Do not graze livestock on treated turf or feed treated thatch to livestock.


DORMANT ESTABLISHED TURFGRASS (Bermudagrass, zoysiagrass) NONFOOD OR FEED CROP

For control of emerged annual broadleaf and grass weeds, including Little Barley*, Annual Bluegrass, Bromes including Rescuegrass, Sixweeks fescue, Henbit, Buttercup, and Carolina Geranium in established dormant bermudagrass lawns, parks, golf courses, etc.

Apply 1 to 2 pts. REWARD LS Landscape Herbicide per acre in 20 to 100 gallons spray mix by ground as a broadcast application. Add the labeled rate of a 75% or greater nonionic surfactant per 100 gallons spray mixture.

Bermudagrass must be dormant at application. Application to actively growing bermudagrass may cause delay or permanent injury. Users in the extreme Southern areas should be attentive to the extent of dormancy at the time of application.

*For control of Little Barley, apply REWARD LS Landscape Herbicide prior to the mid-boot stage.



STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Do not contaminate feed, foodstuffs or drinking water. Do not store or transport near feed or food. Store at temperature above 32°F. For help with any spill, leak, fire or exposure involving this material, call CHEMTREC (1-800-424-9300).

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. ~~Triple rinse (or equivalent). Do not reuse container. Incinerate, burn, or puncture and dispose of in a sanitary landfill, or dispose of by other procedures allowed by State and local authorities. If burned, stay out of smoke.~~

REWARD® LS is a trademark of a ZENECA Group company