

10182-450

06/01/2000

1/15



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
10182-450

Date of Issuance:  
JUN 1 2000

NOTICE OF PESTICIDE:  
  x   Registration  
       Reregistration  
  
(under FIFRA, as amended)

Term of Issuance:  
Conditional

Name of Pesticide Product:  
Devrinol Lawn and Ornamental Selective Herbicide

Name and Address of Registrant (include ZIP Code):

Zeneca Ag Products  
1800 Concord Pike  
P.O. Box 15458  
Wilmington, DE 19850-5458

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following label changes listed below before you release the product for shipment:
  - a. Add the phrase, "EPA Reg. No. 10182-450".
  - b. Add the common chemical name, Napropamide, to the ingredient declaration.

Signature of Approving Official:

Date:

6-1-00


page 2

EPA Reg. No. 10182-450

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



James A. Tompkins  
Product Manager (25)  
Herbicide Branch  
Registration Division (7505C)

Enclosure

3/15

**NEW! READY TO USE  
NON-YELLOW GRANULE**

**DEVIRINOL® 2-G LAWN AND ORNAMENTAL  
Selective Herbicide  
FOR HOMEOWNER USE**

~~— A Preemergence Herbicide for Weed Control in Home Flower Gardens and Landscape  
— Plantings. Use in Deciduous and Evergreen Trees and Shrubs, Ground Covers,  
— Perennial Flowers, Fruit and Nut Crops, Container Grown Plants, and Turfgrass~~

~~One This product prevents weeds in flower beds, rose gardens, ground covers, (including  
dichondra), potted and landscape ornamentals, fruit trees and established turfgrasses:  
(bahagrass, bermudagrass, centipedegrass, St. Augustinegrass and fescue).~~

~~Single application prevents both annual grass and broadleaf weeds. Prevents crabgrass,  
goosegrass and annual bluegrass germination.~~

~~Provides soil residual control. Covers up to: 1,000 sq. ft. (2 lb. 4 oz. Container)  
5,000 sq. ft. (11 lb. 4 oz. Container)~~

**ACTIVE INGREDIENT:**

*N,N*-diethyl-2-(1-naphthalenyloxy)-propionamide ..... 2.0%

**INERT INGREDIENTS:** ..... 98.0%

**TOTAL** ..... 100.0%

Contains 2 Pounds of Active Ingredient per 100 Pounds

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

**JUN 1 2000**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide**

**KEEP OUT OF REACH OF CHILDREN**

EPA Reg. No.: 10182-  
EPA Est. No.:

Net Contents:

**CAUTION**

**SEE SIDE PANEL FOR STATEMENT OF PRACTICAL TREATMENT FIRST AID  
AND PRECAUTIONARY STATEMENTS**

Active Ingredient Made in UK, Formulated and Packaged in USA  
ZENECA Ag Products  
ZENECA Ag Products Inc.  
Wilmington, DE 19850

4/15

**STATEMENT OF PRACTICAL TREATMENT FIRST AID**

If an adverse reaction occurs after exposure, promptly initiate the recommended procedures below. If further treatment is required, seek medical attention.

**IN CASE OF CONTACT:** Flush eyes and skin with plenty of water for several minutes. Get medical attention if eye or skin irritation occurs.

**IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF SWALLOWED:** Immediately give several glasses of water and induce vomiting by placing a finger on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

**IF INHALED:** Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

**FIRST AID**

<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a Poison Control Center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a Poison Control Center or doctor.</li> </ul>
<b>If In Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a Poison Control Center or doctor for treatment advice.</li> </ul>
<b>If On Skin:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a Poison Control Center or doctor for treatment advice.</li> </ul>
<b>If Inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a Poison Control Center or doctor for further treatment advice.</li> </ul>

5/15

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

---

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL 1-800-F-A-S-T-M-E-D  
(1-800-327-8633)

---

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

##### CAUTION

MAY BE HARMFUL IF SWALLOWED. Do not breathe dust. Avoid contact with skin, eyes and clothing. Avoid contamination with food and feed.

##### PERSONAL PROTECTIVE EQUIPMENT

~~Applicators and other handlers must wear:~~

- ~~• Long-sleeved shirt and long pants.~~
- ~~• Shoes plus socks.~~

~~Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.~~

~~When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.~~

---

##### User Safety Recommendations

~~Users should:~~

- ~~• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.~~
  - ~~• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.~~
  - ~~• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.~~
-

6/15

## ENVIRONMENTAL HAZARDS

Do not apply directly to water., or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

## GENERAL PRECAUTIONS

- Read all instructions before using.
- Use only for recommended purposes and at recommended rates.
- Do not overdose.

**NOTICE TO BUYER:** Buyer assumes all responsibility for safety and use not in accordance with directions.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product as specified on this label. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses on lawns and grounds.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

---

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

**Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

7/15

~~PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:~~

- ~~● Coveralls~~
- ~~● Waterproof gloves~~
- ~~● Shoes plus socks.~~

---

### STORAGE AND DISPOSAL

~~Do not contaminate water, food or feed by storage or disposal.~~

**STORAGE:** Keep container closed when not in use. Store in a cool dry place. Do not store near seeds, fertilizer or food stuffs.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** ~~Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. Do not reuse empty container. Securely wrap in newspaper and discard in trash.~~

Formulated with natural clay granule for deep penetration through thick ground covers like Ivy, pachysandra and dichondra.

**HOW DEVIRINOL WORKS:** Devrinol controls many annual grass and broadleaf weeds as they germinate. It does not control established weeds. One application properly applied at the recommended use rate will prevent the germination of most weeds up to 90 days, preventing the need for hand weeding.

**RECOMMENDED APPLICATIONS:** Apply Devrinol over small-established ornamental plants or around larger plants in flowerbeds, rose gardens, landscape ornamental environments, ground covers (including dichondra), lawns, or to potted ornamentals grown for esthetic purpose. Do not use where plants are seeded.

Use Devrinol for weed control in the following plants:

8/15

**WHERE TO USE THIS PRODUCT**

~~DEVIRINOL 2-G ORNAMENTAL can be applied over small ornamental plants or around larger plants in rose gardens, container plantings, ornamental landscapes and ground covers including dichondra.~~

~~Use DEVIRINOL 2-G ORNAMENTAL for weed control in the following plants:~~

**Deciduous and Evergreen Trees**

Acer	Eucalyptus	Picea
Ash	Ficus	Pin Oak
Birch	Fir	Pinus
Black Walnut	Gleditsia	Pittosporum
Cedar	Hemlock	Podocarpus
Citrus	Japanese Larch	Populus
Crataegus	Laburnum	Prunus
Cypress	Malus	Pears
Dogwood	Olive (Olea)	Pistachios
Douglas Fir	Palm (Washingtonia robusta)	

**Deciduous and Evergreen Shrubs**

Abelia	Forsythia	Oleander
Asparagus (Sprengeri)	Gardenia	Osmanthus
Aucuba	Heather (Calluna)	Photinia
Azalea	Hibiscus	Pittosporum
Bottlebrush	Honeysuckle (Lonicera)	Podocarpus
Bougainvillea	Hypericum	Pyracantha
Buxus	Ilex	Raphiolepis
Camellia	Juniperus	Rhododendron
Cotoneaster	Leucothe	Rosa
Crepe Myrtle	Ligustrum	Star Jasmine
Dogwood	Nandina	Taxus
Euonymus	Natal Plum	Viburnum
		Xylosema

**Fruit and Nuts**

Almond	Lemon	Pistachio
Apple	Nectarine	Plum
Apricot	Orange	Prune
Cherry	Peach	Tangerine
Filbert	Pear	Walnut
Grape	Pecan	

~~Strawberry: (Applications should be delayed until the desired number of daughter plants have become established. Do not apply from bloom through harvest.)~~



9/15

**Ground Covers**

Ajuga  
Carpobrotus  
Delosperma  
Dichondra  
Erysimum  
Ganzania

Hedera  
Hypericum  
Iceplant  
Ivy  
Lantana

Liriope  
Osteosperma  
Pachysandra  
Sedum  
Vinca

**Container Stock**

Abelia  
Agapanthus  
Ageratum  
Asparagus  
Azalea  
Bottlebrush  
Bougainvillea  
Buxus  
Camellia  
Cedar  
Cotoneaster  
Crepe Myrtle  
Cypress  
Dogwood  
Douglas Fir

Eucalyptus  
Euonymus  
Fir  
Forsythia  
Gardenia  
Geranium  
Heather  
Hedera  
Hibiscus  
Ilex  
Juniperus  
Leucothe  
Ligustrum  
Liriope  
Nandina

Pachysandra  
Palm (Washingtonia  
robusta)  
Photinia  
Pinus spp.  
Pittosporum  
Podocarpus  
Pyracantha  
Raphiolepis  
Rhododendron  
Rosa  
Sedum  
Star Jasmine  
Thuja  
Vinca  
Xylosema

**Flowers (Established)**

African Daisy  
Aster  
Bird of Paradise

Chrysanthemum  
Dahlia  
Daisy  
Geranium  
Gladiolus

Narcissus  
Petunia  
Plantain Lily (Hosta)  
Zinnia

**Lawns (Warm Season)**

Bahiagrass  
Bermudagrass

Centipedegrass  
Fescue

St. Augustinegrass

**WHAT THIS PRODUCT DOES**

DEVTRINOL 2-G ORNAMENTAL controls certain annual grassy and broadleaf weeds as they germinate. It does not control established weeds. The weeds controlled are listed below:

10/15

WEEDS CONTROLLED BY DEVRINOL 2-G ORNAMENTAL	
<b>ANNUAL GRASSES:</b>	
Barnyardgrass, Junglerice	<i>Echinochloa</i> spp.
Wild Barley	<i>Hordeum leporinum</i>
Annual Bluegrass	<i>Poa annua</i>
Bromes	<i>Bromus</i>
— Cheatgrass	— <i>B. secalinus</i>
— Downy Brome	— <i>B. tectorum</i>
— Ripgut Brome	— <i>B. rigidus</i>
— Soft Chess	— <i>B. mollis</i>
Large (Hairy) Crabgrass	<i>Digitaria sanguinalis</i>
Smooth Crabgrass	<i>Digitaria ischaemum</i>
Southwestern Cupgrass	<i>Eriochloa gracilis</i>
Foxtails	<i>Setaria</i>
— Bristly Foxtail	— <i>S. verticilla</i>
— Giant Foxtail	— <i>S. faberi</i>
— Green Foxtail	— <i>S. viridis</i>
— Yellow Foxtail	— <i>S. lutescens</i>
Goosegrass	<i>Eleusine indica</i>
Seedling Johnsongrass	<i>Sorghum halepense</i>
Wild Oats	<i>Avena fatua</i>
Panicums	<i>Panicum</i>
— Guineagrass	— <i>P. Paximum</i>
— Fall Panicum	— <i>P. dichotomiflorum</i>
— Texas Panicum	— <i>P. texanum</i>
— Witchgrass	— <i>P. capillare</i>
Italian Ryegrass	<i>Lolium multiflorum</i>
Sandbur	<i>Cenchrus</i> spp.
Red Sprangletop	<i>Leptochloa filiformis</i>

11/15

WEEDS CONTROLLED BY DEVRINOL 2-G ORNAMENTAL	
Stinkgrass	<i>Eragrostis ciliaris</i>
<b>ANNUAL BROADLEAF WEEDS:</b>	
Carpetweed	<i>Mollugo verticillata</i>
Common Chickweed	<i>Stellaria media</i>
Coast Fiddleneck	<i>Amsinckia intermedia</i>
Filaree	<i>Erodium spp.</i>
Groundsel	<i>Senecio spp.</i>
Prostrate Knotweed	<i>Polygonum aviculare</i>
Lambsquarters	<i>Chenopodium album</i>
Prickly Lettuce	<i>Lactuca scariola</i>
Little Mallow*	<i>Malva parviflora</i>
Redroot Pigweed	<i>Amaranthus retroflexus</i>
Pineappleweed	<i>Matricaria matricarioides</i>
Common Purslane	<i>Portulaca oleracea</i>
Horse Purslane	<i>Trianthema portulacastrum</i>
Common Ragweed**	<i>Ambrosia artemisiifolia</i>
Annual Sowthistle	<i>Sonchus oleraceus</i>

\*From seed  
 \*\*Suppression

**WEEDS CONTROLLED**

**GRASSES (ANNUAL):** Annual Bluegrass (*Poa Annua*), Barnyardgrass, Bromes (Cheatgrass, Downy, Rippgut and Soft Chess), Crabgrass (Large, Hairy and Smooth), Southwestern Cupgrass, Foxtails (Bristly, Giant, Green and Yellow), Goosegrass, Johnsongrass (Seedling), Panicums (Guineagrass, Fall and Texas), Italian Ryegrass, Sandbur, Red Sprangletop, Stinkgrass and Wild Oats.

**BROADLEAF WEEDS:** Carpetweed, Common Chickweed, Coast Fiddleneck, Filaree, Groundsel, Prostrate Knotweed, Lambsquarters, Prickly Lettuce, Little Mallow (Seedling), Redroot Pigweed, Pineappleweed, Common Purslane, Horse Purslane, Common Ragweed (Suppression) and Annual Sowthistle.

12/15

**LAWNS:** Devrinol selectively controls crabgrass, goosegrass and annual bluegrass in Warm Season Turfgrasses only (bahagrass, bermudagrass, centipedegrass, St. Augustinegrass and fescue).

**CONTROL OF CRABGRASS OR GOOSEGRASS:** Application should be made 2-3 weeks prior to weed emergence. Apply using a shaker canister, flow drop or a cyclone type spreader. A more uniform application can be made by spreading half the required amount over the area to be treated, then applying the remainder at right angles. For optimum weed control in lawns, Devrinol requires a minimum of 1/2 inch of rainfall or irrigation within 2-3 days after application. If irrigation methods are not available, the timing of application should be moved back several weeks to allow for some rainfall between application and weed germination.

**SPLIT APPLICATIONS:** For control of both crabgrass and goosegrass, apply the recommended rate for crabgrass followed by a second application 8-10 weeks apart for the control of goosegrass. Timing should be adjusted to the germination of weeds in your area. Split applications generally give longer and better weed control than one application.

**CONTROL OF ANNUAL BLUEGRASS (*Poa Annua*):** Annual bluegrass is a winter annual in most of the Warm Season turf regions; therefore, application should be made in the Fall before the annual bluegrass germinates.

**RESEEDING:** Do not reseed or overseed within six months after application.

**WHEN TO APPLY:** Devrinol can be applied all year to control germinating weeds. Apply Devrinol before weeds germinate in late winter or early spring, after removing established weeds chemically or by hand, to prevent reinfestation of the listed weeds from seed. Fall and winter applications will provide excellent weed control the following spring and summer. When treating plants in containers, apply after setting and watering the plants. See directions under the "Lawns" section for specific lawn applications.

---

#### WHEN TO USE THIS PRODUCT

~~DEVTRINOL 2-G ORNAMENTAL should be used before weeds germinate. It will not control established weeds:~~

~~Application can be made anytime of the year to weed-free soil. Fall and winter applications will provide excellent weed control the following spring and summer.~~

~~In containers, apply after setting and watering the plants.~~

13/15

~~DEVTRINOL 2-G ORNAMENTAL should be spread evenly on the soil surface using a shaker can or granular herbicide applicator.~~

~~Do not apply when weather conditions favor drift from areas treated.~~

~~When weed infestation is heavy or when broadleaves are the primary weeds, use one pound on 145 square feet (6.9 lbs./1,000 sq. ft.). For annual grassy weeds or when weed infestation is light, use one pound on 218 square feet (4.6 lbs/1,000 sq.ft.).~~

~~DEVTRINOL 2-G ORNAMENTAL should be incorporated evenly into the soil using a hand cultivator to a depth of 1-2 inches or irrigation or natural moisture (rainfall or snow) (1/2 inch within 2-3 days) for optimum results.~~

~~For weed control in dichondra, apply at the time of seeding or to establish dichondra including dichondra sod. Irrigate with a minimum of one inch of water.~~

~~RECOMMENDATIONS: TURF GRASS  
(For Use on Warm Season Grasses Only)~~

~~DEVTRINOL 2-G ORNAMENTAL is recommended for annual bluegrass, crabgrass and goosegrass control in the following established turf grasses: bahia, bermudagrass, centipede, St. Augustine and fescue.~~

~~TIMING OF APPLICATION:~~

~~For control of crabgrass and goosegrass, apply DEVTRINOL 2-G ORNAMENTAL at the recommended rate. Application should be made 2-3 weeks prior to weed emergence. DEVTRINOL 2-G ORNAMENTAL should be applied with a gravity flow drop spreader or a cyclone type spreader. A more uniform application can be made by spreading half the required amount over the area to be treated, then applying the remainder at right angles.~~

~~DEVTRINOL 2-G ORNAMENTAL requires a minimum of one half (1/2) inch of rainfall or irrigation 2-3 days after application for adequate control. If irrigation methods are not available, the timing of the application of DEVTRINOL 2-G ORNAMENTAL should be moved back several weeks to allow for some rainfall between application and weed germination.~~

~~RATES OF APPLICATION:~~

~~For Crabgrass Control: Apply 100 to 150 lbs. of DEVTRINOL 2-G ORNAMENTAL uniformly per acre (2.25-3.38\* lbs per 1,000 sq. ft.). Use the higher rate when applying before January 1 (not applicable north of South Carolina or when weed infestation is expected to be heavy).~~

~~\*Where a rate range is given, use the higher rate when weed infestation is expected to be heavy.~~

14/115

~~For Goosegrass Control: Apply 100 to 150 lbs. of DEVIRINOL 2-G ORNAMENTAL per acre (2.25-3.38 lbs. per 1,000 sq. ft.). Goosegrass usually starts to germinate 6 to 7 weeks after the start of crabgrass germination.~~

~~Split Application for Crabgrass and Goosegrass Control: When both of these weeds are to be controlled, apply 100 lbs. of DEVIRINOL 2-G ORNAMENTAL spread uniformly per acre (2.25 lbs. per 1,000 sq. ft.) for crabgrass and another 100 lbs. of DEVIRINOL 2-G ORNAMENTAL per acre for goosegrass. Apply 8-10 weeks apart. Timing should be adjusted to the germination of weeds in your area. Split applications give longer and better weed control than one application.~~

~~For Annual Bluegrass (Poa Annua) Control: Annual bluegrass is a winter annual in most of the warm season turf regions; therefore, application of DEVIRINOL 2-G ORNAMENTAL should be made in the fall before the annual bluegrass germinates. Apply 100-150 lbs. of DEVIRINOL 2-G ORNAMENTAL per acre (2.25-3.38 lbs. per 1,000 sq. ft.).~~

~~Reseeding: Do not reseed or overseed within six months after application.~~

15/15

APPLICATION	RATES	THIS CONTAINER TREATS	
		2 lb. 4 oz. Container	11 lb. 4 oz. Container
<b>ORNAMENTALS</b> Deciduous and Evergreen Trees, Shrubs, Flowers, Ground Covers and Cont- ainer Grown Ornamentals	1 Ounce* per 10 sq. ft.	360 sq. ft.	1,800 sq. ft.
<b>LAWNS: (Warm Season)</b> For Crabgrass or Goosegrass Control Expected Light Infestation: Heavy Infestation:	2.25 lb/1,000 sq. ft. 3.38 lb/1,000 sq. ft.	1,000 sq. ft. 660 sq. ft.	5,000 sq. ft. 3,300 sq. ft.
Split Application: (For control of Crabgrass and Goosegrass)	2.25 lb/1,000 sq. ft. followed 8-10 weeks later with 2.25 lb/1,000 sq. ft.	1,000 sq. ft.	5,000 sq. ft.
For Annual Bluegrass Control: ( <i>Poa Annuua</i> )	2.25 lb/1,000 sq. ft.	1,000 sq. ft.	5,000 sq. ft.

\*2 rounded Tablespoons = 1 Ounce

Do not use spoons for food after use with this product - keep for garden use only.

**HOW TO APPLY:** Spread evenly on the soil surface using the shaker can or granular herbicide applicator. Do not apply when weather conditions favor drift from intended treatment area. For optimum results, incorporate evenly into the soil using a hand cultivator to a depth of 1-2 inches or by irrigation or natural moisture (rainfall or snow) with ½ inch of water within 2-3 days after application. For weed control in Dichondra, apply at the time of seeding or to established Dichondra including Dichondra sod. Irrigate with a minimum of 1 inch of water.

DEVRINOL® is a trademark of a ZENECA company.