

TABLETED DURACHLOR - Slowly dissolve tablets in water to provide continuous disinfection to control bacteria and algae.

DIRECTIONS

Usually pre-treat your pool water with 1/2 TABLET per 10,000 GALLONS OF POOL WATER. Follow directions on the label for best results.

RECOMMENDED DOSE: For each 15-foot pool, 10 tablets in the skimmer basket. Allow 8 to 12 hours for maximum disinfection. The effect will last from 2 to 4 days depending on water temperature and flow rate. The appropriate chlorine residual should be maintained between 1.0 and 1.5 ppm as determined by a suitable test kit. If daily testing indicates the chlorine level has fallen below the minimum level, place 10 tablets in the skimmer basket. If daily testing indicates the chlorine level is high, reduce the number of tablets accordingly.

For every two weeks, a low dose of 1/2 ppm should be maintained to prevent the development of resistant species of algae. TABLET is 90% available for this purpose. Results of TABLET use in the skimmer basket for 10,000 gallons of water will satisfy the requirement when used in accordance with the label. TABLET is a product of water by 1/2 ppm at 20°C (68°F) and by a suitable test kit.

DANGER: Avoid contact with eyes and skin. Avoid inhalation of dust. Avoid contact with clothing. Avoid contact with food and drink. Avoid contact with children.

DANGER: Avoid contact with eyes and skin. Avoid inhalation of dust. Avoid contact with clothing. Avoid contact with food and drink. Avoid contact with children.


Avoid contact with eyes and skin. Avoid inhalation of dust. Avoid contact with clothing. Avoid contact with food and drink. Avoid contact with children.

IF CONTACTED WITH EYES: Rinse eyes with large amounts of water. If contacted with skin: Wash with cold water. If contacted with clothing: Remove clothing and wash with cold water. If swallowed: Drink plenty of water. Do not induce vomiting. If inhaled: Move to fresh air. If symptoms persist, seek medical attention.


IF CONTACTED WITH EYES: Rinse eyes with large amounts of water. If contacted with skin: Wash with cold water. If contacted with clothing: Remove clothing and wash with cold water. If swallowed: Drink plenty of water. Do not induce vomiting. If inhaled: Move to fresh air. If symptoms persist, seek medical attention.

This product is not for use in swimming pools. It is for use in lakes, streams, ponds, and public wells.


© 1974 Haviland Products, Inc.




TABLETED DURACHLOR
EPA Reg. No. 10183-18



NET WT. 7 LBS.

HAVILAND  PRODUCTS



TABLETED DURACHLOR -- Slow dissolving tablets to provide continuous chlorination to control bacteria and algae.

DIRECTIONS

Initially pre-condition your pool water with ECONOCHLOR liquid or POWDERED ECONOCHLOR. Follow directions on the label of these products.

RECOMMENDED DOSAGE: For each 15,000 gals. of pool water, place 1 tablet in the skimmer basket. Allow 8 to 12 hrs. for chlorine residual to equalize. Initial treatment will last from 2 to 4 days depending on water temperature and flow rate through the skimmer. Chlorine residual should be maintained between 1.0 and 1.5 ppm as determined by a suitable test kit. If daily testing indicates the chlorine level to be low, increase the number of tablets in the skimmer basket. If daily testing indicates the chlorine level to be higher than 1.5 ppm, reduce the number of tablets accordingly.

Once every two weeks a shock dose of 5 ppm chlorine should be made to prevent the development of resistant species of algae. TABLETED DURACHLOR is not recommended for this purpose. Regular DURACHLOR used at the rate of 1/2 lb. per each 10,000 gallons of water will satisfy this requirement when used in conjunction with TABLETED DURACHLOR. Keep pH of water between 7.2 and 7.6 as determined by a suitable test kit.

DANGER: May be fatal or harmful if swallowed. Causes eye damage. May cause irritation of intact skin, nose and throat or burns on broken skin. Do not get in eyes, on skin or on clothing. Dust may be harmful if inhaled. Avoid breathing dust.

DANGER: Strong Oxidant. Moisture contamination may liberate irritating and hazardous chlorine containing gases. Decomposes at 350-475°F. with liberation of heat and harmful gases.

Avoid contact with oil, grease, other combustible matter. Store in cool, dry, well-ventilated area. Keep container tightly closed when not in use. Protect from moisture contamination. Keep away from open flames, heaters and other heat sources. Rinse empty container thoroughly with water and discard. Do not burn.

IN CASE OF PERSONAL CONTACT: Remove excess material and flush skin or eyes with cold water for at least 15 minutes. For eyes, get medical attention. Clean contaminated clothing before re-use. **If Swallowed:** Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available drink large quantities of water. Call a physician immediately.

IN CASE OF FIRE OR SUSTAINED DECOMPOSITION: If possible isolate container in open air or well-ventilated area. Flood with large volume of water.

This product is toxic to fish. Do not discharge treated effluent where it will drain into lakes, streams, ponds or public water.



EPA Reg. No. 10183-18

 **HAVILAND** PRODUCTS COMPANY
421 ANN ST. N.W. GRAND RAPIDS, MICH. 49504

ACCEPTED

FFR :

TABLETED DURACHLOR
EPA Reg. No. 10183-18



TABLETED DURACHLOR - Slow dissolving tablets to provide continuous chlorination to control bacteria and algae.

DIRECTIONS

Initially pre-condition your pool water with ECONOCHLOR liquid or POWDERED ECONOCHLOR. Follow directions on the label of these products.

RECOMMENDED DOSAGE: For each 15,000 gallons of pool water, place 1 tablet in the skimmer basket. Allow 8 to 12 hrs. for chlorine residual to stabilize. Initial treatment will last from 2 to 4 days depending on water temperature and filtration through the skimmer. Chlorine residual should be maintained between 1.0 and 1.5 ppm as determined by a suitable test kit. If daily testing indicates the chlorine level to be low, increase the number of tablets in the skimmer basket. If daily testing indicates the chlorine level to be higher than 1.5 ppm, reduce the number of tablets accordingly.

Once every two weeks a shock dose of 5 ppm chlorine should be made to prevent the development of resistant species of algae. TABLETED DURACHLOR is not recommended for this purpose. Regular DURACHLOR used at the rate of 1 1/2 lb. per each 10,000 gallons of water will satisfy this requirement when used in conjunction with TABLETED DURACHLOR. Keep pH of water between 7.2 and 7.6 as determined by a suitable test kit.

DANGER: May be fatal or harmful if swallowed. Causes eye damage. May cause irritation of intact skin, nose and throat or burns on broken skin. Do not get in eyes, on skin or on clothing. Dust may be harmful if inhaled. Avoid breathing dust.

DANGER: Strong Oxidant. Moisture contamination may liberate irritating and hazardous chlorine containing gases. Decomposes at 300-475°F. with liberation of heat and harmful gases.

Avoid contact with oil, grease, other combustible matter. Store in cool, dry, well-ventilated area. Keep container tightly closed when not in use. Protect from moisture contamination. Keep away from open flames, heaters and other heat sources. Rinse empty container thoroughly with water and discard. Do not burn.

IN CASE OF PERSONAL CONTACT: Remove excess material and flush skin or eyes with cold water for at least 15 minutes. For eyes, get medical attention. Clean contaminated clothing before re-use. If Swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available drink large quantities of water. Call a physician immediately.

IN CASE OF FIRE OR SUSTAINED DECOMPOSITION: If possible isolate container in open air or well-ventilated area. Flood with large volume of water.

This product is toxic to fish. Do not discharge treated effluent where it will drain into lakes, streams, ponds or public water.

EPA Reg No. 10183-18

