

Net Content

One Pound

ITKILLS

WHEN USED AS DIRECTED

RAT AND MOUSE KILLER

Follow Directions Carefully

ACTIVE INGREDIENTS: WARFARIN

(3-a-acetylbenzyl)-4-hydroxycoumarin) 0.025%

INERT INGREDIENTS:

99.975%

Total 100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See other precautions on back/side panel

Distributed by

Best-Way Exterminating Company

3081 Cropsey Ave., Brooklyn, N.Y. 11224

Telephone: ES 2-4742

USDA Reg. #10199-2

RATS: 4 oz. to 1 pound of mixed bait at 8-12 foot intervals in locations frequented by rats and protect from children, dogs, cats, and livestock by means of bait boxes or cages, where necessary. The minimum time for exposed bait should be 10 days. Baits should be replaced as consumed. Use no decomposed, moldy or sour baits.

Where a continuous source of infestation prevails from nearby dumps or fields, permanent bait stations should be used and baits replenished as needed.

MICE: For controlling mice follow the same general directions as for rats except bait replacements should be 1/2 oz. to 1 oz. and more placements should be made. Mice are more difficult to control than rats and complete control may take a longer period of baiting. The minimum time for exposed bait should be 15 days. Baits should be replaced as consumed. Use no decomposed, moldy or sour baits.

IMPORTANT NOTICE

BAITING SHOULD CONTINUE UNTIL COMPLETE LACK OF FEEDING IS NOTED . . . THIS SHOULD BE FROM 10 TO 15 DAYS

CAUTION: Contains as its active ingredient an anticoagulant chemical which if taken by human, domestic animals or pets may reduce the clotting ability of the blood and serious hemorrhage may result. In case baits are eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS: Vitamin K₁ has been known to be an antidote; blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdose of Dicumarol®.

10179-2