

ACCEPTED

ROCKWOOD BRAND

PERTHANE

4 - EC

CAUTION

KEEP OUT OF REACH OF CHILDREN

KEEP CONTAINER CLOSED

DESTROY CONTAINER WHEN EMPTY

COMPOSITION

ACTIVE INGREDIENTS:

Diethyl diphenyl dichlorethane 45.0%

Related Reaction products..... 2.3%

Aromatic petroleum derivative solvent..... 48.7%

INERT INGREDIENTS:..... 4.0%

TOTAL..... 100.0%

Harmful if swallowed, inhaled, or absorbed through skin.

This solution if brought into repeated or prolonged contact with skin may cause toxic symptoms. Avoid breathing spray mist. Avoid contamination of feed or foodstuffs. Avoid skin contact. In case of skin contact wash with soap and water. Inflammable. Do not use in dairy barns, on dairy animals, animals being finished for slaughter, or in feed or forage to be fed to these animals.

This product is toxic to fish. Keep out of lakes, streams, or ponds.

Apply this product only as specified on this label.

Do not re-use empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

DIRECTIONS

ROCKWOOD BRAND PERTHANE 4 - EC is compatible with other commonly used insecticides and fungicides, such as malathion, parathion, DDT, rhothane (TDE), methoxychlor, dithane, captan, ferbam, glyodin, sulfur.

Through, full-coverage of spray application is important and nozzles should be adjusted properly for uniform distribution.

LETTUCE: For control of leafhoppers, imported cabbage worm, cabbage looper, and alfalfa loopers, (California). Use 1 qt. per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial sprayers.

Apply when the insects are found to be present, making thorough application. Repeat at weekly to 10-day intervals or as required for adequate insect control. Do not apply within 4 days of harvest.

SPINACH: For control of leahopper, cabbage loopers. Use 1 qt. per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial application. Apply when insects are first noted, making thorough application. Repeat applications at weekly to 10-day intervals as required for adequate insect control. Do not apply within 7 days of harvest.

CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI: For control of cabbage looper, imported cabbage worms. Use 1 qt. per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial applications. Repeat when insects are first noted, making thorough applications at weekly to 10-day intervals as required for adequate insect control. Do not apply within 3 days of harvest.

ROCKWOOD BRAND

PERTHANE
4 - EC

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of ROCKWOOD CHEMICAL COMPANY, no guarantee expressed or implied is made as to the effects or such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

ROCKWOOD CHEMICAL COMPANY
P.O. BOX 34, BRAWLEY, CALIFORNIA

2/25/69 DJ/bu

ACCEPTED

MAR 24 1969

UNDER THE FEDERAL
INSECTICIDE ACT
1947
100-8

ROCKWOOD BRAND

PERTHANE

4 - EC

CAUTION

KEEP OUT OF REACH OF CHILDREN

KEEP CONTAINER CLOSED

DESTROY CONTAINER WHEN EMPTY

COMPOSITION

ACTIVE INGREDIENTS:

Diethyl diphenyl dichlorethane 45.0%

Related Reaction products..... 2.3%

Aromatic petroleum derivative solvent..... 48.7%

INERT INGREDIENTS:..... 4.0%

TOTAL..... 100.0%

Harmful if swallowed, inhaled, or absorbed through skin.

This solution if brought into repeated or prolonged contact with skin may cause toxic symptoms. Avoid breathing spray mist. Avoid contamination of feed or foodstuffs. Avoid skin contact. In case of skin contact wash with soap and water. Inflammable. Do not use in dairy barns, on dairy animals, animals being finished for slaughter, or in feed or forage to be fed to these animals.

This product is toxic to fish. Keep out of lakes, streams, or ponds.

Apply this product only as specified on this label.

Do not re-use empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

DIRECTIONS

ROCKWOOD BRAND PERTHANE 4 - EC is compatible with other commonly used insecticides and fungicides, such as malathion, parathion, DDT, rhothane (TDE), methoxychlor, dithane, captan, ferbam, glyodin, sulfur.

Through, full-coverage of spray application is important and nozzles should be adjusted properly for uniform distribution.

LETTUCE: For control of leafhoppers, imported cabbage worm, cabbage looper, and alfalfa loopers, (California). Use 1 qt. per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial sprayers. Apply when the insects are found to be present, making thorough application. Repeat at weekly to 10-day intervals or as required for adequate insect control. Do not apply within 4 days of harvest.

SPINACH: For control of leahopper, cabbage loopers. Use 1 qt. per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial application. Apply when insects are first noted, making thorough application. Repeat applications at weekly to 10-day intervals as required for adequate insect control. Do not apply within 7 days of harvest.

CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI: For control of cabbage looper, imported cabbage worms. Use 1 qt. per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial applications. Repeat when insects are first noted, making thorough applications at weekly to 10-day intervals as required for adequate insect control. Do not apply within 3 days of harvest.