

ACCEPTED
 JUL 25 1975
 U.S. Environmental Protection Agency
 Registration Division
 EPA Reg. No. 10226-29

ROCKWOOD CHEMICAL CO.

P. O. Box 34

Brawley, California

ROCKWOOD BRAND

BAITLOX 3%

Contains 3% Dylox*

DIRECTIONS FOR USE

Rockwood Brand BAITLOX 3% can be used on the following crops for the control of Crickets, Cutworms (both climbing and surface), Grasshoppers (in Cotton only). This material may be applied with either ground or air equipment. To be used only by thoroughly trained operators, familiar with the handling of hazardous insecticides. For brevity days between final treatment and harvest will be in parenthesis. ()

ACTIVE INGREDIENTS:

Dimethyl (2,2,2 - Trichloro-1-hydroxyethyl) phosphonate* 3.0%

INERT INGREDIENTS: 97.0%

Total 100.0%

*Dylox is Reg. TM of the parent company of Farben Fabriken Bayer, GmbH Leverkusen

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food. Product is a cholinesterase inhibitor.

NOTE TO PHYSICIAN: Atropine is antidote. If results in poisoning, obtain prompt medical aid.

NOTICE: Recommendations for the use of this product based upon information believed to be reliable at time of printing. The use of this product to control the control of ROCKWOOD CHEMICAL COMPANY, no warranty, expressed or implied is made as to the effects of such use. The results to be obtained if not used in accordance with directions or established safe and sound practices. THE BUYER, must assume all responsibility including injury and/or damage, resulting from its misuse, as such or in combination with other products.

ENVIRONMENTAL CAUTIONS: This product is toxic to birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from areas treated.

BEI CAUTION: This product is toxic to bees and should not be applied when bees are actively visiting areas.

DO NOT REUSE EMPTY CONTAINER. STAY AWAY FROM SMOKE, OR FUMES. DESTROY IT BY BURYING WITH WASTE OR PULPING.

EPA - Reg. No. 10226-29
 EPA - Est. 10226-CA-1

NET CONTENTS _____

USE	DOSAGE	LIMITATIONS
Alfalfa	33 lbs. / Acre	14 days of cutting or grazing. One application per cutting.
Alfalfa (seed crop)	33 lbs. / Acre	14 days of cutting or grazing. One application per cutting.
Barley	33 lbs. / Acre	21 days. Do not graze treated fields.
Oats	33 lbs. / Acre	21 days. Do not graze treated fields.
Wheat	33 lbs. / Acre	21 days. Do not graze treated fields.
Cabbage	33 lbs. / Acre	21 days.
Cauliflower	33 lbs. / Acre	21 days.
Tomatoes	33 lbs. / Acre	28 days. Do not graze treated fields. Do not apply after plants are two feet tall.
Corn	33 lbs. / Acre	Apply once per season when plants are 3-12 inches high. Do not use corn for food, feed or ensilage within 40 days of treatment.
Cotton	50 lbs. / Acre	Do not apply within 7 days of picking. Do not graze livestock in treated fields within 14 days after application.
Lettuce	33 lbs. / Acre	28 days.
Sugar Beets	50 lbs. / Acre	28 days if tops are to be fed to livestock. 14 days if tops are not to be fed.