

# ROCKWOOD CHEMICAL CO.

P.O. Box 34

Brawley, California

ROCKWOOD BRAND

## Bran-L-Bait 1.5%

### DIRECTIONS FOR USE

Rockwood Brand Bran-L-Bait 1.5% bran bait may be applied with either ground or air equipment. To be used only by thoroughly trained operators, familiar with handling of hazardous insecticides. For brevity,

May be fatal if swallowed, inhaled, or absorbed through skin. In case of contact, wash skin with plenty of soap and water; for eyes flush with water for 15 minutes. Get medical attention. Remove contaminated clothing and wash before reuse.

If swallowed, give a tablespoonful of salt in a glass of water and repeat until vomit is clear. If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. NEVER give anything by mouth to an unconscious person. Wear mask or respirator of a type recommended by the U. S. Bureau of Mines to give adequate protection against this material.

### ANTIDOTE

Atropine is an antidote. Call a physician at once if poisoning is suspected. Do not use morphine or 2-PAM.

### NOTE TO PHYSICIAN

Warning symptoms: methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in chest, constriction in pupils, sweating, slow pulse, muscle tremors.

### RECOMMENDED APPLICATIONS

SIZE & PEST	DOSAGE	LIMITATIONS
<b>TOMATOES:</b>	1 day pre-harvest interval through 3016/A. Foliage application. 2 day pre-harvest interval from above 30 through 60 lbs./A.	
Beet Armyworm	30-60 lbs./A	
Cabbage Looper	30-60 lbs./A	
Tomato Fruitworm	30-60 lbs./A	For tomato fruitworm apply at first fruit set.
<b>COTTON:</b>	40 day pre-harvest interval through 44.5 lbs./A. Rockwood Brand Bran-L-Bait 1.5% may redden cotton foliage. Alternate with other insecticides or discontinue the use of methomyl if reddening is excessive. Do not graze or feed treated plants to livestock.	
Bollworm	30-44.5 lbs./A	Use restricted to areas west of the Rocky Mountains. Begin treatment when an economic population develops (normally 5-15 small caterpillars per 100 terminals).

(continued on side panel)

### ACTIVE INGREDIENT:

Methomyl  
S-methyl-N- (Methylcarbamoyl)  
oxythioacetimidate . . . . . 1.50%

INERT INGREDIENTS . . . . . 98.50%

Total 100.00%

Contains Lannate a Reg. U.S. Pat. Office  
U.S. Patent Pending

**WARNING**

KEEP OUT OF REACH OF CHILDREN

**NOT FOR USE**

**OR STORAGE**

**AROUND THE HOME**

Burn, bury or otherwise dispose of containers and unused pesticides in accordance with local laws and regulations.

This product is toxic to fish, birds, and other wildlife. Keep off of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

See antidote statement and other precautions on the back/side panel.

E.P.A. Reg. No. 10226-30  
E.P.A. Est. No. 10226-CA-1

NET CONTENTS \_\_\_\_\_

C R Printers - P6714 Rev. 7 2 74

SITE & PEST	DOSAGE	LIMITATIONS
<b>ALFALFA:</b>	7 day pre-harvest interval through 60 lbs./A. Do not graze or feed treated alfalfa to livestock within 7 days of last application. In fields visited by honey bees, make application in late evening or early morning. In areas frequented by alkali or leaf-cutting bees, do not apply during bloom or when bees will be present in fields.	
Beet Armyworm	14.6-30 lbs./A	Use restricted to California and Arizona. Apply when insects appear in damaging numbers and repeat as needed.
Beet Armyworm & Egyptian Alfalfa Weevil (larvae)	30-60 lbs./A	Apply when insects appear in damaging numbers and repeat as needed.
Alfalfa Caterpillar	20-30 lbs./A	Use for California and Arizona only. Apply when insects appear in damaging numbers and repeat as needed.
Alfalfa Weevil (larvae)	60 lbs./A.	Apply when 50% or more of tips show feeding injury. Repeat as needed. Treat stubble as soon as crop is removed if infestation is severe at harvest.
<b>BROCCOLI:</b>	7 day pre-harvest interval through 30 lbs./A. 14 day pre-harvest interval from 30-60 lbs./A.	
Cabbage Looper	14.6-60 lbs./A	
<b>CABBAGE:</b>	1 day pre-harvest interval through 60 lbs./A.	
Cabbage Looper	15-60 lbs./A	
<b>CUCUMBERS:</b>	3 day pre-harvest interval through 60 lbs./A. Rockwood Brand Bran-L-Bait is toxic to bees and should not be applied when bees are actively visiting area. For ground application only.	
Cabbage Looper	30-60 lbs./A	

**NOTICE:** Recommendations for the use of this product are based upon information believed to be reliable at the time this product being beyond the control of ROCKWOOD CHEMICAL COMPANY, no guarantee, expressed or implied, of such or the results to be obtained if not used in accordance with directions or established safe and sound practices. Assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other

SITE & PEST	DOSAGE
<b>LETTUCE: (Head)</b>	7 day pre-harvest interval. Foliage application. above 30 through 60
Alfalfa Looper	14.6-60 lbs./A
Beet Armyworm	14.6-60 lbs./A
Cabbage Looper	33.3-60 lbs./A
<b>MELONS:</b>	3 day pre-harvest interval. Brand Bran-L-Bait is applied when bees are on ground application only.
Cabbage Looper	30-60 lbs./A
<b>SUGAR BEETS:</b>	7 day pre-harvest interval. Foliage application.
Beet Armyworm	14.6-60 lbs./A
Beet Webworm	14.6-60 lbs./A
Flea Beetles	14.6-60 lbs./A

# ROCKWOOD CHEMICAL CO.

Brawley, California

ROCKWOOD BRAND

Bran-L-Bait 1.5%

## ACTIVE INGREDIENT:

Methomyl  
S-methyl-N- (Methylcarbamoyl)  
oxy- thioacetimidate . . . . . 1.50%

INERT INGREDIENTS . . . . . 98.50%

Total 100.00%

Contains Lannate a Reg. U.S. Pat. Office  
U.S. Patent Pending

**WARNING**

KEEP OUT OF REACH OF CHILDREN

**NOT FOR USE**

**OR STORAGE**

**AROUND THE HOME**

Burn, bury or otherwise dispose of containers and unused pesticides in accordance with local laws and regulations.

This product is toxic to fish, birds, and other wildlife. Keep away from lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

See antidote statement and other precautions on the back/side panel.

E.P.A. Reg. No. 10226-30  
E.P.A. Est. No. 10226-CA-1

NET CONTENTS \_\_\_\_\_

C R Printers - PG 714 Rev. 7 2 74

SITE & PEST	DOSAGE	LIMITATIONS
ALFALFA:	7 day pre-harvest interval through 60 lbs./A. Do not graze or feed treated alfalfa to livestock within 7 days of last application. In fields visited by honey bees, make application in late evening or early morning. In areas frequented by alkali or leaf-cutting bees, do not apply during bloom or when bees will be present in fields.	
Beet Armyworm	14.6-30 lbs./A	Use restricted to California and Arizona. Apply when insects appear in damaging numbers and repeat as needed.
Beet Armyworm & Egyptian Alfalfa Weevil (larvae)	30-60 lbs./A	Apply when insects appear in damaging numbers and repeat as needed.
Alfalfa Caterpillar	20-30 lbs./A	Use for California and Arizona only. Apply when insects appear in damaging numbers and repeat as needed.
Alfalfa Weevil (larvae)	60 lbs./A.	Apply when 50% or more of tips show feeding injury. Repeat as needed. Treat stubble as soon as crop is removed if infestation is severe at harvest.
BROCCOLI:	7 day pre-harvest interval through 30 lbs./A. 14 day pre-harvest interval from 30-60 lbs./A.	
Cabbage Looper	14.6-60 lbs./A	
CABBAGE:	1 day pre-harvest interval through 60 lbs./A.	
Cabbage Looper	15-60 lbs./A	
CUCUMBERS:	3 day pre-harvest interval through 60 lbs./A. Rockwood Brand Bran-L-Bait is toxic to bees and should not be applied when bees are actively visiting area. For ground application only.	
Cabbage Looper	30-60 lbs./A	

SITE & PEST	DOSAGE	LIMITATIONS
LETTUCE: (Head)	7 day pre-harvest interval through 30 lbs./A. Foliage application. 10 day pre-harvest interval from above 30 through 60 lbs./A.	
Alfalfa Looper	14.6-60 lbs./A	
Beet Armyworm	14.6-60 lbs./A	
Cabbage Looper	33.3-60 lbs./A	
MELONS:	3 day pre-harvest interval through 60 lbs./A. Rockwood Brand Bran-L-Bait is toxic to bees and should not be applied when bees are actively visiting the area. For ground application only.	
Cabbage Looper	30-60 lbs./A	
SUGAR BEETS:	7 day pre-harvest interval through 60 lbs./A. Foliage application. Do not feed tops to livestock.	
Beet Armyworm	14.6-60 lbs./A	Use restricted to Arizona and California. Apply when insects or their damage appears. Make up to 2-3 applications as needed.
Beet Webworm	14.6-60 lbs./A	Use restricted to Arizona and California. Apply when insects or their damage appears. Make up to 2-3 applications as needed.
Flea Beetles	14.6-60 lbs./A	Use restricted to Arizona and California. Apply when insects or their damage appears. Make up to 2-3 applications as needed.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of ROCKWOOD CHEMICAL COMPANY, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.