# ROCKWOOD CHEMICAL CO.

P. O. Box 34

ACCEPTED

MAR 2 1 1975

CAUTIO MUNGICIDE AND RODENTICIDE ACT

KEEP OUT OF REACH OF CHILDREN 10 10 2 1 - 22

Harmful if swallowed. Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using. Do not store near heat or open flame. Use with adequate ventilation. keep container closed. Avoid contamination of feed and foodstuffs.

CAUTION COMBUSTIBLE LIQUID

DO NOT USE, POUR, SPILL OR STORE

NEAR HEAT OR OPEN FLAME.

PROTECT FROM TEMPERATURES BELOW 20° F.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of ROCKWOOD CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER, must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

FISH AND WILDLIFE CAUTIONS: This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**ROCKWOOD BRAND** 

# Terraclor. 2 Lb. E.C.

An Emulsion Concentrate Containing 2 Pounds of P.C.N.B. Per Gallon

#### **Active Ingredients**

Inert Ingredients

73.51%

Total..... 100.00%

® Terraclor - Product of Olin Corporatin

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

**CONVERSION FACTOR:** 

40 inch beds - 13,068 linear feet of row per acre.

CONTAINER DISPOSAL: Do not reuse container. Destroy when empty.

See other precautions on the side panel.

#### DIRECTIONS FOR USE

Brawley, California

Mix ROCKWOOD BRAND Terraclor<sup>®</sup> 2 lb. E.C. with water by agitation; agitate the mixture white spraying. For maximum control apply with uniform coverage.

COTTON: For control of damping-off (Rhizoctonia) apply ½ to 1 gal. per 13,000 linear feet of row based on a 40<sup>th</sup> bed at planting time. Do not allow the feeding or grazing of cotton foilage by livestock. Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil. Apply in furrow over seed and surrounding soil at planting time.

Apply the recommended dosgaes in 20-50 gallons of water by ground.

GARLIC (Whiterot - Sclerotium cepivorum) Use 10 gals. mixed in 100 gals, of water and apply to 21,800 linear feet of row at planting time. Attach a suitable rig to the machine planter, using 2 nozzles per planting furrow. Direct the front nozzle to spray the bottom of the open furrow and the "seed" as it is dropped. Direct the rear nozzle to spray the covering soil. Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

**NET CONTENTS** 

EPA. Reg. No. 10226 - 32 EPA. EST 10226 CA 1