

ROCKWOOD CHEMICAL CO.

P.O. Box 34

ROCKWOOD BRAND

Brawley, California

DIRECTIONS FOR USE

CAUTION May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Do not contaminate food or feed. Keep out of reach of children. If misuse results in poisoning, obtain prompt medical aid. Dylox may cause cholinesterase inhibition.

Do NOT APPLY Rockwood Roclox 5% Dust to, or allow dust to drift onto, varieties of sorghum which are sensitive to phosphates. This dust may cause burning of the sensitive sorghum. Do not use on other crops used for food or feed.

NOTE: This product may leave a poisonous residue on plants to which it is applied. The U.S. Food and Drug Administration has set a limit of a maximum amount of such poisonous residue that may remain on raw agricultural products at harvest, and it is the user's responsibility to see that there is no residue on crops at harvest in excess of this maximum amount.

BEE CAUTION

This product is toxic to bees and should not be applied when bees are actively visiting the area.

NOTICE: This product is a restricted use pesticide. It may be sold only to persons who are registered with the U.S. Environmental Protection Agency. The use of this product is limited to the county of ROCKWOOD CHEMICAL COMPANY, its quarantined agents, and employees of the company. Use of this product is restricted to use in accordance with the label and applicable laws and sound practice. THE USER is responsible for all liability, including the cost of removal, cleanup, and disposal, which may result from the use of this product.

Net Contents

EPA Est. 10226-CA-1

EPA Reg. No. 10226-36

ROCLOX 5% DUST

Contains 5% [®] Dylox

ACTIVE INGREDIENT:

Dimethyl (2, 2, 2-trichloro-1-hydroxyethyl) phosphonate 5%

INERT INGREDIENTS 95%
100%

© Dylox is Reg. TM of the parent company of Farben-fabriken Bayer GmbH, Leverkusen.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Do not apply to, or allow to drift onto, varieties of Sorghum which are sensitive to phosphates.

DISPOSAL: Destroy bags and waste by burying in non-cropland areas (away from water) or by burning.

Note to Physician: Atropine Sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

Wildlife Caution: This product is toxic to birds and wildlife. Keep out of lake, stream, or pond. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only to the extent directed on the label.

RECOMMENDED APPLICATIONS

For Brevity, days between final treatment and harvest will be in parentheses.

ALFALFA: For control of Alfalfa Caterpillar, Alfalfa Webworm, Western Yellowstriped armyworm, Beet armyworm, Variegated cutworm, Lygus bug, Stink bug and Tarnished plant bug apply 20 lbs. per acre. (7 days between treatment and cutting or grazing. One application per cutting).

COTTON: For control of Cotton fleahopper, Cotton leafworm, Darkling ground beetle, Western yellow striped armyworm, Beet armyworm, Southern garden leafhopper, Black fleahopper complex, Cotton leaf perforator, leaf roller, Lygus bug, Stink bug and Saltmarsh caterpillar apply 30 lbs. per acre. (Do not apply within 7 days of picking. Do not graze livestock in treated fields within 14 days after application).

SAFFLOWER: For control of Armyworm, Dipterous leaf miner, Imported cabbageworm, Lygus bug, Stink bug, Thrip and Variegated cutworm apply 30 lbs. per acre. Do not apply after flowering stage. Do not apply more than three times a year.

SUGAR BEETS: For control of Beet webworm, Variegated cutworm, Dipterous leaf miners, Alfalfa caterpillar, Beet armyworm and Saltmarsh caterpillar apply 30 lbs. per acre. (14 days between treatments).

BEANS (Dry, Snap): For control of Western yellow striped armyworm, Imported cabbageworm, Imported leaf miner, Lygus bug, Mexican leaf beetle, Stink bug, Variegated cutworm apply 30 lbs. per acre. (14 days between treatments. Do not apply more than three times a year).

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: For control of Western yellow striped armyworm, Imported cabbageworm, Variegated cutworm and Dipterous leaf miner apply 30 lbs. per acre. (14 days between treatments).

CARROTS: For control of western yellow striped armyworm, Imported cabbageworm, Variegated cutworm, Lygus bug, Stink bug, Thrip, and Variegated cutworm apply 30 lbs. per acre. (14 days between treatments. Do not apply more than three times a year).

PEPPERS: For control of Beet webworm, Variegated cutworm, Dipterous leaf miner, and Lygus bug apply 30 lbs. per acre. (14 days between treatments. Do not apply more than three times a year).

TOMATOES: For control of Imported cabbageworm, Lygus bug, Stink bug, Thrip, and Variegated cutworm apply 30 lbs. per acre. (14 days between treatments. Do not apply more than three times a year).

COLLARDS and LETTUCE: For control of Imported cabbageworm, Lygus bug, Stink bug, Thrip, and Variegated cutworm apply 30 lbs. per acre. (14 days between treatments. Do not apply more than three times a year).