

ROCKWOOD CHEMICAL CO.

P.O. Box 34

Brawley, California

ROCKWOOD BRAND

Jan 13, 1975

EPA Reg. No. 10226-38

EPA Est. 10226 - CA - 1

10226-38

ZINEB DUST No. 6

DIRECTIONS FOR USE

Rockwood Brand ZINEB DUST No. 6 should be applied to dry foliage at time when the air is still and the temperature does not exceed 90 degrees F. Thorough and uniform coverage is absolutely essential for disease control. Begin application when disease is first present and continue as long as disease conditions prevail. Repeat at 7 to 10 days intervals, more frequently under severe disease conditions.

Recommended days to harvest are in parenthesis ().

RECOMMENDED USES

CANTALOUPEs, CUCUMBERS, WATERMELONS
For the control of Downy mildew, Alternaria, anthracnose, and gummy stem blight apply 10-50 lbs. per acre. (5)

CABBAGE, BROCCOLI, CAULIFLOWER - For control of Downy mildew and alternaria apply 10-50 lbs. per acre. (7)

CARROTS - For control of early and late blights, use 10-50 lbs. per acre. Do not apply within 7 days of harvest if tops are to be used for food or feed.

LETTUCE - For control of Downy mildew, Botrytis slime, apply 10-50 lbs. per acre. (10)

ONIONS - For control of Downy mildew, Botrytis leaf blight and purple blotch apply 10-50 lbs. per acre. Do not apply within 7 days of harvest on green onions.

SQUASH (summer and winter): For control of Alternaria blight, blossom blight, downy mildew, gummy stem blight, apply 10-50 lbs. per acre. Begin when plants emerge or when disease first occurs. Repeat at 7 to 10 days intervals or as necessary. (5)

ACTIVE INGREDIENTS:

Zinc ethylene bisdithiocarbamate (zineb) 6%

INERT INGREDIENTS 94%

Total 100%

*Registered trade name of Rohm and Haas Company

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause irritation of nose, throat, and skin. Do not breathe dust. In case of prolonged or frequent use, wear clean protective clothing and bathe after each exposure. Avoid contamination of feed and foodstuffs.

Burn, bury or otherwise dispose of containers and unused pesticides in accordance with local laws and regulations.

RECOMMENDED USES

SPINACH - For control of Downy mildew, and White rust apply 10-50 lbs. per acre. (10)

TOMATOES - For control of Early and Late blight, Anthracnose, Stemphylium, Septoria leaf spot and Cladosporium apply 10-50 lbs. per acre. (5)

NOTE

This product may leave a poisonous residue on plants to which it is applied. The U. S. Food and Drug Administration has established a minimum amount of such poisonous residue that may remain on raw agricultural products at harvest and it is the user's responsibility to see that there is no residue on such crops at harvest in excess of this minimal amount.

REMOVE EXCESSIVE RESIDUE BY
WASHING, BRUSHING OR OTHER
EFFECTIVE MEANS.

NET CONTENTS

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of ROCKWOOD CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER, must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

"Do not contaminate water by cleaning of equipment, or disposal of wastes, and apply this product only as specified on this label."

C/R Printers - P4212 - 11874